

# Constructing Small-Bias Sets from Algebraic-Geometric Codes

Avraham Ben-Aroya  
The Blavatnik School of Computer Science  
Tel-Aviv University  
Israel 69978. \*

Amnon Ta-Shma  
The Blavatnik School of Computer Science  
Tel-Aviv University  
Israel 69978. †

## Abstract

We give an explicit construction of an  $\epsilon$ -biased set over  $k$  bits of size  $O\left(\frac{k}{\epsilon^2 \log(1/\epsilon)}\right)^{5/4}$ . This improves upon previous explicit constructions when  $\epsilon$  is roughly (ignoring logarithmic factors) in the range  $[k^{-1.5}, k^{-0.5}]$ . The construction builds on an algebraic-geometric code. However, unlike previous constructions we use low-degree divisors whose degree is significantly smaller than the genus.

Studying the limits of our technique, we arrive at a hypothesis that if true implies the existence of  $\epsilon$ -biased sets with parameters nearly matching the lower bound, and in particular giving binary error correcting codes beating the Gilbert-Varshamov bound.

## 1 Introduction

Explicitly constructing combinatorial objects with certain properties (such as expander graphs, extractors, error correcting codes and others) is an intriguing challenge in computer science. Often, it is easy to verify that a random object satisfies the required property with high probability, while it is difficult to pin down such an explicit object.

In most cases it is believed (and sometimes proven) that a random object is nearly optimal. Therefore, giving an optimal explicit construction becomes a derandomization problem. There are, however, rare cases in which explicit constructions outperform naive random constructions. Perhaps the most remarkable example of this type is that of Algebraic-Geometric codes (AG codes). In the seminal work of Tsfasman et al. [8] it was shown that there are Algebraic-Geometric codes over constant size alphabets that lie above the Gilbert-Varshamov bound, a bound that was believed to be optimal at the time.

---

\*Email: abraham@tau.ac.il. Supported by the Adams Fellowship Program of the Israel Academy of Sciences and Humanities, and by the European Commission under the Integrated Project QAP funded by the IST directorate as Contract Number 015848.

†Email: amnon@tau.ac.il. Supported by Israel Science Foundation grant 217/05 and by USA Israel BSF grant 2004390.

The important case of *binary* error correcting codes is still open. The Gilbert-Varshamov bound gives the best known (explicit or non-explicit) codes to date. Finding an explicit construction that attains this bound is an open problem as well. The above statements also apply if we restrict ourselves to codes with distance close to half, which is a case of special interest.

Another closely related question is that of finding an  $[n, k, \frac{1}{2} - \epsilon]_2$  binary code, in which the relative weight of every non-zero codeword is in the range  $[\frac{1}{2} - \epsilon, \frac{1}{2} + \epsilon]$ . Such codes are called  $\epsilon$ -balanced and they are related to another kind of combinatorial objects called  $\epsilon$ -biased sets. An  $\epsilon$ -biased set is a set  $S \subseteq \{0, 1\}^k$  such that for every non-empty subset  $T \subseteq [k]$ , the binary random variable  $\bigoplus_{i \in T} s_i$ , where  $s$  is sampled uniformly from  $S$ , has bias at most  $\epsilon$ . It turns out that  $\epsilon$ -biased sets are just  $\epsilon$ -balanced codes in a different guise: the columns of a matrix whose rows generate an  $\epsilon$ -balanced code form an  $\epsilon$ -biased set, and vice versa. In terms of parameters, an  $[n, k]_2$   $\epsilon$ -balanced code is equivalent to an  $\epsilon$ -biased set  $S \subseteq \{0, 1\}^k$  of size  $n$ .

The status of  $\epsilon$ -balanced codes is similar to that of  $[n, k, \frac{1}{2} - \epsilon]_2$  codes. In both cases the probabilistic method gives non-explicit  $[n, k]_2$   $\epsilon$ -balanced codes with  $n = O(\frac{k}{\epsilon^2})$ , whereas the best lower bound is  $n = \Omega(\frac{k}{\epsilon^2 \log(\frac{1}{\epsilon})})$ . For a discussion of these bounds see [1, Section 7].

There are several *explicit* constructions of such codes. Naor and Naor [5] give a construction with  $n = k \cdot \text{poly}(\epsilon^{-1})$ . Alon et al. [1] have the incomparable bound  $n = O(\frac{k^2}{\epsilon^2 \log^2(k/\epsilon)})$ . Concatenating Algebraic-Geometric codes with the Hadamard code gives  $n = O(\frac{k}{\epsilon^3 \log(\frac{1}{\epsilon})})$ . In this paper we show an explicit construction of an  $[n, k]_2$   $\epsilon$ -balanced code with  $n = O(\frac{k}{\epsilon^2 \log(\frac{1}{\epsilon})})^{5/4}$ , which improves upon previous explicit constructions when  $\epsilon$  is roughly (ignoring logarithmic factors) in the range of  $k^{-1.5} \leq \epsilon \leq k^{-0.5}$  (see Figure 1).

The construction is simple and can be described by elementary means. We first take a finite field  $\mathbb{F}_q$  of the appropriate size. We then carefully choose a subset  $A$  of  $\mathbb{F}_q \times \mathbb{F}_q$ . The elements in the  $\epsilon$ -biased set are indexed by pairs  $((a, b), c) \in A \times \mathbb{F}_q$ . For each  $((a, b), c) \in A \times \mathbb{F}_q$  the corresponding element is the bit vector  $((a^i b^j), c)_2)_{i,j}$ , where  $(i, j)$  range over all integers  $i, j$  whose sum is bounded by an appropriately chosen parameter and the inner product is of the binary representation of the elements in  $\mathbb{F}_q$ . The analysis of the construction relies on Bézout's Theorem.

To put the construction in context, we need to move to algebraic function fields terminology. AG codes are *evaluation* codes where a certain set of *evaluation functions* is evaluated at a chosen set of *evaluation points*. The space of evaluation functions used is a vector space (this is the reason we get a linear error correcting code) and is determined by a *divisor*  $G$ . We explain what a divisor is and other terminology in Section 3, and for the time being continue with an intuitive discussion. We denote the code associated with a divisor  $G$  by  $C(G)$ .

The code  $C(G)$  has the following parameters. The *length* of the code is the number of evaluation points and is denoted by  $N = N(F)$  ( $F$  is the algebraic function field). The distance of the code is  $N - \deg(G)$  ( $\deg(G)$  is the degree of  $G$ , we explain what it is in Section 3). The dimension of the code,  $\dim(G)$ , is the dimension of the vector space of evaluation functions. When the "degree" of  $G$  is larger than the *genus* (we explain what the genus is in Section 3), the Riemann-Roch Theorem [6, Thm I.5.17] tells us exactly what the dimension  $\dim(G)$  is, and it turns out to be  $\deg(G) - g + 1$ . This almost matches the Singleton bound, except for a loss of  $1g$  in the dimension. Thus, our goal is to get as many evaluation points while keeping the genus small. Indeed, a lot of research was done on the best possible ratio between the




Figure 1: Constructions of  $\epsilon$ -biased sets for  $\epsilon = k^{-c}$

length of the code  $N(F)$  and the genus. The bottom line of this research, roughly speaking, is that  $N(F)$  can be larger than the genus by at most a multiplicative  $\sqrt{q} - 1$  factor and this is essentially optimal.

A simple check shows that when  $\deg(G)$  is larger than the genus, an AG code concatenated with Hadamard cannot give  $\epsilon$ -balanced codes with  $n$  better than  $O(\frac{k}{\epsilon^3 \log(\frac{1}{\epsilon})})$ . In contrast, our construction takes as an outer code an AG code  $C(G)$  where  $\deg(G)$  is much smaller than the genus, and we show that this leads to a better code. One explanation as to why our improvement was not found before is that previous research often focused on AG codes  $C(G)$  where  $\deg(G)$  is higher than the genus.

A natural question is whether the  $\epsilon$ -balanced codes we achieve are the best binary codes one can achieve using this approach. We do not know the answer to this question. When  $\deg(G)$  is smaller than the genus, one cannot use the Riemann-Roch Theorem, and estimating  $\deg(G)$  is often a challenging task. Furthermore,  $\dim(G)$  now depends on  $G$  itself, and not just on its degree as before. However, we can formulate the question as follows. The important thing to us is not the best possible ratio between the number of rational points  $N(F)$  and the genus. Instead, we are interested in the best possible ratio between  $N(F)$  and  $\deg(G)$ , where  $G$  is a *low-degree* divisor having a *large dimension*.

We show that such a good ratio implies good  $\epsilon$ -balanced codes. Using the Hermitian function field we give the construction with  $n = O(\frac{k}{\epsilon^2 \log(\frac{1}{\epsilon})})^{5/4}$  mentioned above. Furthermore, assuming an  $\Omega(q)$ -ratio we show we can construct binary  $\epsilon$ -balanced codes with  $n = O(\frac{k}{\epsilon^2 \log(\frac{1}{\epsilon})})$ , i.e., matching the known lower bound and outperforming the Gilbert-Varshamov bound. If this is the case, then AG codes would outperform naive random codes even over the binary alphabet. We mention that a simple argument due to Henning Stichtenoth [7] shows the

ratio can never be bigger than  $q + 1$  (we repeat his argument in Section 4).<sup>1</sup>

We hope our paper would lead researchers to study not only the possible gap between the genus and the number of rational points, but also the possible gap between high dimension divisors and the number of rational points – a problem that has been somewhat neglected so far.

The rest of the paper is organized as follows. In Section 2 we describe the construction and its analysis using Bézout’s Theorem. Section 3 contains a description the same construction in algebraic function fields terminology. In Subsections 3.1 and 3.1.1 we give the necessary background on algebraic function fields and geometric Goppa codes. Finally, in Section 4 we describe a possible hypothesis regarding function fields and its implications to  $\epsilon$ -balanced codes.

## 2 A self-contained elementary description of the construction

We first recall the definition of an  $\epsilon$ -biased set:

**Definition 1.** A set  $S \subseteq \{0, 1\}^k$  is  $\epsilon$ -biased if for every nonempty  $T \subseteq [k]$ ,

$$\frac{1}{|S|} \left| \sum_{s \in S} (-1)^{\sum_{i \in T} s_i} \right| \leq \epsilon.$$

**The construction:** Given  $k$  and  $\epsilon$ , let  $p = 2^\ell$  be a power of 2 in the range  $\left[ \left(\frac{k}{\epsilon^2}\right)^{1/4}, 2 \left(\frac{k}{\epsilon^2}\right)^{1/4} \right]$ .

Define  $q = p^2$  and  $r = \epsilon p^3$ . Let  $\mathbb{F}_q$  denote the finite field with  $q$  elements and  $\mathbb{F}_p$  its subfield with  $p$  elements. Consider the vector space of bivariate polynomials over  $\mathbb{F}_q$  with total degree at most  $r/(p+1)$ :

$$V = \left\{ \phi \in \mathbb{F}_q[x, y] : \deg(\phi) \leq \frac{r}{p+1} \right\} = \text{Span} \left\{ x^i y^j : i + j \leq \frac{r}{p+1} \right\}.$$

The dimension of this space (over  $\mathbb{F}_q$ ) is  $k' = \Omega\left(\frac{r^2}{p^2}\right) = \Omega(k)$ .

Let  $A \subseteq \mathbb{F}_q \times \mathbb{F}_q$  be the set of roots of the polynomial  $y^p + y - x^{p+1}$ . The  $\epsilon$ -biased set over  $k'$  bits that we construct is

$$S = \left\{ \left( \langle \text{bin}(a^i b^j), \text{bin}(c) \rangle_2 \right)_{i+j \leq \frac{r}{p+1}} : (a, b) \in A \text{ and } c \in \mathbb{F}_q \right\},$$

where  $\text{bin} : \mathbb{F}_q \rightarrow \mathbb{Z}_2^{2\ell}$  is any isomorphism between the additive group of  $\mathbb{F}_q$  and the vector space  $\mathbb{Z}_2^{2\ell}$  and  $\langle \cdot, \cdot \rangle_2$  denotes inner product over  $\mathbb{Z}_2^{2\ell}$ .

**The analysis:** The following claim will be used to bound the size of  $S$ .

**Claim 1.** The cardinality of  $A$  is  $p^3$ .

---

<sup>1</sup>We also mention that an  $\Omega(\sqrt{q})$ -ratio follows by using function fields where  $N(F)$  is larger than the genus by that factor.

**Proof:** The trace function  $\text{Tr}(y) = y^p + y$  maps  $\mathbb{F}_q$  to  $\mathbb{F}_p$ . We claim that for every  $\alpha \in \mathbb{F}_p$ , the number of solutions in  $\mathbb{F}_q$  to  $\text{Tr}(y) = \alpha$  is  $p$ . To see this, observe that  $\text{Tr}$  is a linear function. Hence, the set of solutions to  $\text{Tr}(y) = 0$  is a subgroup of  $\mathbb{F}_q$  that has at most  $p$  elements. For every  $\alpha \in \mathbb{F}_p$ , the set of solutions to  $\text{Tr}(y) = \alpha$  is either empty or a coset of this subgroup. As every element of  $\mathbb{F}_q$  is in one of these cosets, it must be the case that for every  $\alpha \in \mathbb{F}_p$  there are exactly  $p$  solutions.

The norm function  $N(x) = x^{p+1}$  also maps  $\mathbb{F}_q$  to  $\mathbb{F}_p$ . Thus, for every  $\alpha \in \mathbb{F}_q$  there are exactly  $p$  values  $\beta \in \mathbb{F}_q$  such that  $\text{Tr}(\beta) = N(\alpha)$ . Therefore,  $|A| = p^3$ . ■

We want to apply Bézout's Theorem on the bivariate polynomial  $y^p + y - x^{p+1}$ . However, we first need to show it is irreducible. We need Eisenstein's Criterion for irreducibility:

**Theorem 2** (Eisenstein's Criterion [4, Thm 3.1]). *Let  $U$  be a unique factorization ring and let  $K$  be its field of fractions. Let  $f(x) = \sum_{i=0}^n a_i x^i$  be a polynomial of degree  $n \geq 1$  in  $U[x]$ . Let  $\rho$  be a prime of  $U$ , and assume:*

- $a_n \not\equiv 0 \pmod{\rho}$
- For every  $i < n$ ,  $a_i \equiv 0 \pmod{\rho}$
- $a_0 \not\equiv 0 \pmod{\rho^2}$ .

*Then  $f(x)$  is irreducible in  $K[x]$ .*

With that we conclude:

**Claim 3.** *The polynomial  $y^p + y - x^{p+1}$  is irreducible over  $\mathbb{F}_q$ .*

**Proof:** This follows from Eisenstein's Criterion. The unique factorization ring we consider is  $U = \mathbb{F}_q[y]$ . The prime element we use is  $\rho = y$ . The leading coefficient is  $-1$  and  $-1 \not\equiv 0 \pmod{y}$ . Every other coefficient except the last is 0, hence it is  $0 \pmod{y}$ . The last coefficient is also  $0 \pmod{y}$ . Finally, since  $p \geq 2$ ,  $y^p = 0 \pmod{y^2}$  but  $y \not\equiv 0 \pmod{y^2}$ , hence  $y^p + y \not\equiv 0 \pmod{y^2}$ . Therefore the univariate polynomial (in  $x$ ) is irreducible over the field of fractions and in particular over  $\mathbb{F}_q[y]$ . This implies the bivariate polynomial is irreducible over the field  $\mathbb{F}_q$ . ■

We are now ready to recall Bézout's Theorem and apply it to prove  $S$  is indeed  $\epsilon$ -biased.

**Theorem 4** (Bézout's Theorem [2, Section 5.3]). *Suppose  $\phi$  and  $\psi$  are two bivariate polynomials over some field. If  $\phi$  and  $\psi$  have more than  $\deg(\phi) \cdot \deg(\psi)$  common roots than they have a common factor.*

**Theorem 5.** *For every  $k$  and  $\epsilon$  such that  $\epsilon < \frac{1}{\sqrt{k}}$ ,  $S$  is an  $\epsilon$ -biased set over  $k' = \Omega(k)$  bits of size  $O\left(\frac{k}{\epsilon^2 \log(1/\epsilon)}\right)^{5/4}$ .*

**Proof:** By Claim 1,  $|S| = |A| \cdot q = p^5 = O\left(\frac{k}{\epsilon^2}\right)^{5/4}$ .

Let  $T \subseteq [k']$  be some non-empty set. We identify  $[k']$  with the set  $\{(i, j) : i + j \leq \frac{r}{p+1}\}$  and  $T$  with the corresponding subset.

Let  $s \in S$  be an element specified by the pair  $((a, b), c) \in A \times \mathbb{F}_q$ . Then,

$$\sum_{(i,j) \in T} s_{(i,j)} = \sum_{(i,j) \in T} \langle \text{bin}(a^i b^j), \text{bin}(c) \rangle_2 = \left\langle \text{bin}\left(\sum_{(i,j) \in T} a^i b^j\right), \text{bin}(c) \right\rangle_2.$$

The polynomial  $\phi_T = \sum_{(i,j) \in T} x^i y^j$  is a non-zero polynomial. Clearly, for any  $(a, b)$  which is not a root of  $\phi_T$ , the inner-product will be unbiased when ranging over  $c$  (i.e. exactly half of the values for  $c$  will make the inner product 0). From the assumption  $\epsilon < \frac{1}{\sqrt{k}}$  it follows that  $\deg(\phi_T) < p + 1$ . Hence, by Claim 3 it follows that  $\phi_T$  and  $y^p + y - x^{p+1}$  have no common factors. Therefore, by Bezout's theorem we conclude that the number of roots of  $\phi_T$  that are in  $A$  is at most  $\frac{r}{p+1} \cdot (p + 1) = r$ , and,

$$\frac{1}{|S|} \left| \sum_{s \in S} (-1)^{\sum_{i \in T} s_i} \right| \leq \frac{r}{|A|} = \epsilon.$$

■

**Remark 6.** *The above construction can be improved to an  $\epsilon$ -biased set of size  $O\left(\frac{k}{\epsilon^2 \log(1/\epsilon)}\right)^{5/4}$  for every  $k$  and  $\epsilon$  such that  $\frac{\epsilon}{\sqrt{\log(1/\epsilon)}} < \frac{1}{\sqrt{k}}$ . To achieve this we choose  $p = \Theta\left(\frac{k}{\epsilon^2 \log(1/\epsilon ps)}\right)^{1/4}$ . We then observe that instead of taking a basis for  $V$  over  $\mathbb{F}_q$ , we can actually afford to take a basis over  $\mathbb{F}_2$ . Finally we need to use the fact that by the constraints we have on  $\epsilon$ , it follows that  $\log(1/\epsilon) = \Theta(\log(p))$ . When we restate the construction in algebraic function fields terminology, we also include this improvement.*

### 3 Restating the construction in algebraic function fields terminology

Without putting the above construction in the proper context, it may appear coincidental. We now describe the general framework of algebraic-geometric codes and explain why the above construction fits into this framework.

#### 3.1 Algebraic-Geometry

We recall a few notions from the theory of algebraic function fields. A detailed exposition of the subject can be found, e.g., in [6].

$\mathbb{F}_q$  denotes the finite field with  $q$  elements.  $\mathbb{F}_q(x)$ , where  $x$  is transcendental over  $\mathbb{F}_q$ , is the *rational* function field, and it contains all rational functions in  $x$  with coefficients in  $\mathbb{F}_q$ .  $F/\mathbb{F}_q$  is an algebraic function field, if  $F$  is a finite algebraic extension of  $\mathbb{F}_q(x)$ .

A *place*  $P$  of  $F/\mathbb{F}_q$  is a maximal ideal of some valuation ring  $O$  of the function field. We denote by  $O_P$  the valuation ring that corresponds to the place  $P$ . We denote by  $v_P$  the *discrete valuation* that corresponds to the valuation ring  $O_P$ . Therefore, we can write  $P$  and  $O_P$  as

$$P = \{x \in F : v_P(x) > 0\} \quad \text{and} \quad O_P = \{x \in F : v_P(x) \geq 0\}.$$

Since  $P$  is a maximal ideal,  $F_P = O_P/P$  is a field. For every  $x \in O_P$ ,  $x(P)$  denotes  $x \pmod{P}$  and is an element of  $F_P$ . The degree of a place  $P$  is defined to be  $\deg(P) = [F_P : \mathbb{F}_q]$ . In particular, if a place is of degree 1 then  $F_P$  is isomorphic to  $\mathbb{F}_q$ .  $\mathcal{P}_F$  is the set of places of  $F$ .  $N(F)$  is the number places of degree 1 (also called *rational points*) in  $F/\mathbb{F}_q$  and is always finite.

$\mathcal{D}_F$  is the free abelian group over the places of  $F$ . A *divisor* is an element in this group, i.e., it is a sum  $G = \sum_{P \in \mathcal{P}_F} n_P P$  with  $n_P \in \mathbb{Z}$  and where  $n_P \neq 0$  for only a finite number of places. We also denote  $v_P(G) = n_P$ . The *degree* of the divisor  $\sum_P n_P P$  is defined to be  $\sum_P n_P \cdot \deg(P)$ , and it is always finite. We say  $G_1 \geq G_2$  if  $G_1$  is component-wise larger than  $G_2$ , i.e.,  $v_P(G_1) \geq v_P(G_2)$  for any place  $P$ .

Each element  $0 \neq x \in F$  is associated with two divisors. The first is called the *principal divisor* of  $x$  and it is defined by

$$(x) = \sum_P v_P(x)P.$$

The degree of a principal divisor is always 0. The second is the *pole divisor* of  $x$  and it is defined by

$$(x)_\infty = \sum_{P: v_P(x) < 0} v_P(x)P.$$

If  $x \in F \setminus \mathbb{F}_q$  then  $\deg((x)_\infty) = [F : \mathbb{F}_q(x)]$ .

For a divisor  $G$ , we define the *Riemann-Roch space* is

$$\mathcal{L}(G) = \{x \in F : (x) \geq -G\} \cup \{0\}.$$

We define the dimension of  $G$  by  $\dim(G) = \dim \mathcal{L}(G)$  and we use the two notations interchangeably. The fact that the degree of each principal divisor is 0 implies that if  $\deg(G) < 0$  then  $\dim(\mathcal{L}(G)) = 0$ .

### 3.1.1 Geometric Goppa Codes

A Goppa code is specified by a triplet  $(F, Y, G)$ , where  $F/\mathbb{F}_q$  is a function field,  $Y = \{P_1, \dots, P_n\}$  is a set of places of degree 1 and  $G$  is an arbitrary divisor with no support over any place in  $Y$ . Notice that for any  $x \in \mathcal{L}(G)$ ,  $v_{P_i}(x) \geq 0$  and therefore  $x \in O_{P_i}$  and  $x(P_i) \in \mathbb{F}_q$ . The triplet  $(F, Y, G)$  specifies the code:

$$C(Y; G) = \{(x(P_1), \dots, x(P_n)) : x \in \mathcal{L}(G)\} \subseteq \mathbb{F}_q^n.$$

**Claim 7** ([6, Thm II.2.2]). *If  $\deg(G) < n$  then  $C(Y; G)$  is an  $[n, \dim(\mathcal{L}(G)), n - \deg(G)]$  linear code over  $\mathbb{F}_q$ .*

We want the gap between  $\dim(\mathcal{L}(G))$  and  $\deg(G)$  to be small. It turns out that for any function field  $F/\mathbb{F}_q$  there exists a constant  $g \in \mathbb{N}$ , such that for any divisor  $G \in \mathcal{D}_F$ ,  $\deg(G) - \dim(\mathcal{L}(G)) \leq g - 1$ . The minimal integer with this property is called the *genus* of  $F/\mathbb{F}_q$ . The Riemann-Roch Theorem says that:

**Theorem 8** ([6, Thm I.5.17]). *If  $\deg(G) \geq 2g - 1$  then  $\dim(\mathcal{L}(G)) = \deg(G) - g + 1$ .*

This, in particular, allows one to easily compute the dimension of the code when  $\deg(G) > 2g$ . The only remaining question is how many places of degree 1 exist. Informally, the Drinfeld-Vladut bound tells us that when  $g$  tends to infinity,  $n \leq g(\sqrt{q} - 1)$ , and several explicit constructions meet this bound (see [3, Chapter 1]).

In this paper we look at divisors  $G$  whose degree is smaller than the genus. Much less is known about such small-degree divisors. In this regime,  $\dim(\mathcal{L}(G))$  depends on the divisor  $G$  itself, and not only on its degree, as is the case when  $\deg(G) > 2g$ . For some special algebraic function fields the vector space  $\mathcal{L}(G)$  (and therefore also its dimension) is known in full. We talk more about this below.

### 3.2 Concatenating AG codes with Hadamard

We concatenate an outer code with the Hadamard code. If the outer code is an  $[n_1, k_1, d]_q$  code and  $q$  is a power of two, then concatenating it with the  $[2^{k_2}, k_2 = \log(q), \frac{1}{2}]_2$  Hadamard code gives an  $[n = 2n_1q, k = k_1(1 + \log q)]_2$  code that is  $\epsilon = \frac{n_1 - d}{n_1}$  balanced, because non-zero symbols in the outer code expand by the concatenation to perfectly balanced blocks.

Using Reed-Solomon code as the outer code, one gets an  $[n = 2q^2, k = k_1(\log q + 1)]_2$  code that is  $\epsilon = \frac{k_1}{q}$  balanced. Rearranging parameters, this gives an  $[n, k]_2$   $\epsilon$ -balanced code with  $n = O((\frac{k}{\epsilon \log(\frac{k}{\epsilon})})^2)$ . This is one of the constructions in [1].

Taking the outer code to be an AG code  $C(Y; G)$  over  $\mathbb{F}_q$ , with  $\deg(G) > 2g$  and optimal length  $g\sqrt{q}$ , one gets an  $[n = g\sqrt{q}, k = \deg(G) + 1 - g]_2$  code that is  $\epsilon = \frac{\deg(G)}{g\sqrt{q}}$ -biased. Doing the calculation one sees that  $n = O(\frac{k}{\epsilon^3 \log(1/\epsilon)})$ . As these are the best AG codes possible for the case  $\deg(G) > 2g$ , no improvement is possible here unless we consider low-degree divisors  $G$ .

So we now turn our attention to the case where  $\deg(G) \leq 2g - 1$ . In this case  $\dim \mathcal{L}(G)$  depends on the divisor  $G$  and not just its degree. One special case is the case where  $G = rQ$ ,  $r \in \mathbb{N}$  and  $Q$  is a place of degree 1. For any such  $r$ ,  $\dim \mathcal{L}(rQ)$  is either equal to  $\dim \mathcal{L}((r - 1)Q)$  or to  $\dim \mathcal{L}((r - 1)Q) + 1$ . In the former case  $r$  is said to be a *gap number* of  $Q$ . Weierstrass Gap Theorem [6, Thm I.6.7] says that for any place  $Q$  there are exactly  $g = \text{genus}(F/\mathbb{F}_q)$  gap numbers, and they are all in the range  $[1, 2g - 1]$ .

The non-gap numbers (also called *pole numbers*) form a semigroup of  $\mathbb{N}$  (i.e. a set that is closed under addition). This semigroup is sometimes referred to as the *Weierstrass semigroup* of  $Q$ . We say a semi-group  $S$  is generated by a set of elements  $\{g_i\}$ , if each  $g_i \in S$  and, furthermore, every element  $s \in S$  can be expressed as  $s = \sum a_i g_i$  with  $a_i \in \mathbb{N}$ .

The structure of the Weierstrass semigroup is crucial to our construction. We know that there are exactly  $g$  elements of this semigroup in the range  $[1, 2g]$ . If these elements are too concentrated on the upper side of the range then the behavior of  $\dim \mathcal{L}(rQ)$  will be very similar to the case where  $r > 2g - 1$ . Thus, our goal is to find a function field  $F$  that has many places of degree 1, say,  $N(F) \geq \Omega(g\sqrt{q})$ , while at the same time  $F$  has a degree 1 place  $Q$  with a "good" Weierstrass semigroup.

### 3.3 The Construction

Let  $p$  be a prime power and  $q = p^2$ . The Hermitian function field over  $\mathbb{F}_q$  can be represented as the extension field  $\mathbb{F}_q(x, y)$  of the rational function field  $\mathbb{F}_q(x)$  with  $y^p + y = x^{p+1}$ . This

function field has  $1 + p^3$  places of degree one. First, there is the common pole  $Q_\infty$  of  $x$  and  $y$ . Moreover, for each pair  $(\alpha, \beta) \in \mathbb{F}_q$  with  $\beta^p + \beta = \alpha^{p+1}$  there is a unique place  $P_{\alpha,\beta}$  of degree one such that  $x(P_{\alpha,\beta}) = \alpha$  and  $y(P_{\alpha,\beta}) = \beta$  and we already saw there are  $p^3$  such points. The genus of the Hermitian function field is  $g = p(p-1)/2$ .

For the outer code we take the Goppa code  $C_r = C(Y, G = rQ_\infty)$ , where  $Y$  is the set of all degree 1 places  $P_{\alpha,\beta}$  mentioned above and  $r = \epsilon p^3$ . The Weierstrass semigroup of  $G$  is generated by  $p$  and  $p+1$ , and a basis for  $\mathcal{L}(G) = \mathcal{L}(rQ_\infty)$  is

$$\{x^i y^j : j \leq p-1 \text{ and } ip + j(p+1) \leq r\}.$$

The dimension of the code is

$$|\{(i, j) : j \leq p-1 \text{ and } ip + j(p+1) \leq r\}|.$$

We can now see the similarity between this construction and the one in Section 2. The parameter  $r$  will be chosen such that the constraint  $j \leq p-1$  will be nullified. Therefore, both use evaluations of low degree bivariate polynomials over the same set of  $p^3$  points.<sup>2</sup>

**Theorem 9.** *For every  $k$  and every  $\epsilon$  such that  $\frac{\epsilon}{\sqrt{\log(1/\epsilon)}} \leq \frac{1}{\sqrt{k}}$ , there exists an explicit  $[n, \Omega(k)]_2$  code that is  $\epsilon$ -balanced, with  $n = O\left(\frac{k}{\epsilon^2 \log(1/\epsilon)}\right)^{5/4}$ .*

**Proof:** For a given  $k$  and  $\epsilon$ , let

$$p \in \left[ \left( \frac{k}{\epsilon^2 \log(1/\epsilon)} \right)^{1/4}, 2 \left( \frac{k}{\epsilon^2 \log(1/\epsilon)} \right)^{1/4} \right]$$

be a power of two. It can be verified that  $\frac{1}{p^3} \leq \epsilon \leq \frac{1}{p}$  and so  $\log(1/\epsilon) = \Theta(\log(p))$ .

Let  $r = \epsilon p^3$  and let  $\mathbb{F}_q$  be the field with  $q = p^2$  elements. Let  $F$  denote the Hermitian function field over  $\mathbb{F}_q$  and let  $Y$  denote its set of places of degree 1, excluding  $Q_\infty$ . This implies that  $|Y| = p^3$ . Define the divisor  $G$  to be  $G = rQ_\infty$ . Since  $r \leq p^2$ ,  $\dim \mathcal{L}(rQ_\infty) \geq \left(\frac{r}{2(p+1)}\right)^2 = \Omega(\epsilon^2 p^4) = \Omega\left(\frac{k}{\log(p)}\right)$ . By Claim 7, the Goppa code that is obtained from the triplet  $(F, Y, G)$  is a

$$[p^3, \Omega\left(\frac{k}{\log(p)}\right), p^3 - r]_{p^2}$$

code. Concatenating this code with Hadamard gives a  $[p^5, \Omega(k)]_2$  code that is  $\epsilon$ -balanced (since  $\frac{r}{p^3} = \epsilon$ ). Now, by our choice of  $p$ , it follows that

$$\frac{k}{\epsilon^2 \log\left(\frac{1}{\epsilon}\right)} = \Theta(p^4)$$

and therefore  $n = p^5 = O\left(\left(\frac{k}{\epsilon^2 \log\left(\frac{1}{\epsilon}\right)}\right)^{5/4}\right)$  as desired. ■

<sup>2</sup>The only slight difference is that in this construction we take all bivariate polynomials with bounded *weighted* total degree. However, the weight is nearly identical for both variables and so this does not affect much the parameters of the construction.

## 4 A hypothesis and its consequence

The power of AG codes comes from the fact that the number of rational points  $N(F)$  may be larger by a factor of  $\sqrt{q}$  than the genus  $g(F)$ . The genus measures the maximal loss in dimension compared to the degree. By the Drinfeld-Vladut theorem, this gain is limited to a factor of  $\sqrt{q}$ .

Motivated by the results of the previous section we start with a simple question. What is the maximal gap between the number of rational points  $N(F)$  and the degree of a divisor  $G$  with some *positive* dimension. A few definitions are in place:

**Definition 2.** A sequence  $\mathcal{F}$  over  $\mathbb{F}_q$  is an infinite sequence  $\mathcal{F} = (F_1, F_2, \dots)$  of function fields  $F_i/\mathbb{F}_q$  such that  $\lim_{i \rightarrow \infty} N(F_i) = \infty$ .

**Definition 3.** We say a divisor  $G \in \mathcal{D}_F$  has positive dimension if  $\dim(\mathcal{L}(G)) > 1$ .

**Definition 4.** Let  $\mathcal{F}$  be a sequence over  $\mathbb{F}_q$ , and  $0 \leq b = b(q) \leq 1$ . We say  $\mathcal{F}$  has a  $b$ -gap, if there exist divisors  $G_i \in \mathcal{D}_{F_i}$  with positive dimension such that  $\lim \frac{N(F_i)}{\deg(G_i)} = b$ .

For example, any divisor with degree above the genus has positive dimension. Thus, if  $F$  is asymptotically optimal (meaning that  $\lim N(F_i)/g(F_i) \geq \sqrt{q} - 1$ ) then  $b = \sqrt{q} - 1$ .

In fact, we can do better. We already saw function fields with a larger gap. In the Hermitian function field,  $G = pQ_\infty$  has positive dimension, and  $N(F)/\deg(G) \geq p^3/p = p^2 = q$ . One can also build a tower over the Hermitian function field and get a sequence that preserves this gap.

Can the gap be larger? The following argument, shown to us by Henning Stichtenoth, shows this is not possible:

**Lemma 10.** Let  $F/\mathbb{F}_q$  be a function field and  $G \in \mathcal{D}_F$  a divisor with positive dimension. Then  $N(F) \leq \deg(G) \cdot (q + 1)$ .

**Proof:** As  $\dim(\mathcal{L}(G)) > 1$ , there exists some  $x \in F \setminus \mathbb{F}_q$  such that  $(x) \geq -G$ . Fix any such  $x$ . In particular,  $\deg(x)_\infty \leq \deg(G)$ . Also, by [6, Thm I.4.11],  $\deg(x)_\infty = [F : \mathbb{F}_q(x)]$ . On the other hand, we may view  $F$  as a finite extension over the rational function field  $\mathbb{F}_q(x)$ . Every place of degree 1 of  $F$  lies above some place of degree 1 of  $\mathbb{F}_q(x)$ . There are exactly  $q + 1$  places of degree 1 of  $\mathbb{F}_q(x)$ , and each one of them may split to at most  $[F : \mathbb{F}_q(x)]$  places of degree 1 of  $F$  (by the fundamental equality, [6, Thm III.1.11]). Altogether,  $N(F) \leq (q + 1)[F : \mathbb{F}_q(x)] = (q + 1) \deg(x)_\infty \leq (q + 1) \deg(G)$ . ■

We now move on to what we actually want. We want an  $\Omega(q)$  gap between the number of degree 1 places and a divisor with a *large* dimension. We define:

**Definition 5.** We say a divisor  $G \in \mathcal{D}_F$  is  $c$ -dense, for  $0 \leq c \leq 1$ , if  $G \geq 0$  and  $\dim(\mathcal{L}(G)) \geq c \cdot \deg(G)$ .

**Definition 6.** Let  $\mathcal{F}$  be a sequence over  $\mathbb{F}_q$ , and  $0 \leq b = b(q), c = c(q) \leq 1$ . We say  $\mathcal{F}$  has a  $c$ -dense  $b$ -gap, if there exist  $c$ -dense divisors  $G_i \in \mathcal{D}_{F_i}$  such that  $\lim \frac{N(F_i)}{\deg(G_i)} = b$ .

When we use concatenation of a linear code over a large alphabet and a binary *linear* code and we want the resulting code to be linear, we require that the size of the large alphabet is a power of 2. Without this restriction, we may still construct binary codes with concatenation, but not necessarily linear ones.

**Definition 7.** An (not necessarily linear) code is  $\epsilon$ -balanced if the relative Hamming distance between every two codewords is in the range  $[\frac{1}{2} - \epsilon, \frac{1}{2} + \epsilon]$ .

**Lemma 11.** If  $\mathcal{F}$  has a  $c$ -dense  $b$ -gap, then for infinitely many  $k$  there exists an  $(n, k)_2$  code that is  $\epsilon = \frac{1}{b-1}$  balanced and  $n = \Delta \cdot \frac{k}{\epsilon^2 \log(\frac{1}{\epsilon})}$  where  $\Delta = \frac{1}{c} \frac{q}{b-1} \cdot \frac{\log(b-1)}{\log(q)}$ .

**Proof:** Let  $G_i \in \mathcal{D}_{F_i}$  be the promised divisors. Let  $Y_i$  be the set of all places of degree 1 in  $F_i$  that are disjoint with  $G_i$ .  $|Y_i| \geq N(F_i) - \deg(G_i)$ . The Goppa code that is obtained from the triplet  $(F_i, D_i, G_i)$  is an

$$[N = N(F_i) - \deg(G_i), c \cdot \deg(G_i), N - \deg(G_i)]_q$$

code. Concatenating this with Hadamard gives a

$$(n = q \cdot (N(F_i) - \deg(G_i)), k = c \cdot \deg(G_i) \log(q))_2$$

code which is  $\epsilon = \frac{\deg(G_i)}{N(F_i) - \deg(G_i)}$ -balanced. Taking  $i$  large enough, we essentially get  $\epsilon = \frac{1}{b-1}$ . Thus,

$$n = q(N(F_i) - \deg(G_i)) \approx q \cdot (b-1) \cdot \deg(G_i) \approx q \cdot \frac{\deg(G_i)}{(b-1)\epsilon^2} = \frac{q \log(\frac{1}{\epsilon})}{(b-1)c \log q} \cdot \frac{k}{\epsilon^2 \log(\frac{1}{\epsilon})}. \quad \blacksquare$$

For example, if  $\mathcal{F}$  is asymptotically optimal and  $\deg(G_i) \geq 2g$  then  $b = \Theta(\sqrt{q})$  and  $c \geq \frac{1}{2}$ . Another example is given by the divisors we chose for the Hermitian function field. There we chose  $\deg(G_i) = r$  and we had  $\dim(G_i) = \Theta(\frac{r^2}{p^2})$ , thus  $c = \Theta(\frac{r}{p^2})$  and  $b = \Theta(\frac{p^3}{r})$  and altogether  $bc = \Theta(\sqrt{q})$  (the constants in the big- $O$  notation are independent of  $q$ ).

If  $bc$  can be made  $\Omega(q)$ , with a constant independent of  $q$ , then one would match the lower bound for  $\epsilon$ -balanced codes. We formulate this hypothesis below. We have no idea whether it is valid. On the one hand, we saw that one can find divisors  $G_i \in \mathcal{D}_{F_i}$  with positive dimension and  $q$ -gap. On the other hand, we were unable to find such divisors with *large density* and so, we do not know of an example where  $bc \gg \sqrt{q}$ .

**Hypothesis 12.** There exist positive constants  $\beta, c < 1$  and an infinite set  $Q \subseteq \mathbb{N}$  such that for every  $q \in Q$  there exists a sequence  $\mathcal{F}$  over  $\mathbb{F}_q$  that has a  $c$ -dense  $(\beta q)$ -gap.

**Theorem 13.** Under Hypothesis 12 there are infinitely many parameters  $k, \epsilon$  such that there exists an  $(n, k)_2$  code that is  $\epsilon$ -balanced with  $n = O\left(\frac{k}{\epsilon^2 \log(1/\epsilon)}\right)$ .

## References

- [1] N. Alon, O. Goldreich, J. Håstad, and R. Peralta. Simple constructions of almost  $k$ -wise independent random variables. *Random Structures and Algorithms*, 3(3):289–303, 1992.
- [2] W. Fulton. *Algebraic Curves*. Third edition, 2008.
- [3] A. Garcia and Eds. Stichtenoth, H. *Topics in Geometry, Coding Theory and Cryptography (Algebra and Applications)*. Springer-Verlag, 2006.

- [4] S. Lang. *Algebra*. Springer, revised third edition, 2002.
- [5] J. Naor and M. Naor. Small-bias probability spaces: Efficient constructions and applications. *SIAM Journal on Computing*, 22(4):838–856, 1993.
- [6] H. Stichtenoth. *Algebraic function fields and codes*. Springer, 2003.
- [7] H. Stichtenoth. Private communication, 2009.
- [8] M.A. Tsfasman, S.G. Vladutx, and T. Zink. Modular curves, Shimura curves, and Goppa codes, better than Varshamov-Gilbert bound. *Mathematische Nachrichten*, 109(1), 1982.