

Divisible subdivisions

Noga Alon *

Michael Krivelevich †

Abstract

We prove that for every graph H of maximum degree at most 3 and for every positive integer q there is a finite $f = f(H, q)$ such that every K_f -minor contains a subdivision of H in which every edge is replaced by a path whose length is divisible by q . For the case of cycles we show that for $f = O(q \log q)$ every K_f -minor contains a cycle of length divisible by q , and observe that this settles a recent problem of Friedman and the second author about cycles in (weakly) expanding graphs.

1 Introduction

There are several known results asserting that any graph with a sufficiently large minimum (or average) degree contains a cycle of prescribed length modulo a given parameter. An early result of this form appears in [3]: for every odd k there exists a $c(k)$ so that every graph with minimum degree at least $c(k)$ contains a cycle of length ℓ modulo k for every integer ℓ . A similar result holds for every even k and every even ℓ (but of course not for even k and odd ℓ as shown by dense bipartite graphs.) Thomassen proved in [12] that for non-bipartite 2-connected graphs a result as above exists also for even k and odd ℓ .

For graphs with large chromatic number stronger conclusions hold. Another result established in [12] addresses this case: For any two positive integers m and k there exists a number $c(m, k)$ such that the following holds. For every assignment of two natural numbers $k(e) \leq k$ and $d(e)$ for each edge e of K_m , any graph of chromatic number at least $c(m, k)$ contains a subdivision of K_m in which each edge e corresponds to a path of length $d(e)$ modulo $k(e)$.

*Department of Mathematics, Princeton University, Princeton, NJ 08544, USA and Schools of Mathematics and Computer Science, Tel Aviv University, Tel Aviv 6997801, Israel. Email: nogaa@tau.ac.il. Research supported in part by NSF grant DMS-1855464, BSF grant 2018267 and the Simons Foundation.

†School of Mathematical Sciences, Tel Aviv University, Tel Aviv 6997801, Israel. Email: krivelev@tauex.tau.ac.il. Research supported in part by ISF grant 1261/17 and by USA-Israel BSF grant 2018267.

A more recent result with a similar flavor is proved in [11]: for every k there is a $c(k)$ so that every graph with average degree at least $c(k)$ contains a subdivision of K_k in which every edge is subdivided the same number of times.

A common feature of these results and related ones is that they apply only to graphs with a rather large average degree. In particular, if the average degree is just a bit above 2, then none of these results holds, and indeed there are simple examples showing no such result can hold if we do not make any additional assumptions. In this note we prove a result applicable to very sparse graphs. The sufficient condition we give is based on complete minors; since there exist minors of arbitrarily large complete graphs with average degree arbitrarily close to 2 in every subgraph, this imposes essentially the weakest possible condition in terms of graph density.

Theorem 1. *For every graph H of maximum degree at most 3, and for every positive integer q there is a finite $f = f(H, q)$ such that every K_f -minor G contains a subdivision of H in which every edge is replaced by a path whose length is divisible by q .*

Remark: For every f and q there is a K_f -minor G with maximum degree 3, in which any path between two vertices of degree 3 is of length divisible by q ; obviously such G does not contain a subdivision of any graph H with maximum degree $\Delta(H) > 3$. Hence, the assumption that H has maximum degree at most 3 is needed. Similarly the conclusion about paths of length 0 modulo q cannot be replaced by any other residue modulo q , unlike the results in [3] or [12] mentioned above.

We make essentially no attempt to optimize the value of $f = f(H, q)$, and the problem of determining its asymptotic behavior remains open.

It is well known that graphs without small separators contain large complete minors (the line of research establishing these results started in [2] and culminated with [8]). Our main result is thus applicable to such graphs. Also, non-existence of sublinear separators is essentially equivalent to weak expansion (see [10] for a discussion), hence Theorem 1 can be applied to the class of (weakly) expanding graphs.

In the very special case of a cycle of length divisible by q we get a nearly tight bound, proving the following.

Theorem 2. *For every positive integer q there is a $g = g(q) = O(q \log q)$ such that every K_g -minor G contains a cycle of length divisible by q . Moreover, if q is a prime then $g(q) < 4q$.*

This, together with the fact that expanders contain large clique minors (see [10] for a background), settles a problem raised recently in [6].

For illustrative purposes let us mention that, confirming a conjecture of Thomassen [12], Gao, Huo, Liu and Ma proved [7] that requiring minimum degree $\delta(G) \geq q + 1$ guarantees cycles of all even lengths modulo q , including of course cycles of length divisible by q ; requiring $\delta(G) \geq q$ is necessary to guarantee a cycle of length divisible by q , at least for odd q , as shown by the complete bipartite graph $K_{q-1,n}$ for $n \geq q - 1$. In contrast, our result allows to argue about existence of cycles of length divisible by q in much sparser graphs.

The proofs are described in the next two sections. The final section contains some concluding remarks and open problems.

2 Subdivisions

In this section we prove Theorem 1. We start by proving a key lemma. Here and later by the weight of a path P in an edge-weighted graph G we mean the sum of the weights of all edges in P .

Lemma 2.1. *There is a function $f_1(k, q)$ satisfying the following condition. Let $k, q \geq 2$ be integers. Let T be a tree with edges labeled by the elements of Z_q , and let L be a set of specified leaves of T of cardinality $|L| = f_1(k, q)$. Then there are a subset $L_0 \subset L$ of k leaves and a residue $a \in Z_q$ such that for every three leaves $x_1, x_2, x_3 \in L_0$ there is a vertex $v \in V(T)$ with all paths from v to x_i in T being disjoint outside of v and having all weight a modulo q .*

Proof. We set with foresight

$$f_1 = f_1(k, q) = ((k - 1)q + 1)^{(k-1)q^2+1}.$$

Assume that T, L are as given in the lemma. Observe that if the lemma's conclusion holds for a subtree $T' \subseteq T$ containing L then it holds for T . Hence we can assume that T is a minimal by inclusion tree containing L .

For convenience root T at an arbitrary vertex $r \in V(T)$ with $d_T(r) \geq 3$. For a vertex $v \in V(T)$ denote by T_v the subtree of T rooted at v (with respect to r).

Consider first the case where there is a vertex $v \in V(T)$ with $d_T(v) \geq (k - 1)q + 2$. By the minimality of T , for each child u of v the subtree T_u contains a leaf $\ell(u) \in L$, the paths from v to all such $\ell(u)$ are disjoint outside of v . By the pigeonhole principle there is

a subset U_0 of the children of v in T of cardinality $|U_0| = \lceil \frac{d_T(v)-1}{q} \rceil \geq k$ such that all paths from v to $\ell(u)$, $u \in U_0$, are of the same weight a modulo q . The set $L_0 = \{\ell(u) : u \in U_0\}$ fulfills then the requirement of the lemma.

Now we treat the complementary case $\Delta(T) \leq (k-1)q + 1$. Let t be the maximal number of vertices of degree at least three on a path from r to a leaf $x \in L$. Since every $x \in L$ is uniquely determined by the sequence of edges leaving the vertices of degree at least three on the unique path from r to x in T , and the number of such sequences is obviously at most $(\Delta(T))^t$, we obtain: $|L| \leq (\Delta(T))^t$, implying $t \geq (k-1)q^2 + 1$. Let P be a path from r to a leaf of T with t vertices of degree at least three in T along it, and let $U_1 \subset V(P)$ be these vertices, $|U_1| = t \geq (k-1)q^2 + 1$. By the pigeonhole principle, there is a subset $U_2 \subset U_1$ of cardinality $|U_2| = \lceil \frac{|U_1|}{q} \rceil \geq (k-1)q + 1$ such that all subpaths of P from r to $u \in U_2$ have the same weight modulo q . This implies that for every $u_1 \neq u_2 \in U_2$ the subpath of P between u_1 and u_2 has weight 0 modulo q . By the minimality of T , every vertex $u \in U_2$ contains a leaf $\ell(u) \in L$ in its subtree T_u , where all these leaves are distinct. Applying the pigeonhole principle again, we derive the existence of a subset $U_3 \subset U_2$ of cardinality $|U_3| = k$ such that every path between $u \in U_3$ and the corresponding leaf $\ell(u) \in T_u$ has the same weight a modulo q . Set $L_0 = \{\ell(u) : u \in U_3\}$. We claim that L_0 meets the requirement of the lemma. Indeed, let u_1, u_2, u_3 be distinct vertices in U_3 ordered in the order of their appearance along P . Then the paths from u_2 to $\ell(u_i)$, $i = 1, 2, 3$, are all disjoint outside of u_2 and have total weight a modulo q (the part along P has weight 0 modulo q , and the appended part from u_i to $\ell(u_i)$ has weight $a \pmod{q}$). \square

We can now prove Theorem 1, making no attempt to optimize the estimate for f . Let Γ be a Ramsey graph in q colors for the $(q-1)$ -subdivision of H . Assume that the vertex set of Γ is $[N]$, denote $k = 2|E(\Gamma)|$. Let $M = f_1(k, q)$, with $f_1(k, q)$ from Lemma 2.1. Finally, set:

$$f = M + (N-1)q + 1.$$

Assume G is a minor of K_f with supernodes $X_1, \dots, X_{(N-1)q+1}, Y_1, \dots, Y_M$. For a pair of supernodes X_i, Y_j , if $e = (x, y)$ is an edge connecting X_i to Y_j then split e by a vertex z , assign weights $w((x, z)) = 0$, $w((z, y)) = 1$, and append (x, z) to X_i . We assign weight 1 to all remaining edges of G .

Let T_i be a spanning tree of X_i and let L_i be a set of $M = f_1(q, k)$ distinct leaves of T_i , each connected to a different supernode Y_j . Apply Lemma 2.1 to (T_i, L_i) to get a subset L'_i of cardinality $|L'_i| = k$ and a residue $a_i \in Z_q$ with the properties guaranteed by

the lemma. Invoking the pigeonhole principle with respect to the multiset of residues $\{a_i\}$ we conclude that there exists a subset $I \subset [(N-1)q+1]$ of cardinality $|I| = N$ with all residues $a_i, i \in I$, taking the same value a . By renumbering if necessary we can assume $I = [N]$.

Now we go sequentially over all edges $e = (i_1, i_2) \in E(\Gamma)$ and connect the corresponding supernodes X_{i_1}, X_{i_2} as follows: choose a previously unused supernode Y_{j_1} having a neighbor in L'_{i_1} , choose a distinct and previously unused supernode Y_{j_2} having a neighbor in L'_{i_2} , and then connect Y_{j_1} and Y_{j_2} . Concatenating, we obtain a path P_e in G from a leaf $x_1 \in L'_{i_1}$ to a leaf $x_2 \in L'_{i_2}$, where all these paths are vertex disjoint for different edges $e = (i_1, i_2) \in E(\Gamma)$.

Define a coloring $c : E(\Gamma) \rightarrow Z_q$ as follows: for $e = (i_1, i_2) \in E(\Gamma)$, its color $c(e)$ is equal to the weight modulo q of the path P_e between X_{i_1} and X_{i_2} . By the choice of Γ , the so obtained coloring c induces a monochromatic copy H^* of the $(q-1)$ -subdivision of H , say in color $b \in Z_q$. Let $I_0 \subset [N]$ be the subset of supernodes corresponding to the vertices of H in this subdivision. By construction, for each edge $f = (i_1, i_2) \in E(H)$, $i_1, i_2 \in I_0$, the graph G contains a path Q_f from X_{i_1} to X_{i_2} passing through a sequence of $q-1$ intermediate supernodes X_i , with the intermediate supernodes being distinct for distinct edges. Each such path enters and leaves X_i through vertices of L'_i ; by the definition of L'_i and the choice of a , the entrance and departure point in X_i can be connected by a path of length $2a$ modulo q . The weight of Q_f between two consecutive intermediate supernodes X_i , and also between X_{i_1} and the first intermediate supernode, and between the last intermediate supernode and X_{i_2} is b modulo q . Finally, for each supernode X_i , $i \in I_0$, the (at most three) paths Q_f leaving X_i all depart from the vertices of L'_i , hence we can choose a vertex $v_i \in X_i$ connected to the departure points by disjoint paths of weight a each. Collecting all weights, we conclude that for each edge $f = (i_1, i_2) \in E(H)$, the path between v_{i_1} and v_{i_2} in G has total weight:

$$2a + (q-1) \cdot 2a + q \cdot b = (2a+b)q \equiv 0 \pmod{q}.$$

We have thus obtained the required subdivision of H , completing the proof. \square

3 Cycles

In this section we prove Theorem 2. In the lemma below and later, a complete digraph is a digraph in which every pair of vertices is connected by one edge in each of the two directions.

Lemma 3.1. *Let $q \geq 2$ be an integer, and let $\Gamma = (V, E)$ be a complete digraph on $[2q \ln q]$ vertices with weights $w(e) \in Z_q$ on its edges. Then Γ contains a directed cycle C of total weight divisible by q .*

Proof. The proof borrows its main idea from the argument in [1]. Here though we are in a more favorable situation dealing with the complete digraph and can thus allow ourselves to employ a simpler probabilistic tool for the proof – the union bound (instead of the Local Lemma used in [1]).

Let $c : V \rightarrow Z_q$ be a random labeling of V by the elements of Z_q . For a vertex $u \in V$ denote by A_u the event “ u has an outneighbor v satisfying $c(v) = c(u) + w(uv)$ ”. In order to estimate $Pr[\overline{A_u}]$ observe that by conditioning on the label $c(u)$, the probability that none of the outneighbors v of u satisfies $c(v) = c(u) + w(uv)$ is $(1 - 1/q)^{|V|-1} < 1/|V|$. Hence by the union bound there is a choice of c for which all of the events A_u hold. Fix such a choice, and for every vertex $u \in V$ choose an outgoing edge (u, v) so that $c(v) = c(u) + w(uv)$. In the subgraph of Γ obtained this way every outdegree is 1 and hence it has a directed cycle $C = (u_1, \dots, u_\ell, u_1)$. Summing all weights along the edges of C and denoting $u_{\ell+1} = u_1$ we obtain:

$$\sum_{i=1}^{\ell} w(u_i u_{i+1}) = \sum_{i=1}^{\ell} (c(u_{i+1}) - c(u_i)) \equiv 0 \pmod{q},$$

as required. □

If q is a prime number then the logarithmic term in the above lemma can be omitted, thus giving an asymptotically optimal order of magnitude.

Lemma 3.2. *Let $q \geq 2$ be a prime, and let $\Gamma = (V, E)$ be a complete digraph on $2q - 1$ vertices with weights $w(e) \in Z_q$ on its edges. Then Γ contains a directed cycle C of total weight divisible by q .*

Proof. If there are two vertices u, v so that $w(u, v) = -w(v, u)$ (modulo q) then there is a directed cycle consisting of two edges satisfying the requirement, hence we may and will assume that there is no such pair of vertices. In this case we proceed to prove that for every $k < q$ there are distinct vertices $x_0, x_1, y_1, x_2, y_2, \dots, x_k, y_k$ in Γ and a set S of $k + 1$ distinct residues modulo q so that for every $s \in S$ there is a directed path P_s from x_0 to x_k of total weight s modulo q . Each path P_s consists of k subpaths p_1, p_2, \dots, p_k , in this order, where each p_i is either the single edge $x_{i-1}x_i$ or the two-edge path $x_{i-1}y_i x_i$. This is proved by induction on k . For $k = 0$ the required path is the trivial path with no edges

and $S = \{0\}$. Assuming the result holds for $k < q - 1$ we prove it for $k + 1$. Let u, v be two vertices of Γ that are not in the set $\{x_0, x_1, x_2, \dots, x_k, y_k\}$. If $w(x_k, u) = w(x_k, v) + w(v, u)$ and $w(x_k, v) = w(x_k, u) + w(u, v)$ then $w(u, v) = -w(v, u)$ contradicting the assumption. Thus at least one of these equalities does not hold; by renaming u and v if needed we may assume that $w(x_k, u) \neq w(x_k, v) + w(v, u)$. Define $x_{k+1} = u$ and $y_{k+1} = v$. Let T be the set of the two distinct residues $w(x_k, u) = w(x_k, x_{k+1})$ and $w(x_k, v) + w(v, u) = w(x_k, y_{k+1}) + w(y_{k+1}, x_{k+1})$. Clearly, for every residue $s \in S + T$, there is a path from x_0 to x_{k+1} of total weight s in Γ . By the Cauchy-Davenport theorem (for the easy special case in which one of the sets is of size 2), $|S + T| \geq |S| + |T| - 1 = k + 1$ establishing the induction step. Taking $k = q - 1$, the result shows that for every residue class s modulo q there is a directed simple path from x_0 to x_{q-1} of weight s . Choosing $s = -w(x_{q-1}, x_0)$ and adding the edge $x_{q-1}x_0$ gives the required cycle. \square

We can now prove Theorem 2. (In fact, our proof applies to the general weighted setting.) If q is not a prime define $N = \lceil 2q \log q \rceil$ and $g = 2N$, if it is a prime define $N = 2q - 1$ and $g = 2N$. Given a K_g -minor on the $2N$ supernodes X_i^+ and X_i^- for $1 \leq i \leq N$, assume without loss of generality that the induced subgraph on each supernode is a tree and that there is exactly one edge connecting each pair of supernodes. For each i let b_i be the weight of the unique edge, $x_i^- x_i^+$ connecting X_i^- and X_i^+ (1 if there are no weights). For each $i \neq j$ let $w'(ij)$ be the total weight modulo q of the unique path in the induced tree on the vertices $X_i^+ \cup X_j^-$ from the vertex x_i^+ to the vertex x_j^- . By the two lemmas above applied to the auxiliary complete directed graph on the vertices $\{1, 2, \dots, N\}$ with the weights $w(ij) = b_i + w'(ij)$ there is a directed cycle of total weight 0 modulo q in this auxiliary digraph. This gives the required cycle in the original graph. \square

4 Concluding remarks

- There are several ways to improve the bound for $f(H, q)$ in the proof of Theorem 1. In particular, one can use the constructions of Ramsey graphs for subdivisions given in [9, 5] as a seed for our proof. In addition, it is possible to take $k = \Delta(\Gamma)$, where Γ is the Ramsey graph used, and $M = 2|E(\Gamma)| + f_1(q, k)$. Then we first choose N supernodes X_i with the same value of a_i as in the present proof, fix a bijection between these N nodes and the vertex set of Γ , and then for each $i = 1, \dots, N$, look at the leaves of X_i connected to previously unused supernodes Y_j (altogether we use at most $2|E(\Gamma)|$ such supernodes Y_j , two per each edge of Γ throughout the proof).

There are at least $f_1(q, k)$ of them, this would be our set L_i . Next apply to it Lemma 2.1 to get a set of cardinality k (in fact here the degree of vertex i in Γ suffices), and then put aside the supernodes Y_j to which the edges from this k -set of leaves in L_i lead. This gives some improvement, but as is frequently the case with Ramsey-type results, the bound obtained is still huge, surely far from being optimal. It may be interesting to try to determine or to estimate the asymptotic behavior of the best possible bound for $f(H, q)$.

- An α -expander is a graph on n vertices in which every set X of at most $n/2$ vertices has at least $\alpha|X|$ neighbors outside X , see [10] for a general discussion. It is easy to see that any such graph has no sublinear separators and thus contains a K_f -minor for $f \geq c(\alpha)\sqrt{n}$ by [8]. Our results thus apply to such graphs and in particular imply that any such graph contains cycles of length divisible by q for any $q \leq \tilde{O}(n^{1/2})$. This settles a question posed explicitly in [6]. See [6] for further results about cycle lengths in α -expanders.
- There is a substantial amount of research on Ramsey-type problems for structures labelled by elements of an abelian group. Questions of this type are called zero-sum problems, see [4] for a survey of the subject (until the mid 90s). A typical problem in the subject is to determine or estimate the smallest number f so that any complete graph with edges labelled by the elements of Z_q contains a subgraph of a prescribed type in which the total weight of the edges is 0 modulo q . This problem for complete directed graphs, where the desired subgraph is a directed cycle, is addressed in Lemma 3.1 and Lemma 3.2. It seems plausible to believe that the first lemma is not tight and that the function $g(q)$ in Theorem 2 is linear in q for any integer q .

Acknowledgment

The initial results in this paper were obtained when the second author visited the Department of Mathematics of Princeton University. He would like to thank the department for the hospitality.

References

- [1] N. Alon and N. Linial, *Cycles of length 0 modulo k in directed graphs*, J. Combin. Th. Ser. B 47 (1989), 114–119.

- [2] N. Alon, P. D. Seymour and R. Thomas, *A separator theorem for non-planar graphs*, J. Amer. Math. Soc. 3 (1990), 801-808.
- [3] B. Bollobás, *Cycles modulo k* , Bull. London Math. Soc. 9 (1977), 97–98.
- [4] Y. Caro, *Zero-sum problems — a survey*, Discrete Math. 152 (1996), 93–113.
- [5] N. Draganić, M. Krivelevich and R. Nenadov, *The size Ramsey number of short subdivisions*, Random Struct. Alg., to appear.
- [6] L. Friedman and M. Krivelevich, *Cycle lengths in expanding graphs*, Combinatorica 41 (2021), 53–74.
- [7] J. Gao, Q. Huo, C.-H. Liu and J. Ma, *A unified proof of conjectures on cycle lengths in graphs*, Int. Math. Res. Not., to appear.
- [8] K. Kawarabayashi and B. Reed, *A separator theorem in minor-closed classes*, Proc. 51st Symp. Found. Comp. Sci. (FOCS'10), 2010, 153–162.
- [9] Y. Kohayakawa, T. Retter and V. Rödl, *The size Ramsey number of short subdivisions of bounded degree graphs*, Random Struct. Alg. 54 (2019), 304–339.
- [10] M. Krivelevich, *Expanders — how to find them, and what to find in them*, Surveys in Combinatorics, A. Lo et al., Eds., London Math. Soc. Lecture Notes 456, pp. 115–142, 2019.
- [11] H. Liu and R. Montgomery, *A solution to Erdős and Hajnal’s odd cycle problem*, arXiv: 2010.15802.
- [12] C. Thomassen, *Graph decomposition with applications to subdivisions and path systems modulo k* , J. Graph Theory 7 (1983), 261–271.