

A Proof of Kamp's theorem

Alexander Rabinovich

The Blavatnik School of Computer Science, Tel Aviv University,
email: rabinoa@post.tau.ac.il

Abstract

We provide a simple proof of Kamp's theorem.

1998 ACM Subject Classification F.4.1 Temporal Logic

Keywords and phrases Temporal Logic, Monadic Logic, Expressive Completeness

1 Introduction

Temporal Logic (*TL*) introduced to Computer Science by Pnueli in [8] is a convenient framework for reasoning about “reactive” systems. This has made temporal logics a popular subject in the Computer Science community, enjoying extensive research in the past 30 years. In *TL* we describe basic system properties by *atomic propositions* that hold at some points in time, but not at others. More complex properties are expressed by formulas built from the atoms using Boolean connectives and *Modalities* (temporal connectives): A k -place modality M transforms statements $\varphi_1, \dots, \varphi_k$ possibly on ‘past’ or ‘future’ points to a statement $M(\varphi_1, \dots, \varphi_k)$ on the ‘present’ point t_0 . The rule to determine the truth of a statement $M(\varphi_1, \dots, \varphi_k)$ at t_0 is called a *truth table* of M . The choice of particular modalities with their truth tables yields different temporal logics. A temporal logic with modalities M_1, \dots, M_k is denoted by $TL(M_1, \dots, M_k)$.

The simplest example is the one place modality $\diamond X$ saying: “ X holds some time in the future.” Its truth table is formalized by $\varphi_\diamond(t_0, X) := (\exists t > t_0)X(t)$. This is a formula of the First-Order Monadic Logic of Order (*FOMLO*) - a fundamental formalism in Mathematical Logic where formulas are built using atomic propositions $P(t)$, atomic relations between elements $t_1 = t_2$, $t_1 < t_2$, Boolean connectives and first-order quantifiers $\exists t$ and $\forall t$. Two more natural modalities are the modalities *Until* (“*Until*”) and *Since* (“*Since*”). $X\text{Until}Y$ means that X will hold from now until a time in the future when Y will hold. $X\text{Since}Y$ means that Y was true at some point of time in the past and since that point X was true until (not necessarily including) now. Both modalities have truth tables in *FOMLO*. Most modalities used in the literature are defined by such *FOMLO* truth tables, and as a result, every temporal formula translates directly into an equivalent *FOMLO* formula. Thus, the different temporal logics may be considered as a convenient way to use fragments of *FOMLO*. *FOMLO* can also serve as a yardstick by which one is able to check the strength of temporal logics: A temporal logic is *expressively complete* for a fragment L of *FOMLO* if every formula of L with a single free variable t_0 is equivalent to a temporal formula.

Actually, the notion of expressive completeness refers to a temporal logic and to a model (or a class of models), since the question whether two formulas are equivalent depends on the domain over which they are evaluated. Any (partially) ordered set with monadic predicates is a model for *TL* and *FOMLO*, but the main, *canonical*, linear time intended models are the non-negative integers $\langle \mathbb{N}, < \rangle$ for discrete time and the reals $\langle \mathbb{R}, < \rangle$ for continuous time.

Kamp's theorem [7] states that the temporal logic with modalities *Until* and *Since* is



© Alexander Rabinovich;

licensed under Creative Commons License ND

Leibniz International Proceedings in Informatics

LIPIC Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

expressively complete for *FOMLO* over the above two linear time canonical¹ models.

This seminal theorem initiated the whole study of expressive completeness, and it remains one of the most interesting and distinctive results in temporal logic; very few, if any, similar ‘modal’ results exist. Several alternative proofs of it and stronger results have appeared; none of them are trivial (at least to most people) [6].

The objective of this paper is to provide a simple proof of Kamp's theorem.

The rest of the paper is organized as follows: In Section 2 we recall the definitions of the monadic logic, the temporal logics and state Kamp's theorem. Section 3 introduces formulas in a normal form and states their simple properties. In Section 4 we prove Kamp's theorem. The proof of one proposition is postponed to Section 5. Section 6 comments on the previous proofs of Kamp's theorem.

2 Preliminaries

In this section we recall the definitions of the first-order monadic logic of order, the temporal logics and state Kamp's theorem.

Fix a set Σ of *atoms*. We use $P, Q, R, S \dots$ to denote members of Σ . The syntax and semantics of both logics are defined below with respect to such Σ .

2.1 First-Order Monadic Logic of Order

Syntax: In the context of *FOMLO*, the atoms of Σ are referred to (and used) as *unary predicate symbols*. Formulas are built using these symbols, plus two binary relation symbols: $<$ and $=$, and a set of first-order variables (denoted: x, y, z, \dots). Formulas are defined by the grammar:

$$\text{atomic} ::= x < y \mid x = y \mid P(x) \quad (\text{where } P \in \Sigma)$$

$$\varphi ::= \text{atomic} \mid \neg\varphi_1 \mid \varphi_1 \vee \varphi_2 \mid \varphi_1 \wedge \varphi_2 \mid \exists x\varphi_1 \mid \forall x\varphi_1$$

We will also use the standard abbreviated notation for **bounded quantifiers**, e.g., $(\exists x)_{>z}(\dots)$ denotes $\exists x((x > z) \wedge (\dots))$, and $(\forall x)^{<z}(\dots)$ denotes $\forall x((x < z) \rightarrow (\dots))$, and $((\forall x)_{>z_1}^{<z_2}(\dots))$ denotes $\forall x((z_1 < x < z_2) \rightarrow (\dots))$, etc.

Semantics. Formulas are interpreted over *labeled linear orders* which are called *chains*. A Σ -*chain* is a triplet $\mathcal{M} = (T, <, \mathcal{I})$ where T is a set - the *domain* of the chain, $<$ is a linear order relation on T , and $\mathcal{I} : \Sigma \rightarrow \mathcal{P}(T)$ is the *interpretation* of Σ (where \mathcal{P} is the powerset notation). We use the standard notation $\mathcal{M}, t_1, t_2, \dots, t_n \models \varphi(x_1, x_2, \dots, x_n)$ to indicate that the formula φ with free variables among x_1, \dots, x_n is satisfiable in \mathcal{M} when x_i are interpreted as elements t_i of \mathcal{M} . For atomic $P(x)$ this is defined by: $\mathcal{M}, t \models P(x)$ iff $t \in \mathcal{I}(P)$; the semantics of $<, =, \neg, \wedge, \vee, \exists$ and \forall is defined in a standard way.

2.2 TL(Until, Since) Temporal Logic

In this section we recall the syntax and semantics of a temporal logic with **strict-Until** and **strict-Since** modalities, denoted by *TL(Until, Since)*.

In the context of temporal logics, the atoms of Σ are used as *atomic propositions* (also called *propositional atoms*). Formulas of *TL(Until, Since)* are built using these atoms,

¹ the technical notion which unifies $\langle \mathbb{N}, < \rangle$ and $\langle \mathbb{R}, < \rangle$ is Dedekind completeness.

Boolean connectives and *strict-Until* and *strict-Since* binary modalities. The formulas are defined by the grammar:

$$F ::= \text{True} \mid P \mid \neg F_1 \mid F_1 \vee F_2 \mid F_1 \wedge F_2 \mid F_1 \text{Until} F_2 \mid F_1 \text{Since} F_2,$$

where $P \in \Sigma$.

Semantics. Formulas are interpreted at *time-points* (or *moments*) in chains (elements of the domain). The semantics of $TL(\text{Until}, \text{Since})$ formulas is defined inductively: Given a chain $\mathcal{M} = (T, <, \mathcal{I})$ and $t \in T$, define when a formula F holds in \mathcal{M} at t - denoted $\mathcal{M}, t \models F$:

- $\mathcal{M}, t \models P$ iff $t \in \mathcal{I}(P)$, for any propositional atom P .
- $\mathcal{M}, t \models F \vee G$ iff $\mathcal{M}, t \models F$ or $\mathcal{M}, t \models G$; similarly for \wedge and \neg .
- $\mathcal{M}, t \models F_1 \text{Until} F_2$ iff there is $t' > t$ such that $\mathcal{M}, t' \models F_2$ and $\mathcal{M}, t_1 \models F_1$ for all $t_1 \in (t, t')$.
- $\mathcal{M}, t \models F_1 \text{Since} F_2$ iff there is $t' < t$ such that $\mathcal{M}, t' \models F_2$ and $\mathcal{M}, t_1 \models F_1$ for all $t_1 \in (t', t)$.

We will use standard abbreviations. As usual $\Box F$ (respectively, $\overleftarrow{\Box} F$) is an abbreviation for $\neg(\text{TrueUntil}(\neg F))$ (respectively, $\neg(\text{TrueSince}(\neg F))$), and $\mathbf{K}^+(F)$ (respectively, $\mathbf{K}^-(F)$) is an abbreviation for $\neg((\neg F)\text{UntilTrue})$ (respectively, $\neg((\neg F)\text{SinceTrue})$).

1. $\Box F$ (respectively, $\overleftarrow{\Box} F$) holds at t iff F holds everywhere after (respectively, before) t .
2. $\mathbf{K}^-(F)$ holds at a moment t iff $t = \sup(\{t' \mid t' < t \text{ and } F \text{ holds at } t'\})$.
3. $\mathbf{K}^+(F)$ holds at a moment t iff $t = \inf(\{t' \mid t' > t \text{ and } F \text{ holds at } t'\})$.

Note that $\mathbf{K}^+(\text{True})$ (respectively, $\mathbf{K}^-(\text{True})$) holds at t in \mathcal{M} if t is a right limit (respectively, a left limit) point of the underlining order. In particular, both $\mathbf{K}^+(\text{True})$ and $\mathbf{K}^-(\text{True})$ are equivalent to False in the chains over $(\mathbb{N}, <)$,

2.3 Kamp's Theorem

Equivalence between temporal and monadic formulas is naturally defined: F is equivalent to $\varphi(x)$ over a class \mathcal{C} of structures iff for any $\mathcal{M} \in \mathcal{C}$ and $t \in \mathcal{M}$: $\mathcal{M}, t \models F \Leftrightarrow \mathcal{M}, t \models \varphi(x)$. If \mathcal{C} is the class of all chains, we will say that F is equivalent to φ .

A linear order $(T, <)$ is *Dedekind complete* if for every non-empty subset S of T , if S has a lower bound in T then it has a greatest lower bound, written $\inf(S)$, and if S has an upper bound in T then it has a least upper bound, written $\sup(S)$. The order of the naturals and of the reals are Dedekind complete, while the order of the rationals is not Dedekind complete. A chain is Dedekind complete if its underlying linear order is Dedekind complete.

The fundamental theorem of Kamp's states that $TL(\text{Until}, \text{Since})$ is expressively equivalent to $FOMLO$ over Dedekind complete chains.

- **Theorem 2.1 (Kamp [7]).** 1. Given any $TL(\text{Until}, \text{Since})$ formula A there is a $FOMLO$ formula $\varphi_A(x)$ which is equivalent to A over all chains.
- 2. Given any $FOMLO$ formula $\varphi(x)$ with one free variable, there is a $TL(\text{Until}, \text{Since})$ formula which is equivalent to φ over Dedekind complete chains.

The meaning preserving translation from $TL(\text{Until}, \text{Since})$ to $FOMLO$ is easily obtained by structural induction. The contribution of our paper is a proof of Theorem 2.1 (2). The proof is constructive. An algorithm which for every every $FOMLO$ formula $\varphi(x)$ constructs a $TL(\text{Until}, \text{Since})$ formula which is equivalent to φ over Dedekind complete chains is easily extracted from our proof. However, this algorithm is not efficient in the sense of complexity theory. This is unavoidable because there is a non-elementary succinctness gap between $FOMLO$ and $TL(\text{Until}, \text{Since})$ even over the class of finite chains, i.e., for every $m, n \in \mathbb{N}$ there is a $FOMLO$ formula $\varphi(x_0)$ of size $|\varphi| > n$ which is not equivalent (even over finite chains) to

any $TL(\text{Until}, \text{Since})$ formula of size $\leq \exp(m, |\varphi|)$, where the m -iterated exponential function $\exp(m, n)$ is defined by induction on m so that $\exp(1, n) = 2^n$, and $\exp(m+1, n) = 2^{\exp(m, n)}$.

3 $\vec{\exists}\forall$ formulas

First, we introduce $\vec{\exists}\forall$ formulas which are instances of the Decomposition formulas of [3].

► **Definition 3.1** ($\vec{\exists}\forall$ -formulas). Let Σ be a set of monadic predicate names. An $\vec{\exists}\forall$ -formula over Σ is a formula of the form:

$$\begin{aligned} \psi(z_0, \dots, z_m) := & \exists x_n \dots \exists x_1 \exists x_0 \\ & \left(\bigwedge_{k=0}^m z_k = x_{i_k} \right) \wedge (x_n > x_{n-1} > \dots > x_1 > x_0) && \text{“ordering of } x_i \text{ and } z_j\text{”} \\ & \wedge \bigwedge_{j=0}^n \alpha_j(x_j) && \text{“Each } \alpha_j \text{ holds at } x_j\text{”} \\ & \wedge \bigwedge_{j=1}^n [(\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y)] && \text{“Each } \beta_j \text{ holds along } (x_{j-1}, x_j)\text{”} \\ & \wedge (\forall y)_{>x_n} \beta_{n+1}(y) && \text{“}\beta_{n+1} \text{ holds everywhere after } x_n\text{”} \\ & \wedge (\forall y)^{<x_0} \beta_0(y) && \text{“}\beta_0 \text{ holds everywhere before } x_0\text{”} \end{aligned}$$

with a prefix of $n+1$ existential quantifiers and with all α_j, β_j quantifier free formulas with one variable over Σ . (ψ has $m+1$ free variables z_0, \dots, z_m and $m+1 \leq n+1$ existential quantifiers are dummy and are introduced just in order to simplify notations.)

It is clear that

- **Lemma 3.2. 1.** Conjunction of $\vec{\exists}\forall$ -formulas is equivalent to a disjunction of $\vec{\exists}\forall$ -formulas.
- 2. Every $\vec{\exists}\forall$ -formula is equivalent to a conjunction of $\vec{\exists}\forall$ -formulas with at most two free variables.
- 3. For every $\vec{\exists}\forall$ -formula φ the formula $\exists x\varphi$ is equivalent to a $\vec{\exists}\forall$ -formula.

► **Definition 3.3** ($\vee\vec{\exists}\forall$ -formulas). A formula is a $\vee\vec{\exists}\forall$ formula if it is equivalent to a disjunction of $\vec{\exists}\forall$ -formulas.

► **Lemma 3.4** (closure properties). The set of $\vee\vec{\exists}\forall$ formulas is closed under disjunction, conjunction, and existential quantification.

Proof. By (1) and (3) of Lemma 3.2, and distributivity of \exists over \vee . ◀

The set of $\vee\vec{\exists}\forall$ formulas is not closed under negation². However, we show later (see Proposition 4.3) that the negation of a $\vee\vec{\exists}\forall$ formula is equivalent to a $\vee\vec{\exists}\forall$ formula in the expansion of the chains by all $TL(\text{Until}, \text{Since})$ definable predicates.

The $\vee\vec{\exists}\forall$ formulas with one free variable can be easily translated to $TL(\text{Until}, \text{Since})$.

► **Proposition 3.5** (From $\vee\vec{\exists}\forall$ -formulas to $TL(\text{Until}, \text{Since})$ formulas). Every $\vee\vec{\exists}\forall$ -formula with one free variable is equivalent to a $TL(\text{Until}, \text{Since})$ formula.

² The truth table of $P\text{Until}Q$ is an $\vec{\exists}\forall$ formula $(\exists x')_{>x}(Q(x') \wedge (\forall y)_{\leq x'}^{<x} P(y))$, yet we can prove that its negation is not equivalent to any $\vee\vec{\exists}\forall$ formula.

Proof. By a simple formalization we show that every $\vec{\exists}\forall$ -formula with one free variable is equivalent to a $TL(\text{Until}, \text{Since})$ formula. This immediately implies the proposition.

Let $\psi(z_0)$ be an $\vec{\exists}\forall$ -formula

$$\begin{aligned} & \exists x_n \dots \exists x_1 \exists x_0 z_0 = x_k \wedge (x_n > x_{n-1} > \dots > x_1 > x_0) \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \\ & \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{<x_0} \beta_0(y) \wedge (\forall y)_{>x_n} \beta_{n+1}(y) \end{aligned}$$

Let A_i and B_i be temporal formulas equivalent to α_i and β_i (A_i and B_i do not even use Until and Since modalities). It is easy to see that ψ is equivalent to the conjunction of

$$A_k \wedge (B_{k+1} \text{Until} (A_{k+1} \wedge (B_{k+2} \text{Until} \dots (A_{n-1} \wedge (B_n \text{Until} (A_n \wedge \Box B_{n+1}))))))$$

and

$$A_k \wedge (B_{k-1} \text{Since} (A_{k-1} \wedge (B_{k-2} \text{Since} (\dots A_1 \wedge (B_1 \text{Since} (A_0 \wedge \Box B_0)))))) \quad \blacktriangleleft$$

4 Proof of Kamp's theorem

The next definition plays a major role in the proof of both Kamp's and Stavi's theorems [3].

► **Definition 4.1.** Let \mathcal{M} be a Σ chain. We denote by $\mathcal{E}[\Sigma]$ the set of unary predicate names $\Sigma \cup \{A \mid A \text{ is an } TL(\text{Until}, \text{Since})\text{-formula over } \Sigma\}$. The canonical $TL(\text{Until}, \text{Since})$ -expansion of \mathcal{M} is an expansion of \mathcal{M} to an $\mathcal{E}[\Sigma]$ -chain, where each predicate name $A \in \mathcal{E}[\Sigma]$ is interpreted as $\{a \in \mathcal{M} \mid \mathcal{M}, a \models A\}$ ³. We say that first-order formulas in the signature $\mathcal{E}[\Sigma] \cup \{<\}$ are equivalent over \mathcal{M} (respectively, over a class of Σ -chains \mathcal{C}) if they are equivalent in the canonical expansion of \mathcal{M} (in the canonical expansion of every $\mathcal{M} \in \mathcal{C}$).

Note that if A is a $TL(\text{Until}, \text{Since})$ formula over $\mathcal{E}[\Sigma]$ predicates, then it is equivalent to a $TL(\text{Until}, \text{Since})$ formula over Σ , and hence to an atomic formula in the canonical $TL(\text{Until}, \text{Since})$ -expansions.

From now on we say that “formulas are equivalent in a chain \mathcal{M} ” instead of “formulas are equivalent in the canonical $TL(\text{Until}, \text{Since})$ -expansion of \mathcal{M} .” The $\vec{\exists}\forall$ and $\forall\vec{\exists}\forall$ formulas are defined as previously, but now they can use as atoms $TL(\text{Until}, \text{Since})$ definable predicates.

It is clear that all the results stated above hold for this modified notion of $\forall\vec{\exists}\forall$ formulas. In particular, every $\forall\vec{\exists}\forall$ formula with one free variable is equivalent to an $TL(\text{Until}, \text{Since})$ formula, and the set of $\forall\vec{\exists}\forall$ formulas is closed under conjunction, disjunction and existential quantification. However, now the set of $\forall\vec{\exists}\forall$ formulas is also closed under negation, due to the next proposition whose proof is postponed to Sect. 5.

► **Proposition 4.2.** (Closure under negation) The negation of $\vec{\exists}\forall$ -formulas with at most two free variables is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ -formulas.

As a consequence we obtain

► **Proposition 4.3.** Every first-order formula is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ -formulas.

Proof. We proceed by structural induction.

³ We often use “ $a \in \mathcal{M}$ ” instead of “ a is an element of the domain of \mathcal{M} ”

Atomic It is clear that every atomic formula is equivalent to a disjunction of (even quantifier free) $\vec{\exists}\forall$ -formulas.

Disjunction - immediate.

Negation If φ is an $\vec{\exists}\forall$ -formula, then by Lemma 3.2(2) it is equivalent to a conjunction of $\vec{\exists}\forall$ formulas with at most two free variables. Hence, $\neg\varphi$ is equivalent to a disjunction of $\neg\psi_i$ where ψ_i are $\vec{\exists}\forall$ -formulas with at most two free variables. By Proposition 4.2, $\neg\psi_i$ is equivalent to a disjunction of $\vec{\exists}\forall$ formulas γ_i^j . Hence, $\neg\varphi$ is equivalent to a disjunction $\bigvee_i \bigvee_j \gamma_i^j$ of $\vec{\exists}\forall$ formulas.

If φ is a disjunction of $\vec{\exists}\forall$ formulas φ_i , then $\neg\varphi$ is equivalent to the conjunction of $\neg\varphi_i$. By the above, $\neg\varphi_i$ is equivalent to a $\bigvee\vec{\exists}\forall$ formula. Since, $\bigvee\vec{\exists}\forall$ formulas are closed under conjunction (Lemma 3.4), we obtain that $\neg\varphi$ is equivalent to a disjunction of $\vec{\exists}\forall$ formulas.

\exists -quantifier For \exists -quantifier, the claim follows from Lemma 3.4. \blacktriangleleft

Now, we are ready to prove Kamp's Theorem:

► Theorem 4.4. For every *FOMLO* formula $\varphi(x)$ with one free variable, there is a *TL(Until, Since)* formula which is equivalent to φ over Dedekind complete chains.

Proof. By Proposition 4.3, $\varphi(x)$ is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ formulas $\varphi_i(x)$. By Proposition 3.5, $\varphi_i(x)$ is equivalent to a *TL(Until, Since)* formula. Hence, $\varphi(x)$ is equivalent over Dedekind complete chains to a *TL(Until, Since)* formula. \blacktriangleleft

5 Proof of Proposition 4.2

Let $\psi(z_0, z_1)$ be an $\vec{\exists}\forall$ -formula

$$\begin{aligned} & \exists x_n \dots \exists x_1 \exists x_0 [z_0 = x_m \wedge z_1 = x_k \wedge (x_0 < x_1 < \dots < x_{n-1} < x_n) \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \\ & \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{<x_0} \beta_0(y) \wedge (\forall y)_{>x_n} \beta_{n+1}(y)] \end{aligned}$$

We consider two cases. In the first case $k = m$, i.e., $z_0 = z_1$ and in the second $k \neq m$.

If $k = m$, then ψ is equivalent to $z_0 = z_1 \wedge \psi'(z_0)$, where ψ' is an $\vec{\exists}\forall$ -formula. By Proposition 3.5, ψ' is equivalent to an *TL(Until, Since)* formula A' . Therefore, ψ is equivalent to an $\vec{\exists}\forall$ -formula $\exists x_0 [z_0 = x_0 \wedge z_1 = x_0 \wedge A'(x_0)]$.

If $k \neq m$, w.l.o.g. we assume that $m < k$. Hence, ψ is equivalent to a conjunction of

1. $\psi_0(z_0)$ defined as:

$$\begin{aligned} & \exists x_0 \dots \exists x_{m-1} \exists x_m [z_0 = x_m \wedge (x_0 < x_1 < \dots < x_m) \wedge \bigwedge_{j=0}^m \alpha_j(x_j) \\ & \wedge \bigwedge_{j=1}^m (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{<x_0} \beta_0(y) \end{aligned}$$

2. $\psi_1(z_1)$ defined as:

$$\begin{aligned} & \exists x_k \dots \exists x_{k+1} \exists x_n [z_1 = x_k \wedge (x_k < x_{k+1} < \dots < x_n) \wedge \bigwedge_{j=k}^n \alpha_j(x_j) \\ & \wedge \bigwedge_{j=k+1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{>x_n} \beta_{n+1}(y)] \end{aligned}$$

3. $\varphi(z_0, z_1)$ defined as:

$$\exists x_m \dots \exists x_k [(z_0 = x_m < x_{m+1} < \dots < x_k = z_1) \wedge \bigwedge_{j=m}^k \alpha_j(x_j) \\ \wedge \bigwedge_{j=m+1}^k (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y)]$$

The first two formulas are $\vec{\exists}\forall$ -formulas with one free variable. Therefore, (by Proposition 3.5) they are equivalent to a $TL(\text{Until}, \text{Since})$ formulas (in the signature $\mathcal{E}[\Sigma]$). Hence, their negations are equivalent (over the canonical expansions) to atomic (and hence to $\vec{\exists}\forall$) formulas.

Therefore, it is sufficient to show that the negation of the third formula is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ -formulas. This is stated in the the following lemma:

► **Lemma 5.1.** The negation of any formula of the form

$$\exists x_0 \dots \exists x_n [(z_0 = x_0 < \dots < x_n = z_1) \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y)] \quad (1)$$

where α_i, β_i are quantifier free, is equivalent (over Dedekind complete chains) to a disjunction of $\vec{\exists}\forall$ -formulas.

In the rest of this section we prove Lemma 5.1.

First, we introduce some helpful notations.

► **Notations 5.2.** We use the abbreviated notation $[\alpha_0, \beta_1 \dots, \alpha_{n-1}, \beta_n \alpha_n](z_0, z_1)$ for the $\vec{\exists}\forall$ -formula as in (1).

In this notation Lemma 5.1 can be rephrased as $\neg[\alpha_0, \beta_1 \dots, \alpha_{n-1}, \beta_n \alpha_n](z_0, z_1)$ is equivalent (over Dedekind complete chains) to a $\vee \vec{\exists}\forall$ formula.

We start with the instance of Lemma 5.1 where all β_i are True.

► **Lemma 5.3.** $\neg \exists x_1 \dots \exists x_n (z_0 < x_1 < \dots < x_n < z_1) \wedge \bigwedge_{i=1}^n P_i(x_i)$ is equivalent over Dedekind complete chains to a $\vee \vec{\exists}\forall$ formula $O_n(P_1, \dots, P_n, z_0, z_1)$.

Proof. We proceed by induction.

Basis: $\neg(\exists x_1)_{>z_0}^{<z_1} P_1(x_1)$ is equivalent to $(\forall y)_{>z_0}^{<z_1} \neg P_1(y)$.

Inductive step: $n \mapsto n+1$. We assume that a $\vee \vec{\exists}\forall$ formula O_n was defined and construct a $\vee \vec{\exists}\forall$ formula O_{n+1} .

Observe that if the interval (z_0, z_1) is non-empty, then one of the following cases holds:

Case 1 There is no occurrence of P_1 in (z_0, z_1) , i.e. $(\forall y)_{>z_0}^{<z_1} \neg P_1(y)$. In this case $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ should be equivalent to True.

Case 2 If case 1 does not hold then let $r_0 = \inf\{z \in (z_0, z_1) \mid P_1(z)\}$ (such r_0 exists by Dedekind completeness. Note that $r_0 = z_0$ iff $\mathbf{K}^+(P_1)(z_0)$. If $r_0 > z_0$ then $r_0 \in (z_0, z_1)$ and r_0 is definable by the following $\vee \vec{\exists}\forall$ formula:

$$\text{INF}(z_0, r_0, z_1, P_1) := z_0 < r_0 < z_1 \wedge (\forall y)_{>z_0}^{<r_0} \neg P_1(y) \wedge \\ \wedge (P_1(r_0) \vee \mathbf{K}^+(P_1)(r_0)) \quad (2)$$

Subcase $r_0 = z_0$ In this subcase $O_n(P_2, \dots, P_n, z_0, z_1)$ and $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ should be equivalent.

Subcase $r_0 \in (z_0, z_1)$ In this subcase $O_n(P_2, \dots, P_n, r_0, z_1)$ and $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ should be equivalent.

Hence, $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ can be defined as the disjunction of “ (z_0, z_1) is empty” and the following formulas:

1. $(\forall y)_{> z_0}^{< z_1} \neg P_1(y)$
2. $\mathbf{K}^+(P_1)(z_0) \wedge O_n(P_2, \dots, P_n, z_0, z_1)$
3. $(\exists r_0)_{> z_0}^{< z_1} (INF(z_0, r_0, z_1, P_1) \wedge O_n(P_2, \dots, P_n, r_0, z_1))$

The first formula is a $\forall \vec{\exists} \forall$ formula. By the inductive assumptions O_n is a $\forall \vec{\exists} \forall$ formula. $\mathbf{K}^+(P_1)(z_0)$ is an atomic (and hence a $\forall \vec{\exists} \forall$) formula in the canonical expansion, and $INF(z_0, r_0, z_1, P_1)$ is a $\forall \vec{\exists} \forall$ formula. Since $\forall \vec{\exists} \forall$ formulas are closed under conjunction, disjunction and the existential quantification, we conclude that O_{n+1} is a $\forall \vec{\exists} \forall$ formula. ◀

As a consequence we obtain

- **Corollary 5.4. 1.** $\neg(\exists z)_{> z_0}^{< z_1} [\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z)$ is equivalent over Dedekind complete chains to a $\forall \vec{\exists} \forall$ formula.
2. $\neg(\exists z)_{> z_0}^{< z_1} [\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z, z_1)$ is equivalent over Dedekind complete chains to a $\forall \vec{\exists} \forall$ formula.

Proof. (1) Define

$$F_n := \alpha_n$$

$$F_{i-1} := \alpha_{i-1} \wedge \beta_i \mathbf{Until} F_i \quad \text{for } i = 1, \dots, n$$

Observe that there is $z \in (z_0, z_1)$ such that $[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z)$ iff $F_0(z_0)$ and there is an increasing sequence $x_1 < \dots < x_n$ in an open interval (z_0, z_1) such that $F_i(x_i)$ for $i = 1, \dots, n$. Indeed, the direction \Rightarrow is trivial. The direction \Leftarrow is easily proved by induction.

The *basis* is trivial.

Inductive step: $n \mapsto n + 1$. Assume $F_0(z_0)$ holds and that (z_0, z_1) contains an increasing sequence $x_1 < \dots < x_{n+1}$ such that $F_i(x_i)$ for $i = 1, \dots, n + 1$. By the inductive assumption there is $y_1 \in (z_0, x_{n+1})$ such that

$$[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \beta_{n-1} \alpha_{n-1}, \beta_n, (\alpha_n \wedge \beta_{n+1} \mathbf{Until} \alpha_{n+1})](z_0, y_1).$$

In particular, y_1 satisfies $(\alpha_n \wedge \beta_{n+1} \mathbf{Until} \alpha_{n+1})$. Hence, there is $y_2 > y_1$ such that y_2 satisfies α_{n+1} and β_{n+1} holds along (y_1, y_2) .

If $y_2 \leq x_{n+1}$ then the required $z \in (z_0, z_1)$ equals to y_2 , and we are done. Otherwise, $x_{n+1} < y_2$. Therefore, $x_{n+1} \in (y_1, y_2)$ and β_{n+1} holds along (y_1, x_{n+1}) . Hence, the required z equals to x_{n+1} .

From the above observation and Lemma 5.3, it follows that $\neg F_0(z_0) \vee O_n(F_1, \dots, F_n, z_0, z_1)$ is a $\forall \vec{\exists} \forall$ formula that is equivalent to $\neg(\exists z)_{> z_0}^{< z_1} [\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z)$.

(2) is the mirror image of (1) and is proved similarly. ◀

Now we are ready to prove Lemma 5.1, i.e.,

$$\neg[\alpha_0, \beta_1, \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1) \text{ is equivalent}$$

$$\text{over Dedekind complete chains to a } \forall \vec{\exists} \forall \text{ formula.}$$

If the interval (z_0, z_1) is empty then the assertion is immediate. We assume that (z_0, z_1) is non-empty. Hence, at least one of the following cases holds:

Case 1 $\neg \alpha_0(z_0)$ or $\neg \alpha_n(z_1)$ or $\neg(\beta_1 \mathbf{Until} \alpha_1)(z_0)$ or $\neg(\beta_n \mathbf{Since} \alpha_{n-1})(z_1)$.

Case 2 $\alpha_0(z_0)$, and β_1 holds along (z_0, z_1) .

Case 3 1. $\alpha_0(z_0) \wedge (\beta_1 \text{Until} \alpha_1)(z_0)$, and

2. there is $x \in (z_0, z_1)$ such that $\neg\beta_1(x)$.

For each of these cases we construct a $\bigvee \vec{\exists} \forall$ formula $Cond_i$ which describes it (i.e., Case i holds iff $Cond_i$ holds) and show that if $Cond_i$ holds, then $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula $Form_i$. Hence, $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to $\bigvee_i [Cond_i \wedge Form_i]$ which is a $\bigvee \vec{\exists} \forall$ formula.

Case 1 This case is already explicitly described by the $\bigvee \vec{\exists} \forall$ formula (in the canonical expansion). In this case $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to True.

Case 2 This case is described by a $\bigvee \vec{\exists} \forall$ formula $\alpha_0(z_0) \wedge (\forall z)_{>z_0}^{<z_1} \beta_1$. In this case $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to “there is no $z \in (z_0, z_1)$ such that $[\alpha_1, \beta_2 \dots, \beta_n, \alpha_n](z, z_1)$.” By Corollary 5.4 this is expressible by a $\bigvee \vec{\exists} \forall$ formula.

Case 3 The first condition of Case 3 is already explicitly described by a $\bigvee \vec{\exists} \forall$ formula. When the first condition holds, then the second condition is equivalent to “there is (a unique) $r_0 \in (z_0, z_1)$ such that $r_0 = \inf\{z \in (z_0, z_1) \mid \neg\beta_1(z)\}$ ” (If $\beta_1 \text{Until} \alpha_1$ holds at z_0 and there is $x \in (z_0, z_1)$ such that $\neg\beta_1(x)$, then such r_0 exists because we deal with Dedekind complete chains.) This r_0 is definable by the following $\bigvee \vec{\exists} \forall$ formula, i.e., it is a unique z which satisfies it⁴:

$$INF^{\neg\beta_1}(z_0, z, z_1) := z_0 < z < z_1 \wedge (\forall y)_{>z_0}^{<z} \beta_1(y) \wedge (\neg\beta_1(z) \vee \mathbf{K}^+(\neg\beta_1)(z)) \quad (3)$$

Hence, Case 3 is described by $\alpha_0(z_0) \wedge (\beta_1 \text{Until} \alpha_1)(z_0) \wedge (\exists z)_{>z_0}^{<z_1} INF^{\neg\beta_1}(z_0, z, z_1)$. Since the set of $\bigvee \vec{\exists} \forall$ formulas is closed under conjunction, disjunction and \exists , this case is described by a $\bigvee \vec{\exists} \forall$ formula.

It is sufficient to show that $(\exists z)_{>z_0}^{<z_1} INF^{\neg\beta_1}(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula.

We prove this by induction on n .

The *basis* is trivial.

Inductive step $n \mapsto n + 1$.


Define:

$$\begin{aligned} A_i^-(z_0, z) &:= [\alpha_0, \beta_1, \dots, \beta_i, \alpha_i](z_0, z) & i = 1, \dots, n \\ A_i^+(z, z_1) &:= [\alpha_i, \beta_{i+1}, \dots, \beta_{n+1}, \alpha_{n+1}](z, z_1) & i = 1, \dots, n \\ A_i(z_0, z, z_1) &:= A_i^-(z_0, z) \wedge A_i^+(z, z_1) & i = 1, \dots, n \\ B_i^-(z_0, z) &:= [\alpha_0 \beta_1, \dots, \beta_{i-1}, \alpha_{i-1}, \beta_i, \beta_i](z_0, z) & i = 1, \dots, n+1 \\ B_i^+(z, z_1) &:= [\beta_i, \beta_i, \alpha_i \beta_{i+1} \alpha_{i+1}, \dots, \beta_{n+1}, \alpha_{n+1}](z, z_1) & i = 1, \dots, n+1 \\ B_i(z_0, z, z_1) &:= B_i^-(z_0, z) \wedge B_i^+(z, z_1) & i = 1, \dots, n+1 \end{aligned}$$

If the interval (z_0, z_1) is non-empty, these definitions imply

$$\begin{aligned} [\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1) &\Leftrightarrow (\forall z)_{>z_0}^{<z_1} \left(\bigvee_{i=1}^n A_i \vee \bigvee_{i=1}^{n+1} B_i \right) \\ [\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1) &\Leftrightarrow (\exists z)_{>z_0}^{<z_1} \left(\bigvee_{i=1}^n A_i \vee \bigvee_{i=1}^{n+1} B_i \right) \end{aligned}$$

⁴ We will use only existence and will not use uniqueness.



■ **Figure 1** $B_2(z_0, z, z_1) := [\alpha_0, \beta_1, \alpha_1, \beta_2, \beta_2](z_0, z) \wedge [\beta_2, \beta_2, \alpha_2, \beta_3, \alpha_3](z, z_1)$

Hence, for every φ

$$(\exists z)_{>z_0}^{\leq z_1} \varphi(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$$

is equivalent to

$$(\exists z)_{>z_0}^{\leq z_1} (\varphi(z) \wedge \bigwedge_{i=1}^n \neg A_i \wedge \bigwedge_{i=1}^{n+1} \neg B_i)$$

In particular,

$$(\exists z)_{>z_0}^{\leq z_1} INF^{\neg\beta_1}(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$$

is equivalent to

$$(\exists z)_{>z_0}^{\leq z_1} (INF^{\neg\beta_1}(z) \wedge \bigwedge_{i=1}^n \neg A_i \wedge \bigwedge_{i=1}^{n+1} \neg B_i),$$

where $INF^{\neg\beta_1}(z)$ was defined in equation (3).

By the inductive assumption

(a) $\neg A_i$ is equivalent to a $\vee \vec{\exists} \forall$ formula for $i = 1, \dots, n$.

(b) $\neg B_i$ is equivalent to a $\vee \vec{\exists} \forall$ formula for $i = 2, \dots, n$.

Recall $B_1 := B_1^- \wedge B_1^+$ and $B_{n+1} := B_{n+1}^- \wedge B_{n+1}^+$.

(c) $\neg B_1^-$ and $\neg B_{n+1}^+$ are equivalent to $\vee \vec{\exists} \forall$ formulas, by the induction basis.

(d) $INF^{\neg\beta_1}(z) \wedge \neg B_1^+(z, z_1)$ is equivalent to $INF^{\neg\beta_1}(z)$, because if $INF^{\neg\beta_1}(z)$, then for no $x > z$, β_1 holds along $[z, x)$.

(e) $INF^{\neg\beta_1}(z) \wedge \neg B_{n+1}^-(z_0, z)$ is equivalent to $INF^{\neg\beta_1}(z) \wedge (\text{"}\beta_1 \text{ holds on } (z_0, z)\text{"} \wedge \neg B_{n+1}^-(z_0, z))$. Since, by case 2, " β_1 holds on (z_0, z) " $\wedge \neg B_{n+1}^-(z_0, z)$ is equivalent to a $\vee \vec{\exists} \forall$ formula, and $INF^{\neg\beta_1}(z)$ is a $\vee \vec{\exists} \forall$ formula, we conclude that $INF^{\neg\beta_1}(z) \wedge \neg B_{n+1}^-(z_0, z)$ is equivalent to a $\vee \vec{\exists} \forall$ formula.

Since the set of $\vee \vec{\exists} \forall$ formulas is closed under conjunction, disjunction and \exists , by (a)-(e) we obtain that $(\exists z)_{>z_0}^{\leq z_1} (INF^{\neg\beta_1}(z) \wedge \bigwedge_{i=1}^n \neg A_i \wedge \bigwedge_{i=1}^{n+1} \neg B_i)$ is equivalent to a $\vee \vec{\exists} \forall$ formula.

Therefore, $(\exists z)_{>z_0}^{\leq z_1} INF^{\neg\beta_1}(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$ is also a $\vee \vec{\exists} \forall$ formula.

This completes our proof of Lemma 5.1 and of Proposition 4.2.

6 Related Works

Kamp's theorem was proved in

1. Kamp's thesis [7] (proof > 100pages).
2. Outlined by Gabbay, Pnueli, Stavi and Shelah [3] (Sect. 2) for \mathbb{N} and stated that it can be extended to Dedekind complete orders using game arguments.
3. Was proved by Gabbay [1] by separation arguments for \mathbb{N} , and extended to Dedekind complete order in [2].
4. Was proved by Hodkinson [4] by game arguments and simplified in [5] (unpublished).

A temporal logic has the *separation* property if its formulas can be equivalently rewritten as a boolean combination of formulas, each of which depends only on the past, present or future. The separation property was introduced by Gabbay [1], and surprisingly, a temporal logic which can express \Box and \Box has the separation property (over a class \mathcal{C} of structures) iff it is expressively complete for *FOMLO* over \mathcal{C} .

The separation proof for $TL(\text{Until}, \text{Since})$ over \mathbb{N} is manageable; however, over the real (and over Dedekind complete) chains it contains many rules and transformations and is not easy to follow. Hodkinson and Reynolds [6] write:

The proofs of theorems 18 and 19 [Kamp's theorem over naturals and over reals, respectively] are direct, showing that each formula can be separated. They are tough and tougher, respectively. Nonetheless, they are effective, and so, whilst not quite providing an algorithm to determine if a set of connectives is expressively complete, they do suggest a potential way of telling in practice whether a given set of connectives is expressively complete – in Gabbay's words, *try to separate and see where you get stuck!*

The game arguments are easier to grasp, but they use complicated inductive assertions. The proof in [5] proceeds roughly as follows. Let \mathcal{L}_r be the set of $TL(\text{Until}, \text{Since})$ formulas of nesting depth at most r . A formula of the form: $\exists \bar{x} \forall y \chi(\bar{x}, y, \bar{z})$ where \bar{x} is an n -tuple of variables and χ is a quantifier free formula over $\{<, =\}$ and \mathcal{L}_r -definable monadic predicates is called $\langle n, r \rangle$ -decomposition formula. The main inductive assertion is proved by “unusual back-and-forth games” and can be rephrased in logical terms as there is a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that for every $n, r \in \mathbb{N}$, the negation of positive Boolean combinations $\langle n, r \rangle$ -decomposition formula is equivalent to a positive Boolean combination of $\langle f(n), (n+r) \rangle$ -decomposition formulas.

Our proof is inspired by [3] and [5]; however, it avoids games, and it separates general logical equivalences and temporal arguments.

Many temporal formalisms studied in computer science concern only future formulas - whose truth value at any moment is determined by what happens from that moment on. A formula (temporal, or monadic with a single free first-order variable) F is (*semantically*) *future* if for every chain \mathcal{M} and moment $t_0 \in \mathcal{M}$:

$$\mathcal{M}, t_0 \models F \text{ iff } \mathcal{M}|_{\geq t_0}, t_0 \models F,$$

where $\mathcal{M}|_{\geq t_0}$ is the subchain of \mathcal{M} over the interval $[t_0, \infty)$. For example, $P\text{Until}Q$ and $\mathbf{K}^+(P)$ are future formulas, while $P\text{Since}Q$ and $\mathbf{K}^-(P)$ are not future ones.

It was shown in [3] that over the *discrete* chains Kamp's theorem holds also for *future formulas* of *FOMLO*:

► **Theorem 6.1.** Every future *FOMLO* formula is equivalent over discrete orders (Natural, Integer, finite) to a $TL(\text{Until})$ formula.

Theorems 6.1 can be easily obtained from our proof of Kamp's theorem.

The temporal logic with the modalities **Until** and **Since** is not expressively complete for *FOMLO* over the rationals. Stavi introduced two additional modalities **Until^s** and **Since^s** and proved that $TL(\text{Until}, \text{Since}, \text{Until}^s, \text{Since}^s)$ is expressively complete for *FOMLO* over all linear orders [2]. In the full version of this paper we prove Stavi's theorem. The proof is similar to our proof of Kamp's theorem; however, it treats some additional cases related to gaps in orders, and replaces $\exists\forall$ -formulas by slightly more general formulas.

Acknowledgments We are very grateful to Yoram Hirshfeld for numerous insightful discussions, and to the anonymous referees for their helpful suggestions.

References

- 1 D. Gabbay. Expressive functional completeness in tense logic (preliminary report). In U. Monnich, editor, *Aspects of Philosophical Logic*, pages 91-117. Reidel, Dordrecht, 1981.
- 2 D. Gabbay, I. Hodkinson, and M. Reynolds. *Temporal logic: Mathematical Foundations and Computational Aspects*. Oxford University Press, 1994.
- 3 D. Gabbay, A. Pnueli, S. Shelah and J. Stavi. On the Temporal Analysis of Fairness. In *POPL 1980*, pp. 163-173, 1980. *Information and Computation*, 187(2003), 196-208.
- 4 I. Hodkinson. Expressive completeness of Until and Since over dedekind complete linear time. *Modal logic and process algebra*, ed. A. Ponse, M. de Rijke, Y. Venema, *CSLI Lecture Notes 53*, 1995.
- 5 I. Hodkinson. Notes on games in temporal logic. *Lecture notes for LUATCS meeting, Johannesburg, Dec 1999*. <http://www.doc.ic.ac.uk/~imh/index.html>
- 6 I. Hodkinson and M. Reynolds. Temporal Logic Chapter 11 (pp. 655-720) in *Handbook of Modal Logic*, Patrick Blackburn, Johan van Benthem, and Frank Wolter, eds., Elsevier Science, 2006.
- 7 H. W. Kamp. Tense logic and the theory of linear order. *Phd thesis, University of California, Los Angeles, 1968*.
- 8 A. Pnueli (1977). The temporal logic of programs. In *Proc. IEEE 18th Annu. Symp. on Found. Comput. Sci.*, pages 46-57, New York, 1977.