

Walking in Circles

Noga Alon* Michal Feldman† Ariel D. Procaccia‡ Moshe Tennenholtz §

Abstract

We show that for every $x_1, \dots, x_n, y_1, \dots, y_n \in S^1$ there exists $i \in \{1, \dots, n\}$ such that $\sum_{k=1}^n d(x_i, x_k) \leq \sum_{k=1}^n d(x_i, y_k)$, where S^1 is the unit circle and d is the distance on S^1 . We also discuss a game theoretic interpretation of this result.

1 Introduction

Let $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{R}$, and denote $N = \{1, \dots, n\}$. We claim that there exists $i \in N$ such that

$$\sum_{k \in N} |x_i - x_k| \leq \sum_{k \in N} |x_i - y_k|. \quad (1)$$

To see this, assume without loss of generality that $x_1 \leq x_2 \leq \dots \leq x_n$. For every $k \in N$

$$|x_1 - x_k| + |x_n - x_k| = |x_1 - x_n| \leq |x_1 - y_k| + |x_n - y_k|,$$

and by summing over all $k \in N$ we get

$$\sum_{k \in N} |x_1 - x_k| + \sum_{k \in N} |x_n - x_k| \leq \sum_{k \in N} |x_1 - y_k| + \sum_{k \in N} |x_n - y_k|.$$

It immediately follows that (1) holds with respect to $i = 1$ or $i = n$, that is, with respect to one of the extreme points.

Next, let $x_1, \dots, x_n, y_1, \dots, y_n \in S^1$, where S^1 is the unit circle. Let $d : S^1 \times S^1 \rightarrow \mathbb{R}_+$ be the distance on S^1 , i.e., the distance between two points is the length of the shorter arc between them. If x_1, \dots, x_n cannot be placed on one semicircle then there are no longer points that can easily be identified as “extreme”. Is it still true that there exists $i \in N$ such that

$$\sum_{k \in N} d(x_i, x_k) \leq \sum_{k \in N} d(x_i, y_k)?$$

*Microsoft Israel R&D Center, 13 Shenkar Street, Herzeliya 46725, Israel, and Schools of Mathematics and Computer Science, Tel Aviv University, Tel Aviv, 69978, Israel, Email: nogaa@tau.ac.il. Research supported in part by a USA Israeli BSF grant, by an ERC advanced grant and by the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University.

†School of Business Administration and Center for the Study of Rationality, The Hebrew University of Jerusalem, Jerusalem 91904, Israel, and Microsoft Israel R&D Center. Email: mfeldman@huji.ac.il. Research supported in part by the Israel Science Foundation (grant number 1219/09) and by the Leon Recanati Fund of the Jerusalem school of business administration.

‡School of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA, Email: arielpro@seas.harvard.edu. This work was done while the author was at Microsoft Israel R&D Center.

§Microsoft Israel R&D Center, 13 Shenkar Street, Herzeliya 46725, Israel, and Technion, IIT, Haifa 32000, Israel. Email: mshet@microsoft.com

Put another way, if we have n people walking on a circle from x_1, \dots, x_n to y_1, \dots, y_n , is it true that they cannot jointly move closer (in terms of the sum of distances) to *every* starting point?

In Section 2 we answer this question in the affirmative. Although our main result is formulated with respect to S^1 , it clearly also holds for any closed curve that is homeomorphic to S^1 . On the other hand it is easy to see that it does not hold for any graph embedded in the plane that contains a vertex v of degree at least 3. This is shown by taking $n = 3$ and letting x_1, x_2 and x_3 be three points that lie on different edges incident with v , each being of equal distance from v , with $y_1 = y_2 = y_3 = v$. In Section 3 we briefly discuss a game theoretic implication of this result.

2 Main Theorem

We first introduce some notations. Let $x, y \in S^1$; we denote the shorter open arc between x and y by (x, y) , and the shorter closed arc between x and y by $[x, y]$.¹ For every $x \in S^1$ we let \hat{x} be the antipodal point of x on S^1 , i.e., the diametrically opposite point. Finally, given $x, y \in S^1$ we denote the ‘‘clockwise operator’’ by \succeq , and its strong version by \succ . Without being very formal, $x \succeq y$ means that x is clockwise of y on the circle; this operator is well-defined in the context of an arc of length at most π .

We are now ready to formulate and prove our main result.

Theorem 1. *Let $x_1, \dots, x_n, y_1, \dots, y_n \in S^1$. Then there exists $i \in N$ such that*

$$\sum_{k \in N} d(x_i, x_k) \leq \sum_{k \in N} d(x_i, y_k). \quad (2)$$

Proof. Let $x_1, \dots, x_n \in S^1$, and define a multiset X by $X = \{x_1, \dots, x_n\}$. We first note that we can assume that there are no $x_i, x_j \in X$ such that $x_j = \hat{x}_i$. Indeed, in this case the claim holds trivially with respect to either i or j , since for all $z \in S^1$,

$$d(x_i, z) + d(x_j, z) = \pi.$$

In particular, for every $x_i, x_j \in X$, (x_i, x_j) and (\hat{x}_i, \hat{x}_j) are well-defined.

We say that two points $x_i, x_j \in X$ are *nearly antipodal* if there is no point $x_k \in X$ such that $x_k \in (x_i, \hat{x}_j)$ or $x_k \in (x_j, \hat{x}_i)$; let $A \subseteq X^2$ be the set of all nearly antipodal pairs. Given a nearly antipodal pair $\langle x_i, x_j \rangle \in A$, let the *critical arc of $\langle x_i, x_j \rangle$* , denoted $\text{crit}(x_i, x_j)$, be the *long* open arc between \hat{x}_i and \hat{x}_j , that is,

$$\text{crit}(x_i, x_j) = S^1 \setminus [\hat{x}_i, \hat{x}_j] = (x_i, \hat{x}_j) \cup [x_i, x_j] \cup (x_j, \hat{x}_i).$$

See Figure 1 for an illustration of the construction given above.

Let $y_1, \dots, y_n \in S^1$, and define a multiset Y by $Y = \{y_1, \dots, y_n\}$. It is sufficient to prove that there exists a pair of nearly antipodal points $\langle x_i, x_j \rangle \in A$ such that

$$\sum_{k \in N} d(x_i, x_k) + \sum_{k \in N} d(x_j, x_k) \leq \sum_{k \in N} d(x_i, y_k) + \sum_{k \in N} d(x_j, y_k).$$

¹If x and y are antipodal then these arcs are ambiguously defined.




Figure 1: An illustration of the construction in the proof of Theorem 1, for $n = 4$. The nearly antipodal pairs are $A = \{\langle x_1, x_3 \rangle, \langle x_2, x_4 \rangle, \langle x_3, x_4 \rangle\}$.

Indeed, in this case we get that Equation (2) holds with respect to either x_i or x_j . Therefore, assume for the purpose of contradiction that for every pair of nearly antipodal points $\langle x_i, x_j \rangle \in A$,

$$\sum_{k \in N} d(x_i, x_k) + \sum_{k \in N} d(x_j, x_k) > \sum_{k \in N} d(x_i, y_k) + \sum_{k \in N} d(x_j, y_k). \quad (3)$$

We claim that Equation (3) implies that for every pair of nearly antipodal points $\langle x_i, x_j \rangle \in A$, the number of points from Y on $\text{crit}(x_i, x_j)$ is strictly greater than the number of points from X on the same arc. Formally, for $\langle x_i, x_j \rangle \in A$, let

$$\alpha_{ij}^X = |\{x_k \in X : x_k \in \text{crit}(x_i, x_j)\}|,$$

and

$$\alpha_{ij}^Y = |\{y_k \in Y : y_k \in \text{crit}(x_i, x_j)\}|.$$

We have the following claim.

Lemma 1. *Let $\langle x_i, x_j \rangle \in A$. Then $\alpha_{ij}^Y > \alpha_{ij}^X$.*

Proof. For every point $z \in [x_i, x_j]$, we have that

$$d(x_i, z) + d(x_j, z) = d(x_i, x_j).$$

Let $d'(x_i, x_j)$ be the length of the longer arc $S^1 \setminus [x_i, x_j]$ between x_i and x_j , namely

$$d'(x_i, x_j) = d(x_i, \hat{x}_j) + d(\hat{x}_j, \hat{x}_i) + d(\hat{x}_i, x_j) = d(x_i, x_j) + 2 \cdot d(x_i, \hat{x}_j) > d(x_i, x_j).$$

For every point $z \in [\hat{x}_i, \hat{x}_j]$ it holds that

$$d(x_i, z) + d(x_j, z) = d'(x_i, x_j).$$

Finally, it holds that for every $z \in (x_i, \hat{x}_j) \cup (x_j, \hat{x}_i)$,

$$d(x_i, x_j) < d(x_i, z) + d(x_j, z) < d'(x_i, x_j).$$

Since x_i and x_j are nearly antipodal, there are no points from X in (x_i, \hat{x}_j) and (x_j, \hat{x}_i) . Therefore,

$$\sum_{k \in N} d(x_i, x_k) + \sum_{k \in N} d(x_j, x_k) = \alpha_{ij}^X \cdot d(x_i, x_j) + (n - \alpha_{ij}^X) \cdot d'(x_i, x_j). \quad (4)$$

On the other hand,

$$\sum_{k \in N} d(x_i, y_k) + \sum_{k \in N} d(x_j, y_k) \geq \alpha_{ij}^Y \cdot d(x_i, x_j) + (n - \alpha_{ij}^Y) \cdot d'(x_i, x_j). \quad (5)$$

Using Equations (4) and (5), we get that (3) directly implies that $\alpha_{ij}^Y > \alpha_{ij}^X$, as claimed. \square

From Lemma 1, we immediately get that

$$\sum_{(x_i, x_j) \in A} \alpha_{ij}^X < \sum_{(x_i, x_j) \in A} \alpha_{ij}^Y. \quad (6)$$

In order to derive a contradiction, we also need the following lemma.

Lemma 2. *There exists $r \in \mathbb{N}$ such that*

$$\sum_{(x_i, x_j) \in A} \alpha_{ij}^X = r \cdot n, \quad (7)$$

whereas

$$\sum_{(x_i, x_j) \in A} \alpha_{ij}^Y \leq r \cdot n, \quad (8)$$

Proof. It is easy to see that $|A|$ is odd (e.g., by induction on n); let $|A| = 2s + 1$, for some $s \in \mathbb{N}$. We first wish to claim that every $x_i \in X$ is a member of exactly $s + 1$ critical arcs, which directly proves Equation (7) with $r = s + 1$.

Without loss of generality we prove the claim with respect to $x_1 \in X$. Consider the clockwise closed arc between x_1 and \hat{x}_1 . Let $Z = \{z_1, \dots, z_t\}$ be all the points x_i or \hat{x}_i on this arc, where for all k , $z_{k+1} \succeq z_k$. In particular, $z_1 = x_1$ and $z_t = \hat{x}_1$. For instance, in Figure 1 we have that $Z = \{x_1, x_2, \hat{x}_4, x_3, \hat{x}_1\}$.

Now, we have that the set of nearly-antipodal pairs A is exactly the set of pairs $\langle x_i, x_j \rangle$ such that z_k is a point x_i and z_{k+1} is an antipodal point \hat{x}_j (this is a *type 1* nearly-antipodal pair), or z_k is an antipodal point \hat{x}_i and z_{k+1} is a point x_j (this is a *type 2* nearly-antipodal pair). If $\langle x_i, x_j \rangle$ is a nearly-antipodal pair of type 1, we have that $x_1 \in [x_i, x_j]$, and hence $x_1 \in \text{crit}(x_i, x_j)$. On the other hand, if $\langle x_i, x_j \rangle$ is a nearly-antipodal pair of type 2, then $x_1 \notin \text{crit}(x_i, x_j)$. Since $z_1 = x_1$ is a point from X and $z_t = \hat{x}_1$ is an antipodal point, the number of nearly-antipodal pairs of type 1 is exactly $s + 1$, which proves the claim.

In order to prove Equation (8), let $y \in S^1$. It is sufficient to prove that there exists $x_i \in X$ such that y appears in at most as many critical arcs as x_i , since we already know that x_i is a member of exactly $s + 1$ critical arcs. We consider the two points or antipodal points that are adjacent to y , and briefly examine four cases.

1. $x_i \preceq y \preceq x_j$: y appears in exactly the critical arcs that contain x_i (these are also exactly the critical arcs that contain x_j).

2. $x_i \preceq y \prec \hat{x}_j$: y appears in exactly the critical arcs that contain x_i .
3. $\hat{x}_i \prec y \preceq x_j$: y appears in exactly the critical arcs that contain x_j .
4. $\hat{x}_i \preceq y \preceq \hat{x}_j$: When walking counterclockwise from \hat{x}_i , let $x_k \in X$ be the first point from X , and let \hat{x}_l be the last antipodal point such that $x_k \prec \hat{x}_l \preceq \hat{x}_i$. Then y is contained in all the critical arcs that contain x_k , except for $\text{crit}(x_k, \hat{x}_l)$, that is, in exactly s critical arcs.

We deduce that every y_i is contained in at most $r = s + 1$ critical arcs, which implies the validity of Equation (8). \square

It follows from Lemma 2 that

$$\sum_{(x_i, x_j) \in A} \alpha_{ij}^X \geq \sum_{(x_i, x_j) \in A} \alpha_{ij}^Y,$$

in contradiction to Equation (6). \square

3 A Game Theoretic Interpretation

Consider a facility location setting where the facility is to be located on a network. Each player $i \in N$ has an *ideal* location on the network; the player's *cost* is the distance between its ideal location and the location that was selected for the facility. A *mechanism* is a function that receives the reported ideal locations of the players as input, and returns the location of the facility.

From the game theoretic point of view it is desirable that mechanisms be immune to manipulation by rational players. A mechanism is *strategyproof* if players can never benefit by misreporting their ideal location, regardless of the reports of the other players. In other words, by misreporting his location a player cannot influence the facility location in a way that it becomes closer to his ideal location. Schummer and Vohra [2] establish a characterization of deterministic strategyproof facility location mechanisms on networks. In particular, they show that if the network is a circle then the only deterministic strategyproof and onto mechanism is a *dictatorship* of one of the players, i.e., given any constellation of ideal locations the mechanism selects the ideal location of a fixed player.

Randomization provides a way around this negative result. Indeed, under the *random dictator* mechanism the ideal location of one of the agents is selected uniformly at random. This mechanism is strategyproof: if an agent was chosen as the dictator then it could not have gained from lying, whereas if it was not chosen then it could not have affected the outcome. Random dictator is also “fair” compared to a deterministic dictatorship, and in particular produces an outcome that yields a good approximation to the optimal facility location in terms of minimizing the sum of players’ costs².

Taking our game theoretic requirements a step further, we say that a mechanism is *group strategyproof* if even a coalition of agents cannot all benefit by lying, that is, for every joint deviation by a coalition there is a member of the coalition whose expected distance from the facility does not decrease. Group strategyproofness is a highly desirable property, but is rarely satisfied by nontrivial mechanisms. We can derive the following result as an immediate corollary of Theorem 1.

²Specifically, it gives a $2 - 2/n$ approximation, where approximation is defined in the usual sense by looking at the worst-case ratio between the expected cost of the mechanism’s solution and the cost of the optimal solution [1].

Corollary 1. *Assume that the network is a circle. Then the random dictator mechanism is group strategyproof.*

To see this, note that we can assume without loss of generality that the deviating coalition contains all the players. Indeed, the expected cost of a player given that a nondeviating player is selected by the mechanism, and the probability that a nondeviating player is selected by the mechanism, are both independent of the reports of the deviating players. The corollary follows after scaling by a factor of $1/n$. For more details, including the formal facility location model, the reader is referred to [1].

References

- [1] N. Alon, M. Feldman, A. D. Procaccia, and M. Tennenholtz. Strategyproof approximation mechanisms for location on networks. arXiv:0907.2049, 2009.
- [2] J. Schummer and R. V. Vohra. Strategy-proof location on a network. *Journal of Economic Theory*, 104(2):405–428, 2002.