

A Characterization of the (natural) Graph Properties Testable with One-Sided Error*

Noga Alon[†] Asaf Shapira[‡]

Abstract

The problem of characterizing all the testable graph properties is considered by many to be the most important open problem in the area of property-testing. Our main result in this paper is a solution of an important special case of this general problem; Call a property tester *oblivious* if its decisions are independent of the size of the input graph. We show that a graph property \mathcal{P} has an oblivious one-sided error tester, **if and only if** \mathcal{P} is *semi hereditary*. We stress that any “natural” property that can be tested (either with one-sided or with two-sided error) can be tested by an oblivious tester. In particular, all the testers studied thus far in the literature were oblivious. Our main result can thus be considered as a precise characterization of the natural graph properties, which are testable with one-sided error.

One of the main technical contributions of this paper is in showing that any hereditary graph property can be tested with one-sided error. This general result contains as a special case all the previous results about testing graph properties with one-sided error. These include the results of [23] and [6] about testing k -colorability, the characterization of [25] of the graph-partitioning problems that are testable with one-sided error, the induced vertex colorability properties of [4], the induced edge colorability properties of [17], a transformation from two-sided to one-sided error testing [25], as well as a recent result about testing monotone graph properties [11]. More importantly, as a special case of our main result, we infer that some of the most well studied graph properties, both in graph theory and computer science, are testable with one-sided error. Some of these properties are the well known graph properties of being Perfect, Chordal, Interval, Comparability, Permutation and more. None of these properties was previously known to be testable.

1 Introduction

1.1 Definitions and background

The meta problem in the area of property testing is the following: Given a combinatorial structure S , distinguish if S satisfies some property \mathcal{P} or if S is ϵ -far from satisfying \mathcal{P} , where S is said to be

*A preliminary version of this paper appeared in the Proc. of the 46th Annual IEEE Symposium on Foundations of Computer Science (FOCS), IEEE (2005), 429-438.

[†]Schools of Mathematics and Computer Science, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel, and IAS, Princeton, NJ 08540, USA. Email: nogaa@tau.ac.il. Research supported in part by a grant from the Israel Science Foundation, by the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University, and by the Von Neumann Fund.

[‡]School of Computer Science, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel. Email: asafico@tau.ac.il. This work forms part of the author’s Ph.D. thesis. Research supported by a Charles Clore Foundation Fellowship and an IBM Ph.D. Fellowship.

ϵ -far from satisfying \mathcal{P} if an ϵ -fraction of its representation should be modified in order to make S satisfy \mathcal{P} . The main goal is to design randomized algorithms, which look at a very small portion of the input, and using this information distinguish with high probability between the above two cases. Such algorithms are called *property testers* or simply *testers* for the property \mathcal{P} . Preferably, a tester should look at a portion of the input whose size is a function of ϵ only. Blum, Luby and Rubinfeld [14] were the first to formulate a question of this type, and the general notion of property testing was first formulated by Rubinfeld and Sudan [36], who were motivated in studying various algebraic properties such as linearity of functions.

The main focus of this paper is in testing properties of graphs, where a property of graphs is simply a family of graphs closed under isomorphism. Throughout the paper we only consider decidable graph properties, that is, properties \mathcal{P} for which it is possible to decide in finite time whether an input graph G satisfies \mathcal{P} . In this case a graph G , is said to be ϵ -far from satisfying a property \mathcal{P} , if one needs to add/delete at least ϵn^2 edges to/from G in order to turn it into a graph satisfying \mathcal{P} . A tester for \mathcal{P} should distinguish with high probability, say $2/3$, between the case that G satisfies \mathcal{P} from the case that G is ϵ -far from satisfying \mathcal{P} . Here we assume that the tester can query some oracle, whether a pair of vertices, i and j , are adjacent in the input graph G . In what follows we will say that a tester for a graph property \mathcal{P} has *one-sided error* if it accepts any graph satisfying \mathcal{P} with probability 1 (and still rejects those that are ϵ -far with probability at least $2/3$). If the tester may reject graphs satisfying \mathcal{P} with non-zero probability then it is said to have *two-sided error*.

The study of the notion of testability for combinatorial structures, and mainly for labelled graphs, was introduced in the seminal paper of Goldreich, Goldwasser and Ron [23]. In this paper it was shown that many natural graph properties such as k -colorability, having a large clique and having a large cut, have a tester, whose *query complexity*, that is, the number of oracle queries of type “does (i, j) belong to $E(G)$ ”, can be upper bounded by a function that depends only on ϵ and is independent of the size of the input (the dependence on ϵ may vary between properties). In this paper we will say that properties having such efficient testers, that is, whose query complexity can be upper bounded by a function of ϵ only, are simply *testable*. Following [23], many other graph properties were shown to be testable, while others were shown to be non-testable. We note that the model of graph property testing that we study here is the so called *dense graph* model as defined in [23]. Graph property testing has also been studied in other models, such as the *bounded degree* model [24] and the *general density* model [31].

1.2 Related work

The most interesting results in property-testing are those that show that large families of problems are testable. The main result of [23] states that a certain abstract graph partition problem, which includes as special cases k -colorability, having a large cut and having a large clique, is testable. The authors of [25] gave a characterization of the partition problems discussed in [23] that are testable with one-sided error. In [4], a logical characterization of a family of testable graph properties was obtained. According to this characterization, every first order graph-property of type $\exists\forall$ (see Subsection 2.3.2) is testable, while there are first-order graph properties of type $\forall\exists$ that are not testable. These results were extended in [17]. There are also several general testability and non-testability results in other areas besides testing graph properties. In [5] it is proved that every regular language is testable. This result was extended to any read-once branching program in [30]. On the other hand, it was

proved in [20], that there are read-twice branching programs that are not-testable. The main result of [8] states that any constraint satisfaction problem is testable.

With this abundance of general testability results, a natural question is what makes a combinatorial property testable. As graphs are the most well studied combinatorial structures in the theory of computation, it is natural to consider the problem of characterizing the testable graph properties, as the most important open problem in the area of property testing. Regretfully, though, finding such a characterization remains a challenging open problem. The main result of this paper, Theorem 2, resolves an important natural special case of this open problem, which concerns property testers with one-sided error. For additional results and references on graph property-testing as well as on testing properties of other combinatorial structures, the reader is referred to [18], [22], [35] and [12].

2 The New Results

2.1 The main technical result and its immediate applications

A graph property is *hereditary* if it is closed under removal of vertices (and *not* necessarily under removal of edges). Equivalently, such properties are closed under taking induced subgraphs. The main technical result of this paper is the following:

Theorem 1 (Main Technical Result) *Every hereditary graph property is testable with one-sided error.*

As we will see later, the testing algorithms we design for a given hereditary property \mathcal{P} , simply sample a set of vertices S and accept if and only if the graph induced by S satisfies \mathcal{P} . This immediately implies that these testers have one-sided error. Of course, the main difficulty lies in proving that if the input is ϵ -far from satisfying \mathcal{P} then the graph induced by a large enough S (but only large enough as a function of ϵ) will not satisfy \mathcal{P} with high probability.

We note that besides certain partition properties such as having a large cut and having a large clique, which were proved to be testable with two-sided error in [23], essentially any graph property that was studied in the literature is hereditary. Thus, Theorem 1 combined with the graph partition problems of [23] imply the testability of (nearly) any natural graph property. To demonstrate the generality of Theorem 1, we use it to infer that many graph properties, which prior to this paper were not known to be testable, are in fact testable with one-sided error. These include the following hereditary properties:

- **Perfect Graphs:** A graph G is perfect if for every *induced* subgraph of G , denoted G' , the chromatic number of G' equals the size of the largest clique in G' .
- **Chordal Graphs:** A graph is chordal if it contains no *induced* cycle of length at least 4.
- **Interval Graphs:** A graph G on n vertices is an interval graph if there are closed intervals on the real line I_1, \dots, I_n such that $(i, j) \in E(G)$ if and only if $I_i \cap I_j \neq \emptyset$.
- **Circular-Arc Graphs:** A graph G on n vertices is a circular-arc graph if there are closed intervals on a cycle I_1, \dots, I_n such that $(i, j) \in E(G)$ if and only if $I_i \cap I_j \neq \emptyset$.

- **Comparability Graphs:** A graph G is a comparability graph if its edges can be oriented such that if there is a directed edge from i to j and from j to k , then there is one from i to k .
- **Permutation Graphs:** A graph G on n vertices is a permutation graph if there is a permutation σ of $\{1, \dots, n\}$ such that $(i, j) \in E(G)$ iff (i, j) is an inversion under σ .
- **Asteroidal Triple-Free Graphs:** G is asteroidal triple-free if it contains no independent set of 3 vertices such that each pair is joined by a path that avoids the neighborhood of the third.
- **Split Graphs:** G is a split graph if $V(G)$ can be split into a clique and an independent set.

Another abstract family of hereditary graph properties, which have been extensively studied, are the so called *intersection graph properties*. In this case we fix a certain “type” T of sets and say that a graph G on n vertices has the intersection property defined by T , if there are n sets S_1, \dots, S_n of type T , such that vertices i and j are connected in G if and only if $S_i \cap S_j \neq \emptyset$. For example, the property of being a d -Box (see [16] and its references) is obtained by letting the “type” of the sets be axis parallel boxes in R^d . See the monograph [29] for more information and examples of intersection graph properties.

It is clear that the above surveyed properties are some of the most well-studied properties in graph-theory as well as in theoretical and applied computer-science. These properties also arise naturally in Chemistry, Biology, Social Sciences, Statistics as well as in many other areas. See [26], [29], [32] and their references, where other hereditary properties and their applications are also discussed.

To further convey the reader of the power of Theorem 1 we mention that Lemma 4.2, which is the main result needed to obtain Theorem 1, immediately implies, for example, that for every ϵ there is $c = c(\epsilon)$, such that if a graph G is ϵ -far from being Chordal then G contains an **induced** cycle of length at most c , and that similar results hold for any other hereditary property. This is non-trivial as it is not clear a priori that there is no graph that is, say, $\frac{1}{100}$ -far from being Chordal and yet contains only induced cycles of length at least, say, $\Omega(\log n)$. Put in other way, if G has the property that all its induced subgraphs of size $c = c(\epsilon)$ are chordal, then G is ϵ -close to being Chordal. This gives a strong connection between the *local* properties of a graph and its *global* properties. In fact, we can show that an analogous result holds for any graph property, see Theorem 6.

2.2 The main result: oblivious testing with one-sided error

By a result of [4] and [25], it is possible to assume that a property tester works by making its queries non-adaptively. In other words, the tester first picks a random subset of vertices S , then queries all pairs (i, j) $i, j \in S$ and then continues without making additional queries. Inspecting previous results on property-testing, motivates the following notion of a slightly more restricted tester, which works while being “oblivious” to the size of the input¹.

Definition 2.1 (Oblivious Tester) *A tester (one-sided or two-sided) is for a property \mathcal{P} is said to be oblivious if it works as follows: given ϵ the tester computes an integer $Q = Q(\epsilon)$ and asks an oracle for a subgraph induced by a set of vertices S of size Q , where the oracle chooses S randomly*

¹The tester implied by the results of [25] and [4] may use the size of the input in order to determine both the query complexity and in order to make its decisions

and uniformly from the vertices of the input graph. If Q is larger than the size of the input graph then the oracle returns the entire graph. The tester then accepts or rejects (possibly randomly) according to ϵ and the graph induced by S .

Note, that by insisting that the oracle chooses the set of vertices S , an oblivious tester indeed operates without knowing the size of the input, because if the tester had to choose S then it would have to know the size of the input graph in order to specify a vertex of the graph. We believe that the above definition captures the essence of property testing in the dense graph model as essentially all the testers that have been analyzed in this model were in fact oblivious, or could trivially be turned into oblivious testers. Even the testers for properties such as having an independent set of size $\frac{1}{2}n$ or a cut with at least $\frac{1}{8}n^2$ edges (see [23]), whose definition involves the size of the graph, have oblivious testers. The reason is simply that these properties can easily be expressed without using the size of the graph. For example, in order to test if a graph has a cut with at least $\frac{1}{8}n^2$ edges one can sample some $Q = Q(\epsilon)$ vertices and accept the input if and only if the graph induced on the sample has a cut of size at least $(\frac{1}{8} - \frac{\epsilon}{2})Q^2$ (of course, one needs to prove that this sampling scheme indeed works, see [23]). Another family of graph properties for which we can confine ourselves to oblivious testers is the family of hereditary properties, which is shown to be testable by an oblivious tester in the present paper. We finally note that most “applications” of property-testing (see [18] and [35]) involve testing properties of huge networks such as the Internet, whose size is anyway unknown.

Observe, that there are two restrictions that the above definition imposes on an oblivious tester. The first is that it cannot use the size of the input in order to determine the size Q , of the sample of vertices. In other words, Q is only a function of ϵ and not a function of ϵ and n . The reader should note that a tester for a testable graph property (as defined in the Section 1) may have a query complexity that is *bounded* by a function of ϵ but one that *depends* on the size of the graph (e.g. $Q(\epsilon, n) = 1/\epsilon + (-1)^n$). Though this seems like an annoying technicality, it was proved in [10] that this subtlety may have non-trivial ramifications. The second, seemingly more severe, restriction on an oblivious tester is that it cannot use the size of the input in order to make its decisions after the subgraph induced on the set S of Q vertices has been obtained. One can easily “cook” graph properties that cannot be tested by an oblivious tester. However, these properties are somewhat non-natural. One example out of many is the following property, which we denote by \mathcal{P}' : A graph on an even number of vertices satisfies \mathcal{P}' if and only if it is bipartite, while a graph on an odd number of vertices satisfies \mathcal{P}' if and only if it is triangle-free. A tester for \mathcal{P}' clearly must use the size of the input in order to make its decision regarding the graph induced by the sample.

We now turn to the main result of this paper, which gives a characterization of the graph properties that can be tested with 1-sided error by an oblivious tester. Intuitively, in order to test a property with 1-sided error the tester must “find” some kind of proof that the input does not satisfy the property. Of course the graph itself is such a proof, but as we confine ourselves to testers whose number of queries is independent of the size of the input, the tester must find a *small* proof of this fact. For hereditary properties, such proofs exist, and are in fact (relatively) abundant. These are small induced subgraph that do not satisfy the property. In fact, this is the main idea behind our algorithm for testing hereditary properties. See Lemma 4.2, which is the main technical result of this paper.

A natural question is if other non-hereditary properties have such small proofs. For example, having a clique of size $\frac{1}{2}n$ obviously does not have such small proofs. The reason is that for any fixed graph C there are graphs that contain C as an induced subgraph and have a clique of size $\frac{1}{2}n$, and

graphs that contain C as an induced subgraph and are far from having a clique of size $\frac{1}{2}n$. In [25] it was shown that when considering the partition-problems of [23], which contain the clique property as a special case, then non-hereditary partition properties cannot be tested with 1-sided error. For general properties the situation is much more involved. However, considering only oblivious testers enables us to precisely characterize the graph properties, which are testable with one-sided error. To state this characterization we need the following definition:

Definition 2.2 (Semi-Hereditary) *A graph property \mathcal{P} is semi-hereditary if there exists a hereditary graph property \mathcal{H} such that the following holds:*

1. *Any graph satisfying \mathcal{P} also satisfies \mathcal{H} .*
2. *There is a function $M : (0, 1) \mapsto \mathbb{N}$, such that any graph G of size at least $M(\epsilon)$, which is ϵ -far from satisfying \mathcal{P} , contains an induced subgraph that does not satisfy \mathcal{H} .*

Comment 2.3 *As \mathcal{H} is hereditary, an equivalent and simpler condition equivalent to item 2 above is that G itself does not satisfy \mathcal{H} . However, the statement above will be more convenient for the proof of Theorem 2.*

Clearly, any hereditary graph property \mathcal{P} is also semi-hereditary because we can take \mathcal{H} in the above definition to be \mathcal{P} itself. In simple words, a semi-hereditary \mathcal{P} is obtained by taking a hereditary graph property \mathcal{H} , and removing from it a (possibly infinite) set of graphs. This means that the first item in Definition 2.2 is satisfied. As there are graphs not satisfying \mathcal{P} that do satisfy \mathcal{H} these graphs do not contain any induced subgraph that does not satisfy \mathcal{H} (because \mathcal{H} is hereditary). The only restriction, which is needed in order to get item 2 in Definition 2.2, is that \mathcal{P} will be such that for any $\epsilon > 0$ there will be only finitely many graphs that are ϵ -far from satisfying it, and yet contain no induced subgraph that does not satisfy \mathcal{H} .

We are now ready to state the main result of this paper.

Theorem 2 (Main Result) *A graph property \mathcal{P} has an oblivious one-sided error tester if and only if \mathcal{P} is semi-hereditary.*

Returning to the graph property \mathcal{P}' discussed above, note that by Theorem 1 this property, which is not semi-hereditary, can be tested with one-sided error by a non-oblivious tester. Therefore, it is not the case that a graph property is testable if and only if it is semi-hereditary. However, if we disregard this and other non-natural graph properties then we may assume that in order to test them we can confine ourselves to oblivious testers. Theorem 2 can thus be considered as a *precise characterization* of the “natural” graph properties, which are testable with one-sided error. We believe that it may be very interesting to further study property-testing via the framework of oblivious testers, see Section 7.

Theorems 1 and 2 suggest many questions, some of which we discuss and resolve in the following subsections, while others are discussed in Section 7 and are left as interesting open problems.

2.3 Additional results

2.3.1 On the (im)possibility of relaxing the notion of property-testing

Theorem 1 implies that any hereditary graph property is testable, when one uses the standard notion of ϵ -far as defined in Section 1. Suppose we forbid addition of edges and define a graph G on n vertices to be ϵ -far_{del} from satisfying property \mathcal{P} if one needs to delete from G at least ϵn^2 edges in order to turn it into a graph satisfying \mathcal{P} . We say that property \mathcal{P} is testable_{del} if there is a tester for distinguishing between graphs satisfying \mathcal{P} from those that are ϵ -far_{del} from satisfying it, whose number of queries can be upper bounded by a function of ϵ . A natural question is which graph properties are testable_{del}. Obviously, any hereditary property, which is also closed under removal of edges (such as k -colorability) is testable_{del} as in these cases being ϵ -far_{del} is equivalent to ϵ -far. The following theorem is a sharp contrast to Theorems 1 and 2.

Theorem 3 *For any hereditary property \mathcal{P} , which is not satisfied by all graphs, and is satisfied by any complete graph, there is a constant $\delta = \delta(\mathcal{P}) > 0$ such that testing_{del} property \mathcal{P} (even with two-sided error) requires n^δ queries.*

Note, that any natural hereditary property, such as any of those discussed in Subsection 2.1, is satisfied by any complete graph, thus the above result applies to these properties. We briefly mention that we can also prove a similar statement when one allows only edge additions. See Section 6.

2.3.2 Unbounded first order graph properties

A first order graph property is one involving the boolean operators \wedge, \vee, \neg , the \forall, \exists quantifiers, the equality operator $=$, and the adjacency relation \sim . For example, the triangle-freeness property can be written as $\forall v_1, v_2, v_3 \neg(v_1 \sim v_2 \wedge v_2 \sim v_3 \wedge v_1 \sim v_3)$. The main result of [4] states that every first order graph property without quantification $\forall\exists$ is testable (possibly with two-sided error). The main tool in [4] was a theorem stating that any hereditary graph property, which is expressible in terms of a *finite* family of forbidden induced subgraphs is testable. Theorem 1 is a powerful extension of this result as it allows the family of forbidden induced subgraphs to be infinite. One may thus ask whether Theorem 1 can be used in order to extend the result of [4]. Theorem 4 below gives a positive answer to this question. To state this extension we need the following definition.

Definition 2.4 (Unbounded First-Order Properties of type $\exists\forall$) *An unbounded first order graph property of type $\exists\forall$ is of the form*

$$\exists x_1, \dots, x_t \bigwedge_{i=1}^{\infty} \forall y_1, \dots, y_i A_i(x_1, \dots, x_t, y_1, \dots, y_i) \quad (1)$$

where each $A_i(x_1, \dots, x_t, y_1, \dots, y_i)$ is a quantifier-free first order expression.

The main result of [4] states that any graph property that can be expressed as above while using a *single* relation A_i is testable. Using the main techniques of this paper, we can extend this to expressions containing *infinitely* many expressions A_i .

Theorem 4 *Every graph property describable by an unbounded first order graph property of type $\exists\forall$ is testable (possibly with two-sided error).*

It should be noted that it is proved in [4] that there are first order graph properties with alternation of type $\forall\exists$ which are not testable, thus Theorem 4 is in some sense best possible.

2.3.3 ANDing hereditary graph properties

We next describe a consequence of Theorem 1 (in fact, of the main step of proving Theorem 1), which does not assert the testability of some graph property, but rather one that may be useful in the general study of graph property testing. Suppose $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots\}$ is a (possibly infinite) set of monotone graph properties, that is, properties that are closed under removal of vertices and edges. It was proved in [11] that in this case there is a function $\delta : (0, 1) \mapsto (0, 1)$ such that if a graph G is ϵ -far from satisfying all the properties of \mathcal{P} then for some i it is also $\delta(\epsilon)$ -far from satisfying \mathcal{P}_i . Note, that this statement is non-trivial only when \mathcal{P} is infinite, as if \mathcal{P} contains k properties we can clearly take $\delta(\epsilon) = \epsilon/k$ (in fact, to get the case of finite k the properties only need to be closed under removal of edges). Consider now the case when the properties are assumed to be hereditary properties, which are not necessarily monotone. Now it is not at all clear that a similar statement holds even for $k = 2$, as modifying a graph in order to turn it into a graph satisfying \mathcal{P}_1 may increase its distance from satisfying \mathcal{P}_2 . Using Theorem 1 we can show that a similar result holds even for infinite sets of properties.

Theorem 5 *For any (possibly infinite) set of hereditary graph properties $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots\}$, there is a function $\delta_{\mathcal{P}} : (0, 1) \mapsto (0, 1)$ with the following property: If a graph G is ϵ -far from satisfying all the properties of \mathcal{P} , then for some i , the graph G is $\delta_{\mathcal{P}}(\epsilon)$ -far from satisfying \mathcal{P}_i .*

2.3.4 An extremal result for all graph property

Confirming a conjecture of Erdős, it was shown in [33] that if a graph is ϵ -far from being k -colorable, then it contains a non k -colorable subgraph of size that depends only on ϵ . In [11] this result was extended to any monotone graph property. As we have alluded to in Subsection 2.1, the main technical result of this paper, Lemma 4.2, immediately implies that this result can be extended to any hereditary graph property. In fact, we can show that a similar result holds for *any* graph property.

Theorem 6 *For every graph property \mathcal{P} , there is a function $W_{\mathcal{P}}(\epsilon)$ with the following property: If G is ϵ -far from satisfying \mathcal{P} , then G contains an **induced** subgraph of size at most $W_{\mathcal{P}}(\epsilon)$, which does not satisfy \mathcal{P} .*

2.4 Comparison to previous results

We next survey the previous results on graph property-testing, which were shown to be testable with one-sided error. As all these properties are hereditary, their testability with one-sided error follows as a special case of Theorem 1.

- **H -free:** For every fixed graph H let \mathcal{P}_H be the property of not containing a copy of H , and let \mathcal{P}_H^* be the property of not containing an induced copy of H . The property \mathcal{P}_H was (implicitly) shown to be testable in [3], and \mathcal{P}_H^* was shown to be testable in [4].

- **k -colorability:** The k -colorability property was (implicitly) shown to be testable already in [33]. In [23], a simplified explicit tester was studied with a significantly better query complexity. This result was further improved by [6].
- **Induced vertex colorability:** The main technical step in the proof of the main result of [4] was in showing that for every *finite* set of k -colored graphs \mathcal{K} , one can test the property of a graph being vertex k -colorable with no induced colored graph from the set \mathcal{K} . Note, that any such property is hereditary
- **Induced edge colorability:** Following [4], further induced edge-colorability properties were studied in [17]. In this case we have a *finite* set of k -edge-colored graphs \mathcal{K} , and the property defined by \mathcal{K} is that of having a k -edge-coloring with no induced colored graph from the set \mathcal{K} . Note, that any such property is hereditary, and that by Theorem 1 we can even take \mathcal{K} to be an infinite family of edge-colored graphs.
- **Graph partition problems:** One of the main results of [23] is that any graph-partition problem is testable with *two-sided* error. A characterization of the graph-partition properties that are testable with one-sided error was obtained in [25]. This characterization (essentially) follows as a special case of Theorem 2, as what it (implicitly) states is that a partition problem is testable with one-sided error if and only if it is hereditary.
- **Monotone graph properties:** Very recently, the authors have shown in [11] that any monotone graph property is testable with one-sided error (a graph-property is monotone if it is closed under removal of vertices *and* edges, therefore, any monotone property is in particular hereditary). Though this family of graph properties is very general and contains many interesting graph properties such as k -colorability, being H -free and certain Ramsey properties, it fails to include many interesting hereditary non-monotone properties such as those that were discussed in Subsection 2.1.
- **One-sided vs. two-sided testers:** The first author has shown ([25], Appendix D) that if a hereditary graph property is testable with two-sided error then it is also testable with one-sided error (but not necessarily with the same query complexity). By Theorem 1, this transformation becomes obsolete, as Theorem 1 directly asserts that any hereditary graph property is testable with one-sided error.
- **Bounded first order graph properties:** Theorem 4 extends the main result of [4], where the first order graph-property can contain only a single predicate A_i .

It is important to note that Theorems 1 and 2 do not assert the existence of one-sided error testers, which are as efficient as the ad-hoc testers that were designed for every specific property in the above mentioned papers. For example, the query complexity of the tester for k -colorability that follows as a special case of Theorem 1, is significantly larger than the query complexity which is guaranteed by the main result of [23] and [6]. These large bounds are obviously a consequence of the generality of Theorems 1 and 2. Furthermore, by Theorem 4 of [11], the upper bounds of Theorems 1 and 2 cannot be generally improved even for monotone graph properties. See the precise statement in [11].

2.5 Recent results

Recently, Lovász and B. Szegedy [28] found an alternative proof of Theorem 1 using the method of convergent graph sequences. Their result is slightly weaker than ours as it does not give any explicit upper bound on the query complexity of testing even simple hereditary properties such as triangle-freeness. Rödl and Schacht [34] have generalized Theorem 1 and showed that any hereditary property of k -uniform hypergraphs is testable. This proof applies variants of the recently proved hypergraph regularity lemmas. In a joint work with Fischer and Newman [2] we have managed to give a combinatorial characterization of the testable graph properties (recall that in this paper we mainly deal with one-sided error testers). This characterization implies that the regularity lemma is in some sense essential to graph property testing. Finally, the main techniques developed in this paper have been applied in another recent study [13] of the family of hereditary graph properties.

2.6 Organization

Our main tool in the proof of Theorem 1 is a novel application of a powerful variant of Szemerédi's Regularity Lemma proved in [4]. In Section 3 we introduce the basic notions of regularity and state the regularity lemmas that we use and some of their standard consequences. The proof of Theorem 1 is quite involved technically, and thus we give in Section 4 an overview of it. In this section we also prove Theorem 6. The ideas of this proof, especially the usage of the notion of colored-homomorphism, may be useful for handling other problems involving induced subgraphs. In Section 5 we give the full proof of Theorem 1 as well as the proof of Theorem 2. The proofs of Theorems 3, 4 and 5 appear in Section 6. In Section 7, we describe several possible extensions and open problems that this paper suggests. Throughout the paper, whenever we relate, for example, to a function $f_{3.1}$, we mean the function f defined in Lemma/Claim/Theorem 3.1. We would like to mention that readers that are not familiar with applications of the regularity-lemma may find it useful to refer to [11] prior to reading this paper.

3 Regularity Lemma Background

In this section we discuss the basic notions of regularity, some of the basic applications of regular partitions and state the regularity lemmas that we use in the proof of Theorem 1. See [27] for a comprehensive survey on the regularity-lemma. We start with some basic definitions. For every two nonempty disjoint vertex sets A and B of a graph G , we define $e(A, B)$ to be the number of edges of G between A and B . The *edge density* of the pair is defined as $d(A, B) = e(A, B)/|A||B|$.

Definition 3.1 (γ -regular pair) *A pair (A, B) is γ -regular, if for any two subsets $A' \subseteq A$ and $B' \subseteq B$, satisfying $|A'| \geq \gamma|A|$ and $|B'| \geq \gamma|B|$, the inequality $|d(A', B') - d(A, B)| \leq \gamma$ holds.*

A very useful lemma that we use in this paper is Lemma 3.2 below, which helps us find many induced copies of some fixed graph F , whenever a family of vertex sets are pairwise regular “enough” and their densities correspond to the edge-set of F . Several versions of this lemma were previously proved in papers using the regularity lemma. See [4] for one such proof.

Lemma 3.2 *For every real $0 < \eta < 1$ and integer $f \geq 1$ there exist $\gamma = \gamma_{3.2}(\eta, f)$ and $\delta = \delta_{3.2}(\eta, f)$ with the following property. Suppose U_1, \dots, U_f is an f -tuple of disjoint vertex sets of a graph G*

such that for every $1 \leq i < j \leq f$ the pair (U_i, U_j) is γ -regular. Let F be a graph on f vertices v_1, \dots, v_f such that whenever $(v_i, v_j) \in E(F)$ we have $d(U_i, U_j) \geq \eta$, and whenever $(v_i, v_j) \notin E(F)$ we have $d(U_i, U_j) \leq 1 - \eta$. Then, at least $\delta \prod_{i=1}^f |U_i|$ of the f -tuples $u_1 \in U_1, \dots, u_f \in U_f$ span an **induced** copy of F , where each u_i plays the role of v_i .

Comment 3.3 *Observe, that the functions $\gamma_{3.2}(\eta, f)$ and $\delta_{3.2}(\eta, f)$ may and will be assumed to be monotone non-increasing in f . Also, for ease of future definitions (in particular the one given in (4)) we set $\gamma_{3.2}(\eta, 0) = \delta_{3.2}(\eta, 0) = 1$ for any $0 < \eta < 1$.*

Note, that in terms of regularity, Lemma 3.2 requires all the pairs (U_i, U_j) to be γ -regular. However, and this will be very important later in the paper, the requirements in terms of density are not very restrictive. In particular, if $\eta \leq d(U_i, U_j) \leq 1 - \eta$ then we don't care if (i, j) is an edge of F .

A partition $\mathcal{A} = \{V_i \mid 1 \leq i \leq k\}$ of the vertex set of a graph is called an *equipartition* if $|V_i|$ and $|V_j|$ differ by no more than 1 for all $1 \leq i < j \leq k$ (so in particular each V_i has one of two possible sizes). The Regularity Lemma of Szemerédi can be formulated as follows.

Lemma 3.4 ([37]) *For every m and $\epsilon > 0$ there exists a number $T = T_{3.4}(m, \epsilon)$ with the following property: Any graph G on $n \geq T$ vertices, has an equipartition $\mathcal{A} = \{V_i \mid 1 \leq i \leq k\}$ of $V(G)$ with $m \leq k \leq T$, for which all pairs (V_i, V_j) , but at most $\epsilon \binom{k}{2}$ of them, are ϵ -regular.*

The function $T_{3.4}(m, \epsilon)$ may and is assumed to be monotone nondecreasing in m and monotone non-increasing in ϵ . Another lemma, which will be very useful in this paper is Lemma 3.5 below. Some versions of this lemma appear in various papers applying the Regularity Lemma. See [4] for one such proof.

Lemma 3.5 *For every l and γ there exists $\delta = \delta_{3.5}(l, \gamma)$ such that for every graph G with $n \geq \delta^{-1}$ vertices there exist disjoint vertex sets W_1, \dots, W_l satisfying:*

1. $|W_i| \geq \delta n$.
2. All $\binom{l}{2}$ pairs are γ -regular.
3. Either all pairs have densities at least $\frac{1}{2}$, or all pairs have densities less than $\frac{1}{2}$.

Comment 3.6 *Observe, that the function $\delta_{3.5}(l, \gamma)$ may and will be assumed to be monotone non-increasing in l and monotone non-decreasing in γ . Therefore, for ease of future applications we will assume that for all l and γ we have $\delta_{3.5}(l, \gamma) \leq 1/2$.*

Our main tool in the proof of Theorem 1 in addition to Lemmas 3.2 and 3.5 is Lemma 3.8 below, proved in [4]. This lemma can be considered a variant of the standard regularity lemma, where one can use a function that defines ϵ as a function of the size of the partition, rather than having to use a fixed ϵ as in Lemma 3.4. We denote such functions by \mathcal{E} throughout the paper. To state the lemma we need the following definition.

Definition 3.7 (The function $W_{\mathcal{E}, m}$) *Let $\mathcal{E}(r) : \mathbb{N} \mapsto (0, 1)$ be an arbitrary monotone non-increasing function, and m be an arbitrary positive integer. We define the function $W_{\mathcal{E}, m} : \mathbb{N} \mapsto (0, 1)$ inductively as follows: $W_{\mathcal{E}, m}(1) = T_{3.4}(m, \mathcal{E}(0))$. For any integer $i > 1$ put $R = W_{\mathcal{E}, m}(i - 1)$ and define*

$$W_{\mathcal{E}, m}(i) = T_{3.4}(R, \mathcal{E}(R)/R^2). \quad (2)$$

Lemma 3.8 ([4]) *For every integer m and monotone non-increasing function $\mathcal{E} : \mathbb{N} \mapsto (0, 1)$ define*

$$S = S_{3.8}(m, \mathcal{E}) = W_{\mathcal{E}, m}(100/\mathcal{E}(0)^4).$$

For any graph $G = (V, E)$ on $n \geq S$ vertices, there exists an equipartition $\mathcal{A} = \{V_i \mid 1 \leq i \leq k\}$ of $V(G)$ as well as an equipartition $\mathcal{B} = \{U_i \mid 1 \leq i \leq k\}$ of a subset of vertices $U \subseteq V(G)$, which satisfy the following:

1. $m \leq k \leq S$.
2. $U_i \subseteq V_i$ for all $i \geq 1$, and $|U_i| \geq n/S$.
3. In the equipartition \mathcal{B} , all pairs are $\mathcal{E}(k)$ -regular.
4. All but at most $\mathcal{E}(0) \binom{k}{2}$ of the pairs $1 \leq i < j \leq k$ are such that $|d(V_i, V_j) - d(U_i, U_j)| < \mathcal{E}(0)$.

Comment 3.9 *For technical reasons (see the proof in [4]), Lemma 3.8 requires that for any $r > 0$ the function $\mathcal{E}(r)$ will satisfy $\mathcal{E}(r) \leq \min\{\mathcal{E}(0)/4, 1/4r^2\}$. However, we can always assume wlog that \mathcal{E} satisfies this condition because if it does not, then we can apply Lemma 3.8 with \mathcal{E}' which is defined as $\mathcal{E}'(r) = \min\{\mathcal{E}(r), \mathcal{E}(0)/4, 1/4r^2\}$. We will thus disregard this technicality.*

The main power of Lemma 3.8 is that for *any* function \mathcal{E} it allows us to find k sets of vertices U_1, \dots, U_k of size $\Omega(n)$ such that all pairs (U_i, U_j) are $\mathcal{E}(k)$ -regular. Note, that in Lemma 3.4 we have no “control” on the relation between the number of sets and the regularity measure between them, as we first fix the regularity measure ϵ , and then get via the lemma k sets of vertices, where k can be very large in terms of ϵ . In Lemma 3.8 k can also be very large but the lemma guarantees that so will be the regularity measure between the sets, via the function \mathcal{E} .

One of the difficulties in the proof of Theorem 2, is in showing that all the constants that are used in the course of the proof can be upper bounded by functions depending on ϵ only. The following observation will thus be useful.

Proposition 3.10 *If m is bounded by a function of ϵ only then for any $\mathcal{E} : \mathbb{N} \mapsto (0, 1)$, the integer $S = S_{3.8}(m, \mathcal{E})$ can be upper bounded by a function of ϵ only².*

It should be noted that the dependency of the function $T_{3.4}(m, \epsilon)$ on ϵ is a tower of exponents of height polynomial in $1/\epsilon$ (see the proof in [27]). Thus, even for moderate functions \mathcal{E} the integer S has a huge dependency on ϵ , which is a tower of towers of exponents of height polynomial in $1/\epsilon$.

One of the main results of [4] is that for every *finite* set of graphs \mathcal{F} , the property of not containing any member of \mathcal{F} as an induced subgraph can be tested with one-sided error and with query complexity depending on ϵ only. The proof technique in [4], which applies Lemmas 3.2, 3.5 and 3.8 critically relies on the fact that the family of graphs is finite. The main step in the proof of Theorem 1 is in extending the above to *infinite* families of graphs. To this end, we use the main idea of [11], as well as a new type of homomorphism, in order to prove this stronger result. As in [11] the main idea of the proof is to apply Lemma 3.8 with a suitable function $\mathcal{E}(r)$. However, as it turns out, dealing with hereditary properties, which are not necessarily monotone, is considerably more involved. The techniques we apply in the next section, in particular the notion of colored-homomorphism, may be useful in dealing with other problems involving induced subgraphs.

²In our application of Lemma 3.8 the function $\mathcal{E}(r)$ will (implicitly) depend on the error parameter ϵ and on the fixed property \mathcal{P} being tested. For example, we will set $\mathcal{E}(r) = f(r, \mathcal{P}, \epsilon)$ for some function f . However, that will not change the fact that for a fixed property \mathcal{P} , the integer $S_{3.8}(m, \mathcal{E})$ can be bounded from above by a function of ϵ only.

4 Overview of the Proof of Theorem 1

The proof of Theorem 1 is very technical and rather long and appears in its entirety in Section 5. In this section we try to give an overview of the proof, while keeping out most of the (unnecessary) technical details. We start with an equivalent formulation of Theorem 1. To this end we introduce a convenient way of handling hereditary graph properties.

Definition 4.1 (Forbidden Induced Subgraphs) *For a hereditary graph property \mathcal{P} , define $\mathcal{F} = \mathcal{F}_{\mathcal{P}}$ to be the set of graphs which are minimal with respect to not satisfying property \mathcal{P} . In other words, a graph F belongs to \mathcal{F} if it does not satisfy \mathcal{P} , but any graph obtained from F by removing a vertex, satisfies \mathcal{P} .*

For a (possibly infinite) family of graph \mathcal{F} , a graph G is said to be *induced \mathcal{F} -free* if it contains no induced copy of any graph $F \in \mathcal{F}$. Note, that for any hereditary graph property \mathcal{P} there is a family of graphs $\mathcal{F} = \mathcal{F}_{\mathcal{P}}$ such that a graph satisfies \mathcal{P} if and only if it is induced \mathcal{F} -free. For \mathcal{F} one can simply take the family of forbidden induced subgraphs as in Definition 4.1. For example, when \mathcal{P} is the property of being Chordal (see Subsection 2.1) then $\mathcal{F}_{\mathcal{P}}$ is the set of cycles of length at least 4. As another example note that if \mathcal{P} is the property of being bipartite then $\mathcal{F}_{\mathcal{P}}$ is the family of odd cycles. Observe, that \mathcal{F} may contain *infinitely* many graphs. Clearly for any family \mathcal{F} , the property of being induced \mathcal{F} -free is hereditary, thus, the hereditary graph properties are precisely the graph properties, which are equivalent to being induced \mathcal{F} -free for some family \mathcal{F} . For ease of presentation, it will be more convenient to derive Theorem 1 from the following (essentially equivalent³) lemma, whose proof is the main technical step in this paper.

Lemma 4.2 (Main Technical Result) *For every (possibly infinite) family of graphs \mathcal{F} , there are functions $N_{\mathcal{F}}(\epsilon)$, $f_{\mathcal{F}}(\epsilon)$ and $\delta_{\mathcal{F}}(\epsilon)$ such that the following holds for any $\epsilon > 0$: If a graph G on $n \geq N_{\mathcal{F}}(\epsilon)$ vertices is ϵ -far from being induced \mathcal{F} -free, then G contains δn^f induced copies of a graph $F \in \mathcal{F}$ of size f , where $f \leq f_{\mathcal{F}}(\epsilon)$ and $\delta \geq \delta_{\mathcal{F}}(\epsilon)$.*

Before continuing with the overview of the proof of Theorem 1, we note that the above lemma immediately implies Theorem 6. Indeed, given any graph property \mathcal{P} let \mathcal{F} be the family of graphs not satisfying \mathcal{P} . Observe, that if a graph is ϵ -far from satisfying \mathcal{P} then it is also ϵ -far from being induced \mathcal{F} -free and thus by Lemma 4.2 it contains an induced subgraph $F \in \mathcal{F}$ of size at most $f_{\mathcal{F}}(\epsilon)$, and by our choice of \mathcal{F} the graph F does not satisfy \mathcal{P} . Therefore, as the function $W_{\mathcal{P}}(\epsilon)$ in the statement of Theorem 6 we can take the function $f_{\mathcal{F}}(\epsilon)$.

4.1 Homomorphism and colored homomorphism

For the proof of Lemma 4.2 we will need a new type of homomorphism, which is suitable for handling induced subgraph.

Definition 4.3 (Colored-Homomorphism) *Let K be a complete graph whose vertices are colored black or white, and whose edges are colored black, white or grey (neither the vertex coloring nor the edge coloring is assumed to be proper in the standard sense). A colored-homomorphism from a graph F to a graph K is a mapping $\varphi : V(F) \mapsto V(K)$, which satisfies the following:*

³See Section 5 for a discussion about the subtle difference due to the possible necessity of testing some properties non-uniformly.

1. If $(u, v) \in E(F)$ then either $\varphi(u) = \varphi(v) = t$ and t is colored black, or $\varphi(u) \neq \varphi(v)$ and $(\varphi(u), \varphi(v))$ is colored black or grey.
2. If $(u, v) \notin E(F)$ then either $\varphi(u) = \varphi(v) = t$ and t is colored white, or $\varphi(u) \neq \varphi(v)$ and $(\varphi(u), \varphi(v))$ is colored white or grey.

If there is a colored-homomorphism from a graph F to a colored complete graph K , we write for brevity $F \mapsto_c K$. Some explanation is in place as to the meaning of the colors in the above definition. To this end, it is instructive to compare the definition of a colored-homomorphism to the standard notion of homomorphism.

Definition 4.4 (Homomorphism) *A homomorphism from a graph F to a graph K is a mapping $\varphi : V(F) \mapsto V(K)$, which maps edges to edges, namely $(v, u) \in E(F)$ implies $(\varphi(v), \varphi(u)) \in E(K)$.*

For brevity, we denote by $F \mapsto K$ the fact that there is a homomorphism from F to K . The fact that $F \mapsto K$, simply means that we can partition the vertex set of F into $k = |V(K)|$ subsets V_1, \dots, V_k , such that each V_i is edgeless and if $(i, j) \notin E(K)$ then none of the vertices of F that belong to V_i is connected to any of the vertices of F that belong to V_j . In particular, note that $F \mapsto K_k$ if and only if F is k -colorable (where K_k is a clique of size k). The standard notion of homomorphism is sufficient for dealing with not necessarily induced subgraphs as was carried out in [11]. The reason is that having a homomorphism to a graph K is “closed under removal of vertices and edges” in the sense that if $F \mapsto K$ and F' is a subgraph of F then $F' \mapsto K$. When one wants to handle *induced* subgraphs it soon turns out that standard homomorphism is not sufficient as it does not supply enough information about F . The reason for that is that a standard homomorphism has no requirement about the non-edges of the graph. Returning to the colored-homomorphism from Definition 4.3, suppose we interpret the colors of K as follows: A white edge of K represents a non-edge, a black edge of K represents an existing edge and a grey edge represents a “don’t care.” As for the vertex colors, we think of a black vertex as a complete graph, and a white vertex as an edgeless graph. Thus, the fact that $F \mapsto_c K$ where K is a colored complete graph of size k is equivalent to the following: There is a partition of $V(F)$ into k subsets V_1, \dots, V_k such that each V_i is either complete or edgeless, where V_i is complete if $i \in V(K)$ is black and edgeless if $i \in V(K)$ is white. Also, if (i, j) is colored white then none of the vertices of F that belong to V_i is connected to any of the vertices of F that belong to V_j . Similarly, if (i, j) is colored black then all the vertices of F that belong to V_i are connected to all the vertices of F that belong to V_j . Finally, if (i, j) is colored grey then there is no restriction on pairs $(v \in V_i, u \in V_j)$ (or in our “formal” notation, we “don’t care” if $(v \in V_i, u \in V_j)$ is an edge of F). It is clear that a colored-homomorphism carries a lot more information about the structure of F than a standard homomorphism.

Our definition of colored-homomorphism should also be thought of with Lemma 3.2 in mind. Note, that in this lemma we only require $d(U_i, U_j) \geq \eta$ when $(i, j) \in E(F)$ and $d(U_i, U_j) \leq 1 - \eta$ when $(i, j) \notin E(F)$. In particular, if $\eta \leq d(U_i, U_j) \leq 1 - \eta$ then we “don’t care” whether $(i, j) \in E(F)$. In fact, as the details of the proof of Lemma 4.2 reveal, the possibility of having grey edges in the coloring of K in the definition of the colored-homomorphism is unavoidable (at least in our proof). Note, that as far as Lemma 3.2 is concerned, we only need the edge coloring in the colored-homomorphism. The details below supply some explanation for the need of the vertex coloring.

We now turn to discuss the relation between the standard regularity lemma (Lemma 3.4), the stronger regularity lemma (Lemma 3.8) and colored-homomorphism. We stress that some of the

explanations we give below are not completely accurate, and are given in order to explain the main ideas of the proof. The formal proof appears in Section 5. Given $\epsilon > 0$ and a graph G , Lemma 3.4 returns an equipartition of $V(G)$ of size k . Let the *regularity graph* of G , denoted $R = R(G)$, be the following graph. R is a graph on k vertices, where vertices i and j are connected if and only if (V_i, V_j) is a dense regular pair (with the appropriate parameters). In some sense, the regularity graph is an approximation of the original graph, up to ϵn^2 modifications. This approximation was good enough when considering monotone properties in [11] (this notion of regularity graph is standard when applying Lemma 3.4) but it is not good enough when dealing with induced graphs, which is the case we consider here. The reason is that R only approximates the dense pairs of the equipartition, while it carries no restriction or information on the sparse pairs in this equipartition. This is somewhat analogous to the fact that standard homomorphism is not good enough for dealing with induced subgraphs. Just like we defined colored-homomorphism we introduce *colored regularity graphs* as follows; Let R be a complete graph on k vertices. Color (i, j) black if (V_i, V_j) is a very dense pair, white if (V_i, V_j) is a very sparse pair, and grey if (V_i, V_j) is neither very dense nor very sparse (we omit the precise definition of “very”). Note, that a colored regularity graph carries a lot more information about G . Note also how this definition relates to a colored-homomorphism.

4.2 Proof strategy

Our overall strategy for the proof of Lemma 4.2 can be described as follows: given a graph G , which is ϵ -far from being induced \mathcal{F} -free, we will construct another “well structured” graph \tilde{G} by making less than ϵn^2 edge modifications. This will guarantee that \tilde{G} spans an induced copy of some $F \in \mathcal{F}$. One of the key ingredients of the proof will be that as G is close to \tilde{G} and \tilde{G} is “well structured” we will be able to infer that G also contains a copy of some graph $F' \in \mathcal{F}$ in a way that will allow us to infer that G actually contains *many* induced copies of some (possibly other) $F'' \in \mathcal{F}$. For simplicity, in the following overview, we will briefly argue how to find many induced copies of some graph $F \in \mathcal{F}$ in the graph \tilde{G} rather than G . The way to argue why G must also contain many copies of some $F \in \mathcal{F}$ is that the densities of the a partition of G and \tilde{G} are very similar. See the next section for the full details.

Suppose a graph G is ϵ -far from being induced \mathcal{F} -free. We would want to apply Lemma 3.4 on G , then construct the colored regularity graph, and then argue that if we make few (less than ϵn^2) modifications in G then the new graph \tilde{G} , contains an induced copy of a graph $F \in \mathcal{F}$. Furthermore, as we make very few changes, the colored regularity graph is also a “good” approximation of \tilde{G} . We would thus want to use Lemma 3.2, where for the sets U_1, \dots, U_f we take the clusters V_1, \dots, V_k of the equipartition in order to get that there are many induced copies of F in \tilde{G} . However, we are faced with the following two problems: (i) As \mathcal{F} may be infinite, we don’t know the size of the graph $F \in \mathcal{F}$ that we may expect to find in \tilde{G} . As Lemma 3.2 needs to know the size of F in advance, we don’t know how small a γ should we choose in order to apply Lemma 3.4. (ii) Note that Lemma 3.2 allows the copies of F to have only one vertex in each of the sets U_i . However, the induced copy of the graph $F \in \mathcal{F}$ that we may find in \tilde{G} may have many vertices in each cluster V_i . Note further, that Lemma 3.4 does not guarantee anything about the graphs induced by each V_i .

The main idea of the proof is to overcome the first problem by applying Lemma 3.8 with a suitable function \mathcal{E} that will guarantee that the partition is regular enough even for the largest graph $F \in \mathcal{F}$ we may expect to find in \tilde{G} . For the second problem we apply Lemma 3.5 on each of the clusters V_i in order to find subsets $W_{i,1}, \dots, W_{i,f} \subset V_i$. Note that by Lemma 3.2, if for all j', j'' we have

$d(W_{i,j'}, W_{i,j''}) \geq 1/2$ then $W_{i,1}, \dots, W_{i,f}$ span many cliques of size f , while if for all j', j'' we have $d(W_{i,j'}, W_{i,j''}) \leq 1/2$ then they span many independent sets of size f . This is the main reason for the vertex coloring of R , that is, we color vertex i of R black, if the sets returned by Lemma 3.5 are very dense, and white if they are sparse. We note that overcoming both problems mentioned above *simultaneously* adds another level of complication.

An important ingredient in the proof of Lemma 4.2 will be the following function. The reader should think of the graphs R considered below as the set of colored regularity graphs discussed above, and the parameter r as representing the size of R .

Definition 4.5 (The family \mathcal{F}_r) For any (possibly infinite) family of graphs \mathcal{F} , and any integer r let \mathcal{F}_r be the following set of colored complete graphs: A colored complete graph R belongs to \mathcal{F}_r if and only if it has at most r vertices and there is at least one $F \in \mathcal{F}$ such that $F \mapsto_c R$.

In the proof of Lemma 4.2, the set \mathcal{F}_r , defined above, will represent a subset of the colored regularity graphs of size at most r . Namely, those R for which there is at least one $F \in \mathcal{F}$ such that $F \mapsto_c R$. We now arrive at the key definition used in the proof of Lemma 4.2.

Definition 4.6 (The function $\Psi_{\mathcal{F}}$) For any family of graphs \mathcal{F} and integer r for which $\mathcal{F}_r \neq \emptyset$, let

$$\Psi_{\mathcal{F}}(r) = \max_{R \in \mathcal{F}_r} \min_{\{F \in \mathcal{F}: F \mapsto_c R\}} |V(F)|. \quad (3)$$

Define $\Psi_{\mathcal{F}}(r) = 0$ if $\mathcal{F}_r = \emptyset$. Therefore, $\Psi_{\mathcal{F}}(r)$ is monotone non-decreasing in r .

One of the key definitions in [11], is a function analogous to $\Psi_{\mathcal{F}}$ but with respect to standard homomorphism. In our case as well, $\Psi_{\mathcal{F}}$ is one of the main tools with which we apply Lemma 3.8. As by Lemma 3.4 we can upper bound the size of the regularity graph R , we can also upper bound the size of the smallest graph $F \in \mathcal{F}$ for which $F \mapsto_c R$.

As we have mentioned in the previous section, the main difficulty that prevents one from proving Theorem 1 using Lemma 3.2 is that one does not know a priori the size of the graph that one may expect to find in the equipartition. This leads us to define the following function

$$\mathcal{E}(r) = \gamma_{3.2} \left(\frac{\epsilon}{6}, \Psi_{\mathcal{F}}(r) \right) \cdot \delta_{3.5} \left(\Psi_{\mathcal{F}}(r), \gamma_{3.2} \left(\frac{\epsilon}{6}, \Psi_{\mathcal{F}}(r) \right) \right). \quad (4)$$

We next try to explain why the above defined $\mathcal{E}(r)$ when applied with Lemma 3.8 is useful in resolving the two difficulties mentioned above. Recall that r stands for the size of the regularity graph returned by Lemma 3.8. If we apply Lemma 3.8 with the above \mathcal{E} then by the first term in the definition of \mathcal{E} we know that the sets U_i (recall the statement of Lemma 3.4) are regular enough to allow one to apply Lemma 3.2 with the largest member of \mathcal{F} , which we may need to work with. This is due to invoking $\Psi_{\mathcal{F}}(r)$. The reason we need the second term in the definition of \mathcal{E} is that we intend to apply Lemma 3.5 on each of the sets U_i in order to obtain certain subsets of U_i . This term guarantees that even the subsets of U_i will be “regular-enough” for our purposes (via Claim 5.1).

5 Proofs of Main Results

We start with the proof of Lemma 4.2, which is the main technical step in the proof of Theorem 1. We then use Theorem 1 in order to prove Theorem 2. We assume the reader is familiar with the

overview of the proof of Lemma 4.2 given in Section 4. For the proof we need the following simple fact, which states that large enough subsets of a regular pair are themselves somewhat regular.

Claim 5.1 *If (A, B) is a γ -regular pair with density η , and $A' \subseteq A$ and $B' \subseteq B$ satisfy $|A'| \geq \xi|A|$ and $|B'| \geq \xi|B|$ for some $\xi \geq \gamma$, then (A', B') is a $\max\{2\gamma, \gamma/\xi\}$ -regular pair.*

Proof: As (A, B) is a γ -regular pair with density η , then by definition of a regular pair, for every pair of subsets of $A' \subseteq A$ with $|A'| \geq \xi|A| \geq \gamma|A|$ and $B' \subseteq B$ with $|B'| \geq \xi|B| \geq \gamma|B|$ we have $|d(A', B') - d(A, B)| \leq \gamma$. Note, that if A' and B' are as above, then for every pair of subsets $A'' \subseteq A'$ and $B'' \subseteq B'$ satisfying $|A''| \geq \frac{\gamma}{\xi}|A'|$ and $|B''| \geq \frac{\gamma}{\xi}|B'|$ also satisfy $|A''| \geq \gamma|A|$ and $|B''| \geq \gamma|B|$. Therefore, by the γ -regularity of (A, B) we have $|d(A'', B'') - d(A, B)| \leq \gamma$. We thus conclude that $|d(A'', B'') - d(A', B')| \leq 2\gamma$. Hence, (A', B') is $\max\{2\gamma, \gamma/\xi\}$ -regular. \blacksquare

Proof of Lemma 4.2: Fix any family of graphs \mathcal{F} . Let $\Psi_{\mathcal{F}}(r)$ be the function from Definition 4.6 and define the following functions of r :

$$\alpha(r) = \alpha_{\mathcal{F}}(r) = \delta_{3.5}(\Psi_{\mathcal{F}}(r), \gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(r))), \quad (5)$$

$$\beta(r) = \beta_{\mathcal{F}}(r) = \alpha(r) \cdot \gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(r)), \quad (6)$$

and

$$\mathcal{E}(r) = \mathcal{E}_{\mathcal{F}}(r) = \begin{cases} \epsilon/6, & r = 0 \\ \min\{\beta(r), \epsilon/6\}, & r \geq 1 \end{cases} \quad (7)$$

For the rest of the proof set

$$S(\epsilon) = S_{\mathcal{F}}(\epsilon) = S_{3.8}(6/\epsilon, \mathcal{E}), \quad (8)$$

and note that as we define $S(\epsilon)$ in terms of $m = 6/\epsilon$ we get by Proposition 3.10 that $S(\epsilon)$ is indeed a function of ϵ only. We now set $N_{\mathcal{F}}(\epsilon)$ to be the following function of ϵ

$$N = N_{\mathcal{F}}(\epsilon) = S(\epsilon) \quad (9)$$

(as we have just argued, $S(\epsilon)$ and therefore also N can be upper bounded by functions of ϵ only). We postpone the definition of $f_{\mathcal{F}}(\epsilon)$ and $\delta_{\mathcal{F}}(\epsilon)$ till the end of the proof.

In the rest of the proof we consider any graph G on n vertices, with $n \geq N \geq S(\epsilon)$, which is ϵ -far from being induced \mathcal{F} -free. Given G , we can use Lemma 3.8 with $m = 6/\epsilon$ and $\mathcal{E}(r)$ as defined in (7), in order to obtain an equipartition \mathcal{A} of $V(G)$ into $6/\epsilon \leq k \leq S(\epsilon)$ clusters V_1, \dots, V_k (this is possible by item (1) in Lemma 3.8). Throughout the rest of the proof, k will denote the size of the equipartition returned by Lemma 3.8. By item (2) of Lemma 3.8, for every $1 \leq i \leq k$ we have sets $U_i \subseteq V_i$ each of size at least $n/S(\epsilon)$. Also, by item (3) of Lemma 3.8, **every** pair of these sets is at least $\beta(k)$ -regular (recall that $\mathcal{E}(k) \leq \beta(k)$). For each $1 \leq i \leq k$, apply Lemma 3.5 on the subgraph induced by G on each U_i with $\ell = \Psi_{\mathcal{F}}(k)$ and $\gamma = \gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k))$ in order to obtain the appropriate sets $W_{i,1}, \dots, W_{i,\Psi_{\mathcal{F}}(k)} \subset U_i$, all of size at least $\alpha(k)|U_i|$ (recall the definition of $\alpha(r)$ in (5)). It is crucial to note that we apply Lemma 3.5 on each of the sets U_1, \dots, U_k *after* we apply Lemma 3.8 on G , thus we “know” the value of k . The following observation will be useful for the rest of the proof:

Claim 5.2 *All the pairs $(W_{i,i'}, W_{j,j'})$ are $\gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k))$ -regular. Also, if $i \neq j$ then we also have $|d(W_{i,i'}, W_{j,j'}) - d(U_i, U_j)| \leq \epsilon/6$.*

Proof: Consider first pairs that belong to the same set U_i . In this case, the fact that any pair $(W_{i,i'}, W_{i,j'})$ is $\gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k))$ -regular follows immediately from our choice of these sets, as we applied Lemma 3.5 on each set U_i with $\gamma = \gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k))$. Consider now pairs that belong to different sets U_i, U_j . As was mentioned above, any pair (U_i, U_j) is $\beta(k)$ -regular. As each set $W_{i,j}$ satisfies $|W_{i,j}| \geq \alpha(k)|U_i|$, we get from Claim 5.1 and the definition of $\beta(k)$ that any pair $(W_{i,i'}, W_{j,j'})$ is at least $\max\{2\beta(k), \beta(k)/\alpha(k)\} \leq \gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k))$ -regular (here we use the fact that $\alpha(k) \leq 1/2$, which is guaranteed by Comment 3.6). Finally, as each of the sets $W_{i,j}$ satisfies $|W_{i,j}| \geq \alpha(k)|U_i| \geq \beta(k)|U_i| \geq \mathcal{E}(k)|U_i|$ we get from the fact that each pair (U_i, U_j) is $\mathcal{E}(k)$ -regular that $|d(W_{i,i'}, W_{j,j'}) - d(U_i, U_j)| \leq \mathcal{E}(k) \leq \epsilon/6$, thus completing the proof. ■

Recall that our goal is to show that G contains many induced copies of some graph $F \in \mathcal{F}$. To this end, we would like to apply Lemma 3.2 on some appropriately chosen subset of the sets $W_{i,j}$ defined above. As by Claim 5.2 all the pairs of sets $W_{i,j}$ are regular (we will later infer that they are regular enough for our purposes), we just have to find sets, whose densities will correspond to the edge set of some graph $F \in \mathcal{F}$ (recall the statement of Lemma 3.2). To this end, we define a graph \tilde{G} that will help us in choosing the sets $W_{i,j}$. The graph \tilde{G} is obtained from G by adding and removing the following edges, in the following order:

1. For $1 \leq i < j \leq k$ such that $|d(V_i, V_j) - d(U_i, U_j)| > \epsilon/6$, for all $v \in V_i$ and $v' \in V_j$ the pair (v, v') becomes an edge if $d(U_i, U_j) \geq \frac{1}{2}$, and becomes a non-edge if $d(U_i, U_j) < \frac{1}{2}$.
2. For $1 \leq i < j \leq k$ such that $d(U_i, U_j) < \frac{1}{3}\epsilon$, all edges between V_i and V_j are removed. For all $1 \leq i < j \leq k$ such that $d(U_i, U_j) > 1 - \frac{1}{3}\epsilon$, all non-edges between V_i and V_j become edges.
3. If for a fixed i all densities of pairs from $W_{i,1}, \dots, W_{i,l}$ are less than $\frac{1}{2}$, all edges within the vertices of V_i are removed. Otherwise, all the above densities are at least $\frac{1}{2}$ (by the choice of $W_{i,1}, \dots, W_{i,l}$ through Lemma 3.5), in which case all non-edges within V_i become edges.

In what follows we denote by $d(A, B)$ and $\tilde{d}(A, B)$ the edge density of the pair (A, B) in G and \tilde{G} , respectively. The following claim states several relations between the densities of G and \tilde{G} .

Claim 5.3 *For any i and $i' < j'$ we either have $\tilde{d}(W_{i,i'}, W_{i,j'}) = 1$ and $d(W_{i,i'}, W_{i,j'}) \geq \frac{1}{2}$ or $\tilde{d}(W_{i,i'}, W_{i,j'}) = 0$ and $d(W_{i,i'}, W_{i,j'}) \leq \frac{1}{2}$. Also, for any $i < j$ and any i', j' precisely one of the following holds:*

- (1) $\tilde{d}(V_i, V_j) = 1$ and $d(W_{i,i'}, W_{j,j'}) \geq \epsilon/6$.
- (2) $\tilde{d}(V_i, V_j) = 0$ and $d(W_{i,i'}, W_{j,j'}) \leq 1 - \epsilon/6$.
- (3) $\epsilon/6 \leq \tilde{d}(V_i, V_j) \leq 1 - \epsilon/6$ and $\epsilon/6 \leq d(W_{i,i'}, W_{j,j'}) \leq 1 - \epsilon/6$.

Proof: The proof follows easily from the three steps for obtaining \tilde{G} from G . The first assertion of the claim (concerning the relation between $\tilde{d}(W_{i,i'}, W_{i,j'})$ and $d(W_{i,i'}, W_{i,j'})$) follows directly from the third step of obtaining \tilde{G} . As for the second assertion (concerning the relation between $\tilde{d}(V_i, V_j)$ and $d(W_{i,i'}, W_{j,j'})$), assume the first step was applied to a pair (V_i, V_j) . In this case either $\tilde{d}(V_i, V_j) = 1$ and $d(U_i, U_j) \geq 1/2$ or $\tilde{d}(V_i, V_j) = 0$ and $d(U_i, U_j) \leq 1/2$. By Claim 5.2 we get that in the former case for any i', j' we have $d(W_{i,i'}, W_{j,j'}) \geq 1/2 - \epsilon/6 \geq \epsilon/6$, while in the latter $d(W_{i,i'}, W_{j,j'}) \leq 1/2 + \epsilon/6 \leq$

$1 - \epsilon/6$, as needed. Note, that if the first step was applied to a pair (V_i, V_j) then the second step has no effect, thus either (1) or (2) will hold at the end of the process. Assume the second step was applied to a pair (V_i, V_j) . In this case either $\tilde{d}(V_i, V_j) = 1$ and $d(U_i, U_j) \geq 1 - \epsilon/3$ or $\tilde{d}(V_i, V_j) = 0$ and $d(U_i, U_j) \leq \epsilon/3$. Again, by Claim 5.2, we get that in the former case $d(W_{i,i'}, W_{j,j'}) \geq 1 - \epsilon/3 - \epsilon/6 \geq \epsilon/6$ while in the latter $d(W_{i,i'}, W_{j,j'}) \leq \epsilon/3 + \epsilon/6 \leq 1 - \epsilon/6$. If none of the two steps was applied to (V_i, V_j) , then we initially had $|d(V_i, V_j) - d(U_i, U_j)| \leq \epsilon/6$ and $\epsilon/3 \leq d(U_i, U_j) \leq 1 - \epsilon/3$. Thus, item (3) holds as in this case we have $\epsilon/6 \leq d(V_i, V_j) = \tilde{d}(V_i, V_j) \leq 1 - \epsilon/6$ and by Claim 5.2 for any i', j' we have $\epsilon/6 \leq d(W_{i,i'}, W_{j,j'}) \leq 1 - \epsilon/6$. ■

Claim 5.4 *The graphs G and \tilde{G} differ by less than ϵn^2 edges.*

Proof: As the number of pairs $v \in V_i, v' \in V_j$ is n^2/k^2 , and by item (4) of Lemma 3.8 the number of pairs $1 \leq i < j \leq k$ for which $|d(V_i, V_j) - d(U_i, U_j)| > \epsilon/6 = \mathcal{E}(0)$ is at most $\mathcal{E}(0) \binom{k}{2} = \frac{1}{6} \epsilon \binom{k}{2}$, in the first step we changed less than $\frac{1}{6} \epsilon \binom{k}{2} \frac{n^2}{k^2} \leq \frac{1}{6} \epsilon n^2$ edges. In the second stage, if $d(U_i, U_j) < \frac{1}{3} \epsilon$ then by the modifications made in the first step, we have $d(V_i, V_j) < \frac{1}{2} \epsilon$. Similarly if $d(U_i, U_j) > 1 - \frac{1}{3} \epsilon$ then by the modifications made in the first step, we have $d(V_i, V_j) > 1 - \frac{1}{2} \epsilon$. Thus in this step we make at most $\binom{k}{2} \frac{1}{2} \epsilon (n^2/k^2) \leq \frac{1}{2} \epsilon n^2$ modifications. Finally, in the third step we make at most $k \binom{n/k}{2} \leq n^2/k$ modifications. As we apply Lemma 3.8 with $m = 6/\epsilon$, we have $n^2/k \leq \frac{1}{6} \epsilon n^2$. Altogether, we make less than ϵn^2 modifications. ■

We now turn to use the notion of colored-homomorphism, which was introduced in Section 4. For the rest of the proof, let R be the following colored complete graph on k vertices. We color $i \in V(R)$ white if V_i is edgeless in \tilde{G} . Otherwise (i.e. V_i is a complete graph in \tilde{G} , by step (3) in obtaining \tilde{G} from G) we color v_i black. If $\tilde{d}(V_i, V_j) = 0$ we color (i, j) white, if $\tilde{d}(V_i, V_j) = 1$ we color (i, j) black, otherwise (i.e. $\epsilon/6 \leq \tilde{d}(V_i, V_j) \leq 1 - \epsilon/6$, by Claim 5.3) we color (i, j) grey. Our goal in the following two claims is to identify a graph $F \in \mathcal{F}$, which we will later show to be abundant in G .

Claim 5.5 *\tilde{G} spans an induced copy of a graph $F' \in \mathcal{F}$. Moreover, $F' \mapsto_c R$.*

Proof: As G is by assumption ϵ -far from being induced \mathcal{F} -free, and by Claim 5.4 \tilde{G} is obtained from G by making less than ϵn^2 modifications (of adding and removing edges) \tilde{G} spans an induced copy of a graph $F' \in \mathcal{F}$. We claim that there is a colored-homomorphism from F' to R . Indeed, consider a mapping $\varphi : V(F') \mapsto V(R)$ which maps all the vertices of F' that belong to V_i to vertex i of R . We claim that this is a colored-homomorphism from F' to R . Suppose first that (u, v) is an edge of F' . If u and v belong to the same vertex set V_i , then V_i must be complete in \tilde{G} . By definition of φ they are both mapped to $i \in V(R)$ and by our coloring of R , vertex i is colored black. If $u \in V_i$ and $v \in V_j$ then it cannot be the case that $\tilde{d}(V_i, V_j) = 0$, hence $(i, j) \in E(R)$ was not colored white. Similarly, if (u, v) is not an edge of F' , then if u and v belong to the same vertex set V_i , then V_i must be edgeless. Hence, vertex i is colored white. If $u \in V_i$ and $v \in V_j$ then it cannot be the case that $\tilde{d}(V_i, V_j) = 1$, hence $(i, j) \in E(R)$ was not colored black. We thus get that φ satisfies the definition of a colored-homomorphism. ■

Claim 5.6 *There is a graph $F \in \mathcal{F}$ of size $f \leq \Psi_{\mathcal{F}}(k)$ for which $F \mapsto_c R$.*

Proof: By Claim 5.5, there is a graph $F' \in \mathcal{F}$ for which $F' \mapsto_c R$. Therefore, R belongs to \mathcal{F}_k (recall Definition 4.5 and the fact that R is of size k). It thus follows from the definition of $\Psi_{\mathcal{F}}$ (see Definition 4.6) that \mathcal{F} contains a graph of size at most $\Psi_{\mathcal{F}}(k)$ such that $F \mapsto_c R$. ■

The reader may want to recall at this stage that in order to apply Lemma 3.2 with respect to a graph on f vertices we need f distinct vertex sets. The following proposition will enable us to apply Lemma 3.2 on an appropriately chosen f sets of vertices in order to infer that G contains many induced copies of F .

Proposition 5.7 *Let F be the graph from Claim 5.6 and denote its vertex set by $\{1, \dots, f\}$ with $f \leq \Psi_{\mathcal{F}}(k)$. Let $\varphi : V(F) \mapsto V(R)$ be the colored homomorphism from F to R , which is guaranteed to exist by Claim 5.6, and put $t_i = \varphi(i)$ for every $i \in V(F)$. The following holds with respect to the sets $W_{t_1,1}, \dots, W_{t_f,f}$:*

- If $(i, j) \in E(F)$ then $d(W_{t_i,i}, W_{t_j,j}) \geq \epsilon/6$.
- If $(i, j) \notin E(F)$ then $d(W_{t_i,i}, W_{t_j,j}) \leq 1 - \epsilon/6$.

Proof: First, note that we choose the sets as $W_{t_1,1}, \dots, W_{t_f,f}$ in order to make sure that we do not choose the same $W_{i,i'}$ twice, because as we may need to use several sets $W_{i,j}$ from the same set U_i (in the case that $t_i = t_j$ for some $i \neq j$). Also, observe that as $f \leq \Psi_{\mathcal{F}}(k)$ and we obtained through Lemma 3.5 $\ell = \Psi_{\mathcal{F}}(k)$ sets $W_{i,j}$ from each U_i , we can indeed choose the sets in the above manner, even if all the chosen sets $W_{i,j}$ belong to the same U_i .

Assume that $(i, j) \in E(F)$. As φ is a colored homomorphism from F to R we conclude that either $\varphi(i) = \varphi(j) = t$ and $t \in V(R)$ is colored black or $\varphi(i) = t \neq t' = \varphi(j)$ and $(t, t') \in E(R)$ is colored black or grey. By the way we colored R in the paragraph preceding Claim 5.5 this means that either $\varphi(i) = \varphi(j) = t$ and V_t is a complete graph in \tilde{G} or $\varphi(i) = t \neq t' = \varphi(j)$ and $\tilde{d}(V_t, V_{t'}) \geq \epsilon/6$. Finally, by Claim 5.3 this means that in both cases $d(W_{t_i,i}, W_{t_j,j}) \geq \epsilon/6$. The case of $(i, j) \notin E(F)$ is analogous. ■

The proof of Lemma 4.2 now follows easily from the above proposition. Consider the sets $W_{t_1,1}, \dots, W_{t_f,f}$ as in Proposition 5.7. By Claim 5.2 any pair of these sets is at least $\gamma_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k))$ -regular in G . Moreover, by Proposition 5.7, these $f \leq \Psi_{\mathcal{F}}(k)$ sets satisfy in G (**not** in \tilde{G}) the edge requirements of Lemma 3.2, which are needed in order to infer that they span many induced copies of F (recall that F has at most $\Psi_{\mathcal{F}}(k)$ vertices). Thus, Lemma 3.2 ensures that $W_{t_1,1}, \dots, W_{t_f,f}$ span in G (**not** in \tilde{G}) at least

$$\delta_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k)) \cdot \prod_{i=1}^f |W_{t_i,i}| \tag{10}$$

induced copies of F . We next show that we can take F as the graph in the statement of Lemma 4.2. To show this, we should only define the functions $f_{\mathcal{F}}(\epsilon)$ and $\delta_{\mathcal{F}}(\epsilon)$ (the function $N_{\mathcal{F}}(\epsilon)$ is defined in (9)). As $|U_i| \geq n/S(\epsilon)$ and $|W_{t_i,i}| \geq \alpha(k)|U_i|$, we conclude from (10) that G contains at least

$$\delta_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(k)) \cdot (\alpha(k)/S(\epsilon))^f \cdot n^f \tag{11}$$

induced copies of F . Thus, as $f \leq \Psi_{\mathcal{F}}(k)$, $k \leq S(\epsilon)$ and by the monotonicity properties of all the functions considered in the proof, we can replace k with $S(\epsilon)$ and f with $\Psi_{\mathcal{F}}(S(\epsilon))$ and thus define

$$f_{\mathcal{F}}(\epsilon) = \Psi_{\mathcal{F}}(S(\epsilon)). \quad (12)$$

Similarly, we can replace k and f in (11) in order to define

$$\delta_{\mathcal{F}}(\epsilon) = \frac{\delta_{3.2}(\epsilon/6, \Psi_{\mathcal{F}}(S(\epsilon)))}{(S(\epsilon)/\alpha(S(\epsilon)))^{\Psi_{\mathcal{F}}(S(\epsilon))}}. \quad (13)$$

This completes the proof of Lemma 4.2. ■

Before proving Theorem 1 we briefly discuss the notions of uniform and non-uniform testing, which were defined and studied in [11] and [10]. We give here only a rough overview of the result of [10]. A tester is defined in [10] as *non-uniform* if it knows ϵ in advance, and therefore should be able to distinguish between graphs that satisfy \mathcal{P} from those that are ϵ -far from satisfying it (only for that specific ϵ). A tester is *uniform* if it can accept ϵ as part of the input. The main result of [10] is that there are monotone graph properties, which have non-uniform one-sided testers but cannot be tested by a uniform (one-sided or two-sided) testers. The main reason is that they are monotone and decidable properties (in fact even in *coNP*) with the property that the query complexity of testing them cannot be computed given only the value of ϵ . It thus follows that we cannot design uniform testers for all the hereditary graph properties.

Note, that in (9), (12) and (13) the only function that may be non-computable is $\Psi_{\mathcal{F}}$. Thus whenever this function is computable so are the three functions of Lemma 4.2. As the proof of Theorem 1 suggests (see below), once these functions are computable, the tester is uniform. Finally, we note that for any reasonable graph property, and in particular those that were discussed in Subsection 2.1, $\Psi_{\mathcal{F}}$ is indeed computable (not necessarily very efficiently). Thus, these properties are testable in the usual sense. We thus assume henceforth that \mathcal{F} is such that the functions $N_{\mathcal{F}}(\epsilon)$, $f_{\mathcal{F}}(\epsilon)$ and $\delta_{\mathcal{F}}(\epsilon)$ are computable. Note however, that even if they are not computable, we still get a non-uniform tester for any (decidable) hereditary graph property.

Proof of Theorem 1: We show that any hereditary property can be tested with one-sided error even by an oblivious tester. Fix any hereditary graph property \mathcal{P} , and let \mathcal{F} be the family of forbidden induced subgraphs of \mathcal{P} as in Definition 4.1. Let $N_{\mathcal{F}}(\epsilon)$, $f_{\mathcal{F}}(\epsilon)$ and $\delta_{\mathcal{F}}(\epsilon)$ be the functions of Lemma 4.2 and assume they are computable. To design our one-sided error tester for \mathcal{P} we just need to note that if a graph on n vertices contains at least δn^f induced copies of a graph F on f vertices, then sampling $2/\delta$ sets of f vertices each, which is a total of $2f/\delta$, finds an induced copy of F with probability at least $2/3$.

Given a graph G the one-sided error tester for \mathcal{P} works as follows; it asks the oracle for a subgraph of G induced by a randomly chosen set of $\max\{N_{\mathcal{F}}(\epsilon), 2f_{\mathcal{F}}(\epsilon)/\delta_{\mathcal{F}}(\epsilon)\}$ vertices. It declares G to be a graph satisfying \mathcal{P} if and only if the induced subgraph on S satisfies \mathcal{P} . Clearly, if G satisfies \mathcal{P} , then as \mathcal{P} is hereditary the algorithm accepts G with probability 1. If G is ϵ -far from satisfying \mathcal{P} and G has less than $N_{\mathcal{F}}(\epsilon)$ vertices, the algorithm answers correctly with probability 1, as in this case S spans G . If G has more than $N_{\mathcal{F}}(\epsilon)$ vertices, then by Lemma 4.2 there is a member of \mathcal{F} of size $f \leq f_{\mathcal{F}}(\epsilon)$ such that G spans $\delta_{\mathcal{F}}(\epsilon)n^f$ induced copies of F . By the observation from the preceding paragraph, S spans an induced copy of F with probability at least $2/3$. As $F \in \mathcal{F}$ and

\mathcal{P} is hereditary, we get that with probability at least $2/3$, the graph spanned by S does not satisfy \mathcal{P} . Hence, the tester rejects G with probability at least $2/3$. Also, its query complexity is always a function of ϵ only. \blacksquare

Proof of Theorem 2: Let \mathcal{P} be a semi-hereditary property and let \mathcal{H} be the hereditary graph property as in Definition 2.2. We next show that \mathcal{P} has an oblivious one-sided error tester. As \mathcal{H} is hereditary we get from Theorem 1 and the fact that its proof actually gives an oblivious tester for \mathcal{H} that there is a function $Q_{\mathcal{H}}(\epsilon)$ such that \mathcal{H} can be tested by an oblivious one-sided error tester with query complexity $Q_{\mathcal{H}}(\epsilon)$. The oblivious tester T we design for testing \mathcal{P} works as follows: its query complexity is $Q(\epsilon) = \max\{M(\epsilon/2), Q_{\mathcal{H}}(\epsilon/2)\}$ (the function M is part of Definition 2.2). After getting from the oracle the randomly chosen induced subgraph, which we denote by G' , the tester T proceeds as follows: If G' is of size strictly smaller than $Q(\epsilon)$, the algorithm accepts if and only if G' satisfies \mathcal{P} . If G' is of size at least $Q(\epsilon)$ the algorithm accepts if and only if G' satisfies \mathcal{H} .

We turn to show that T is indeed an oblivious one-sided error tester for \mathcal{P} . We first observe that T satisfies the definition of an oblivious tester. We also note that if the input graph is of size less than $Q(\epsilon)$ then we accept the input if and only if it satisfies \mathcal{P} because by the definition of an oblivious tester this means that the input graph was of size less than $Q(\epsilon)$ and therefore the oracle returned the entire input graph. Let us now consider an input of size at least $Q(\epsilon)$ and recall that $Q(\epsilon) \geq M(\epsilon/2)$. If this input satisfies \mathcal{P} then by the first item of Definition 2.2 it also satisfies \mathcal{H} , and as in this case we accept if and only if G' satisfies \mathcal{H} this means that T accepts the input. Hence, T has one-sided error. Suppose now that the input is ϵ -far from satisfying \mathcal{P} . This means that after adding/deleting $\frac{1}{2}\epsilon n^2$ edges, the input is still $\frac{\epsilon}{2}$ -far from satisfying \mathcal{P} . By item 2 of Definition 2.2 and as in this case the input must be of size at least $M(\epsilon/2)$, this means that after adding/deleting $\frac{1}{2}\epsilon n^2$ edges, the input still contains an induced subgraph not satisfying \mathcal{H} . In other words, this means that the input is at least $\frac{\epsilon}{2}$ -far from satisfying \mathcal{H} . As $Q(\epsilon) \geq Q_{\mathcal{H}}(\epsilon/2)$ we infer that with probability at least $2/3$ the graph G' spans an induced subgraph not satisfying \mathcal{H} and therefore G' does not satisfy \mathcal{H} (as it is hereditary). As in this case T accepts if and only if G' satisfies \mathcal{H} , this means that T will reject an input that is ϵ -far from satisfying \mathcal{P} with probability at least $2/3$.

Assume now that property \mathcal{P} has a one-sided error oblivious tester T . Our goal is to show the existence of a hereditary property \mathcal{H} as in Definition 2.2. Let \mathcal{F} be the following family of graphs: a graph F on $|V(F)|$ vertices belongs to \mathcal{F} if (i) For some $\epsilon > 0$ the query complexity of T satisfies $Q(\epsilon) = |V(F)|$ (recall that the query complexity of T is a function of ϵ only). (ii) If for this ϵ the sample of vertices spans a graph isomorphic to F , then T rejects the input with positive probability. We claim that we can take \mathcal{H} in Definition 2.2 to be the property of being induced \mathcal{F} -free.

To establish the first item of Definition 2.2 it is enough to show that there is no graph G satisfying \mathcal{P} , which spans an induced subgraph isomorphic to a graph $F \in \mathcal{F}$. Suppose such a G exists, and consider the execution of T on G with an ϵ for which $Q(\epsilon) = |V(F)|$. By definition of \mathcal{F} we get that T asks for a random subgraph of G of size $|V(F)|$, and that if T gets a graph isomorphic to F it rejects G with positive probability. As we assume that G spans an induced copy of a graph isomorphic to F , this means that T has a non-zero probability of rejecting G , contradicting our assumption that T is one-sided.

To establish the second item of Definition 2.2, we claim that we can take $M(\epsilon) = Q(\epsilon)$. Indeed, consider a graph G on at least $Q(\epsilon)$ vertices that is ϵ -far from satisfying \mathcal{P} . As T is a tester for \mathcal{P} it should reject G with non-zero probability. By definition of an oblivious tester and as G has at least $Q(\epsilon)$ vertices, this means that G must contain an induced subgraph F , of size precisely $Q(\epsilon)$, with

the property that if T gets F from the oracle then it rejects G . By definition of \mathcal{F} this means that $F \in \mathcal{F}$. Hence, we can take F itself to be the graph not satisfying \mathcal{H} . ■

6 Proofs of Additional Results

In this Section we give the proofs of Theorems 3, 4 and 5. We start with the proof of Theorem 3. Note that when considering the notion of ϵ -far_{del} there is no sense in considering hereditary properties, which are not satisfied by some independent set, as in this case any graph with even a single independent set (say, of size 3) is arbitrarily far from satisfying the property. Moreover, finding this independent set requires $\Omega(n^2)$ queries.

Our main tool for the proof of Theorem 3 is the following result, which is essentially proved by Frankl and Füredi in [21].

Theorem 7 ([21]) *For any graph $F = (R, T)$, with $|T| = t > 0$ edges there is a constant $\delta = \delta(F)$ with the following property: For any integer n there is a graph $G_n = (V, E)$ on n vertices, which consists of $(1 - n^{-\delta})\binom{n}{2}/t$ induced copies of F , such that no two copies of F share an edge.*

Proof of Theorem 3: By the discussion above we may assume that \mathcal{P} has at least one forbidden induced subgraph $F = (R, T)$ and that F is not an independent set. Put $t = |T|$ and for any n let G_n be the graph, whose existence is guaranteed by Theorem 7. As all these graphs consist of $(1 - n^{-\delta})\binom{n}{2}/t > n^2/4t$ induced copies of F , where non of the copies share an edge, these graphs are all at least $\frac{1}{4t}$ -far_{del} from being induced F free. Hence, they are also at least $\frac{1}{4t}$ -far_{del} from satisfying \mathcal{P} . On the other, as we assume that any clique satisfies \mathcal{P} , and G contains $(1 - n^{-\delta})\binom{n}{2}$ edges, any randomized algorithm with query-complexity much smaller than n^δ cannot test_{del} property \mathcal{P} . The reason is that the algorithm has a negligible probability of distinguishing between the graphs G_n , which are $\frac{1}{4t}$ -far_{del} from satisfying \mathcal{P} , and a clique of size n , which by assumption satisfies \mathcal{P} . ■

Suppose we define ϵ -far_{add} and testable_{add} but now allowing only edge additions. One can easily see that simple modifications of the proof of Theorem 3 imply that the same lower bound can be proved for testing_{add} any hereditary property, which is not closed under edge additions and which is satisfied by any edgeless graph.

We continue with the proof of Theorem 4. As most of the technical details are very similar to those appearing in [4] we only discuss the main idea needed to obtain the extension of the result of [4]. We start with a useful result of [4].

Definition 6.1 (Indistinguishability) *Two graph properties \mathcal{P} and \mathcal{Q} are called indistinguishable if for every $\epsilon > 0$ there exists $N = N(\epsilon)$ satisfying the following; A graph on $n \geq N$ vertices satisfying one of the properties is never ϵ -far from satisfying the other.*

Lemma 6.2 ([4]) *If \mathcal{P} and \mathcal{Q} are indistinguishable graph properties, then \mathcal{P} is testable if and only if \mathcal{Q} is testable.*

We next define an extension of the notion of colorability. A similar notion was used in [4], where \mathcal{F} was restricted to be *finite*.

Definition 6.3 (\mathcal{F} -colorability) Suppose we are given an integer c , and a (possibly infinite) family (with repetitions) \mathcal{F} of graphs, each of which is provided with a c -coloring (i.e. a function from its vertex set to $\{1, \dots, c\}$ which is not necessarily a proper c -coloring in the usual sense). A c -coloring of a graph G is called an \mathcal{F} -coloring if no member of \mathcal{F} appears as an induced subgraph of G with an identical coloring. A graph G is called \mathcal{F} -colorable if there exists an \mathcal{F} -coloring of it.

Note, that for any family of colored graphs \mathcal{F} (finite or infinite), being \mathcal{F} -colorable is a hereditary graph property. We thus get the following from Theorem 1:

Lemma 6.4 For any family of colored graphs \mathcal{F} , being \mathcal{F} -colorable is testable.

Note, that by Theorem 1 being \mathcal{F} -colorable is in fact testable with one-sided error, but we do not need this stronger assertion here. The following lemma shows the relevance of the notion of \mathcal{F} -colorability for the proof of Theorem 4.

Lemma 6.5 For every first order property \mathcal{P} of the form

$$\exists x_1, \dots, x_t \bigwedge_{i=1}^{\infty} \forall y_1, \dots, y_i A_i(x_1, \dots, x_t, y_1, \dots, y_i)$$

there exists a (possibly infinite) family \mathcal{F} , of $(2^{t+\binom{t}{2}} + 1)$ -colored graphs such that the property \mathcal{P} is indistinguishable from the property of being \mathcal{F} -colorable.

Proof: (sketch) The proof uses ideas very similar to those used to prove Lemma 2.2 in [4] and is thus omitted. We briefly mention that one can use the same technique of [4] along with the fact that one is allowed to put in \mathcal{F} *infinitely* many forbidden colored subgraphs. ■

Proof of Theorem 4: Immediate from Lemmas 6.2, 6.4 and 6.5. ■

We conclude this section with the proof of Theorem 5.

Proof of Theorem 5: For each of the hereditary properties \mathcal{P}_i , let \mathcal{F}_i be the family of forbidden induced subgraphs of \mathcal{P}_i as in Definition 4.1, and let $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3 \cup \dots$. Clearly, a graph G satisfies all the properties of \mathcal{P} if and only if it is induced \mathcal{F} -free. Consider a graph G , which is ϵ -far from satisfying all the properties of \mathcal{P} . In this case G is also ϵ -far from being induced \mathcal{F} -free, hence, by Lemma 4.2, there is a graph $F \in \mathcal{F}$ of size $f = f_{\mathcal{F}}(\epsilon)$ such that G contains $\delta_{\mathcal{F}}(\epsilon)n^f$ induced copies of F . Note, that adding or removing an edge from G destroys at most $\binom{n}{f-2} \leq n^{f-2}$ induced copies of F . Thus, one must add or delete at least $\delta_{\mathcal{F}}(\epsilon)n^2$ edges to G in order to turn it into a graph containing no induced copy of F . Let i be such that $F \in \mathcal{F}_i$. We may now infer that G is $\delta_{\mathcal{F}}(\epsilon)$ -far from satisfying \mathcal{P}_i . Finally, note that as \mathcal{F} is determined by \mathcal{P} , we can also say that G is $\delta_{\mathcal{P}}(\epsilon)$ -far from satisfying \mathcal{P}_i . ■

7 Concluding Remarks and Open Problems

- Our main result in this paper can be considered a characterization of the natural graph properties that are testable with one-sided error. Thus, a natural and interesting open problem related to this paper is to complete the characterization of the graph properties that are testable with one-sided error by arbitrary testers, and not just oblivious ones.
- Theorem 1 asserts that any hereditary property is testable with one-sided error. However, the upper bounds on the query complexity, which this theorem guarantees are huge. Even for rather simple properties, these bounds are towers of towers of exponents of height polynomial in $1/\epsilon$. Some specific properties, such as k -colorability, have far more efficient testers, whose query complexity is polynomial in $1/\epsilon$ (see [6]). For others, like being H -free (that is, containing no copy of H as a (not necessarily induced) subgraph), it is known that whenever H is not bipartite, there is no tester (one-sided or two-sided) whose query complexity is polynomial in $1/\epsilon$ (see [1], [9]). Recall that a hereditary property \mathcal{P} is equivalent to being $\mathcal{F}_{\mathcal{P}}$ -free for a possibly infinite family of graphs $\mathcal{F}_{\mathcal{P}}$. The hardness of testing hereditary properties for which $\mathcal{F}_{\mathcal{P}}$ is finite is (relatively) well understood, as it follows from the main result of [7] that if $\mathcal{F}_{\mathcal{P}}$ has a graph on at least 5 vertices, then there is no tester (one-sided or two-sided) for \mathcal{P} , whose query complexity is polynomial in $1/\epsilon$. When $\mathcal{F}_{\mathcal{P}}$ is infinite the situation is much more complicated, and there are no general results which guarantee or rule out the possibility of designing testers with query complexity polynomial in $1/\epsilon$. In particular, a natural intriguing and probably challenging problem is the following:

Which hereditary graph properties can be tested with $\text{poly}(1/\epsilon)$ queries?

As a special case of this problem, it seems interesting to study the query complexity needed to test the natural graph properties that were discussed in Subsection 2.1.

- Theorem 2 gives a precise characterization of the graph properties that have oblivious one-sided testers. As we have explained in Section 1, any natural property that can be tested, can be tested by an oblivious tester. It may thus be simpler, but still very interesting, to resolve the following problem:

Which graph properties have (possibly two-sided) oblivious testers?

Note, that the definition of an oblivious tester implicitly assumes that the query complexity of such a tester is a function of ϵ only.

- Fischer and Newman [19] have recently shown that every testable graph property is also estimable, namely, for any such property one can estimate how far is a given graph from satisfying the property (in this paper this quantity is denoted by ϵ) while making a constant number of queries. Combining Theorem 1 and the result of [19] we get that any hereditary property is estimable.

References

- [1] N. Alon, Testing subgraphs in large graphs, *Random Structures and Algorithms* 21 (2002), 359-370. Also, *Proc. 42nd IEEE FOCS, IEEE* (2001), 434-441.

- [2] N. Alon, E. Fischer, I. Newman and A. Shapira, A combinatorial characterization of the testable graph properties: it's all about regularity, Proc. of STOC'06, 251-260.
- [3] N. Alon, R. A. Duke, H. Lefmann, V. Rödl and R. Yuster, The algorithmic aspects of the Regularity Lemma, Proc. 33rd IEEE FOCS, Pittsburgh, IEEE (1992), 473-481. Also: J. of Algorithms 16 (1994), 80-109.
- [4] N. Alon, E. Fischer, M. Krivelevich and M. Szegedy, Efficient testing of large graphs, Proc. of 40th FOCS, New York, NY, IEEE (1999), 656–666. Also: Combinatorica 20 (2000), 451-476.
- [5] N. Alon, M. Krivelevich, I. Newman and M. Szegedy, Regular languages are testable with a constant number of queries, Proc. 40th FOCS, New York, NY, IEEE (1999), 645–655. Also: SIAM J. on Computing 30 (2001), 1842-1862.
- [6] N. Alon and M. Krivelevich, Testing k-colorability, SIAM J. Discrete Math., 15 (2002), 211-227.
- [7] N. Alon and A. Shapira, A characterization of easily testable induced subgraphs, Proc. of the 15th Annual ACM-SIAM SODA, ACM Press (2004), 935-944. Also, Combinatorics, Probability and Computing, to appear.
- [8] N. Alon and A. Shapira, Testing satisfiability, Proc. 13th Annual ACM-SIAM SODA, ACM Press (2002), 645-654. Also: Journal of Algorithms, 47 (2003), 87-103.
- [9] N. Alon and A. Shapira, Testing subgraphs in directed graphs, JCSS 69/3 (2004), 354-382. Also, Proc. of the 35th STOC, 2003, 700–709.
- [10] N. Alon and A. Shapira, A separation theorem in property-testing, manuscript, 2005.
- [11] N. Alon and A. Shapira, Every monotone graph-property is testable, Proc. of STOC 2005, 128-137. Also, SIAM J. on Computing, to appear.
- [12] N. Alon and A. Shapira, Homomorphisms in graph property testing - A survey, to appear.
- [13] N. Alon and U. Stav, What is the furthest graph from a hereditary property?, manuscript, 2006.
- [14] M. Blum, M. Luby and R. Rubinfeld, Self-testing/correcting with applications to numerical problems, JCSS 47 (1993), 549-595.
- [15] B. Bollobás, **Extremal Graph Theory**, Academic Press, New York (1978).
- [16] T. M. Chan, Polynomial-time approximation schemes for packing and piercing fat objects, Journal of Algorithms 46 (2003), 178-189.
- [17] E. Fischer, Testing graphs for colorability properties, Proc. of the 12th SODA (2001), 873-882.
- [18] E. Fischer, The art of uninformed decisions: A primer to property testing, The Computational Complexity Column of The Bulletin of the European Association for Theoretical Computer Science 75 (2001), 97-126.
- [19] E. Fischer and I. Newman, Testing versus estimation of graph properties, Proc. of STOC 2005, 138-146.

- [20] E. Fischer, I. Newman and J. Sgall, Functions that have read-twice constant width branching programs are not necessarily testable, *Random Structures and Algorithms*, in press.
- [21] P. Frankl and Z. Füredi, Colored packings of sets in combinatorial design theory, *Annals of Discrete Math.* 34 (1987), 165-178.
- [22] O. Goldreich, Combinatorial property testing - a survey, In: *Randomization Methods in Algorithm Design* (P. Pardalos, S. Rajasekaran and J. Rolim eds.), AMS-DIMACS (1998), 45-60.
- [23] O. Goldreich, S. Goldwasser and D. Ron, Property testing and its connection to learning and approximation, *Proc. of 37th Annual IEEE FOCS*, (1996), 339-348. Also, *JACM* 45(4): 653-750 (1998).
- [24] O. Goldreich and D. Ron, Property Testing in Bounded-Degree Graphs, *Proc. of STOC 1997*, 406-415.
- [25] O. Goldreich and L. Trevisan, Three theorems regarding testing graph properties, *Proc. 42nd IEEE FOCS*, IEEE (2001), 460-469. Also, *Random Structures and Algorithms*, 23(1):23-57, 2003.
- [26] M.C. Golumbic, **Algorithmic Graph Theory and Perfect Graphs**, Academic Press, 1980.
- [27] J. Komlós and M. Simonovits, Szemerédi's Regularity Lemma and its applications in graph theory. In: *Combinatorics, Paul Erdős is Eighty*, Vol II (D. Miklós, V. T. Sós, T. Szönyi eds.), János Bolyai Math. Soc., Budapest (1996), 295-352.
- [28] L. Lovász and B. Szegedy, Graph limits and testing hereditary graph properties, manuscript, 2006.
- [29] T. A. McKee and F.R. McMorris, **Topics in Intersection Graph Theory**, SIAM, Philadelphia, PA, 1999.
- [30] I. Newman, Testing of functions that have small width branching programs, *Proc. of 41th FOCS* (2000), 251-258.
- [31] M. Parnas and D. Ron, Testing the diameter of graphs, *Random structures and algorithms*, 20 (2002), 165-183.
- [32] J. L. Ramírez-Alfonsín (Editor), B. A. Reed (Editor), **Perfect Graphs**, Wiley, 2001.
- [33] V. Rödl and R. Duke, On graphs with small subgraphs of large chromatic number, *Graphs and Combinatorics* 1 (1985), 91-96.
- [34] V. Rödl and M. Schacht, Generalizations of the removal lemma, manuscript, 2006.
- [35] D. Ron, Property testing, in: P. M. Pardalos, S. Rajasekaran, J. Reif and J. D. P. Rolim, editors, *Handbook of Randomized Computing*, Vol. II, Kluwer Academic Publishers, 2001, 597-649.
- [36] R. Rubinfeld and M. Sudan, Robust characterization of polynomials with applications to program testing, *SIAM J. on Computing* 25 (1996), 252-271.

- [37] E. Szemerédi, Regular partitions of graphs, In: *Proc. Colloque Inter. CNRS* (J. C. Bermond, J. C. Fournier, M. Las Vergnas and D. Sotteau, eds.), 1978, 399–401.