

# Ordinal Embeddings of Minimum Relaxation: General Properties, Trees, and Ultrametrics\*

Noga Alon<sup>†</sup>    Mihai Bădoiu<sup>‡</sup>    Erik D. Demaine<sup>†§</sup>    Martin Farach-Colton<sup>¶</sup>  
MohammadTaghi Hajiaghayi<sup>†‡</sup>    Anastasios Sidiropoulos<sup>†||</sup>

## Abstract

We introduce a new notion of embedding, called *minimum-relaxation ordinal embedding*, parallel to the standard notion of minimum-distortion (metric) embedding. In an ordinal embedding, it is the relative order between pairs of distances, and not the distances themselves, that must be preserved as much as possible. The (multiplicative) relaxation of an ordinal embedding is the maximum ratio between two distances whose relative order is inverted by the embedding. We develop several worst-case bounds and approximation algorithms on ordinal embedding. In particular, we establish that ordinal embedding has many qualitative differences from metric embedding, and capture the ordinal behavior of ultrametrics and shortest-path metrics of unweighted trees.

## 1 Introduction

The classical field of *multidimensional scaling (MDS)* has witnessed a surge of interest in recent years with a slew of papers on *metric embeddings*; see e.g. [IM04]. The problem of multidimensional scaling is that of mapping points with some measured pairwise distances into some target metric space. Originally, the MDS community considered embeddings into an  $\ell_p$  space, with the goal of aiding in visualization, compression, clustering, or nearest-neighbor searching; thus, low-dimensional embeddings were sought. An *isometric embedding* preserves all distances, while more generally, *metric embeddings* tradeoff the dimension with the fidelity of the embeddings.

Note, however, that the distances themselves are not essential in nearest-neighbor searching and many contexts of visualization, compression, and clustering. Rather, the order of the distances captures sufficient information, that is, we might only need an embedding into a metric space with any monotone mapping of the distances. Such embeddings were heavily studied in the early MDS literature [CS74, Kru64a, Kru64b, She62, Tor52] and have been referred to as *ordinal embeddings*, *nonmetric MDS*, or *monotone maps*. Here, we use the first term.

---

\*A preliminary version of this paper is to appear in the Proceedings of the 16th Annual ACM-SIAM Symposium on Discrete Algorithms, January 2005.

<sup>†</sup>Schools of Mathematics and Computer Science, Tel Aviv University, Tel Aviv, Israel, email: nogaa@tau.ac.il. Research supported in part by a grant from the Israel Science Foundation, and by the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University.

<sup>‡</sup>MIT Computer Science and Artificial Intelligence Laboratory, 32 Vassar St., Cambridge, MA 02139, USA, email: {mihai, edemaine, hajiagha, tasos}@mit.edu

<sup>§</sup>Supported in part by NSF under grant number ITR ANI-0205445.

<sup>¶</sup>Department of Computer Science, Rutgers University, email: farach@cs.rutgers.edu

<sup>||</sup>Supported by the Paris Kanellakis Fellowship Fund and by the Alexandros S. Onassis Public Benefit Foundation.

While the early work on ordinal embeddings was largely heuristic, there has been some work with mathematical guarantees since then. Define a *distance matrix* to be any matrix of pairwise distances, not necessarily describing a metric. In [SFC04], it was shown that it is NP-hard to decide whether a distance matrix can be ordinally embedded into an additive metric, i.e., the shortest-path metric in a tree. Define the *ordinal dimension* of a distance matrix to be the smallest dimension of a Euclidean space into which the matrix can be ordinally embedded. Bilu and Linial [BL04] have shown that every matrix has ordinal dimension at most  $n - 1$ . They also applied the methods of [AFR85] to show that (in a certain well-defined sense) almost every  $n$ -point metric space has ordinal dimension  $\Omega(n)$ . Because ultrametrics can be characterized by the order of distances on all triangles, they are closed under monotone mappings. Holman [Hol72] showed that every  $n$ -point ultrametric can be isometrically embedded into  $(n - 1)$ -dimensional Euclidean space and that  $n - 1$  dimensions are necessary. Combined with the closure property just noted, this shows that the ordinal dimension of every ultrametric is exactly the maximal  $n - 1$ .<sup>1</sup>

Relaxations of ordinal embeddings have involved problems of deciding the realization of partial orders. For example, Opatrny [Opa79] showed that it is NP-hard to decide whether there is an embedding into one dimension satisfying a partial order that specifies the maximum edge for some triangles. Such partial orders on triangles are called *betweenness constraints*. Chor and Sudan [CS98] gave a  $1/2$ -approximation for maximizing the number of satisfied constraints. It is also NP-hard to decide whether there is an embedding into an additive metric that satisfies a partial order defined by the total order of each triangle [SFC04].

## 1.1 Our Results

We take a different approach. We define a metric  $M'$  to be an *ordinal embedding with relaxation*  $\alpha \geq 1$  of a distance matrix  $M$  if  $\alpha M[i, j] < M[k, l]$  implies  $M'[i, j] < M'[k, l]$ . In other words, significantly different distances have their relative order preserved. Note that in an ordinary ordinal embedding, we must respect distance equality, while in an ordinal embedding with relaxation 1, we may break ties. It is now natural to minimize the relaxation needed to embed a distance matrix  $M$  into a target family of metric spaces. Here we optimize the confidence with which we make an ordinal assertion, rather than the number of ordinal constraints satisfied.

In this paper, we prove a variety of results about the *Ordinal Relaxation Problem*. We show that the best relaxation achievable is always at most the best *distortion* of a metric embedding. Furthermore, while the optimal relaxation is bounded by the ratio between the largest and smallest distances in  $M$ , the optimal distortion can grow arbitrarily. Indeed, the ratio between the optimal relaxation and distortion can be arbitrarily large even when embedding into the line, and can be infinite when embedding into cut metrics. (We also give a polynomial-time algorithm to compute the best ordinal embedding into a cut metric.) We show that, if the target class of the embedding is ultrametrics, the relaxation and distortion are equal, and the optimal embedding can be computed in polynomial time. More surprisingly, we show that ultrametrics are the only target metrics for which all distance matrices have a bounded ratio between the best distortion and the best relaxation.

We demonstrate many more differences between ordinal and metric embeddings. While any metric can be isometrically embedded into  $\ell_\infty$ , there are four-point metrics that cannot be so embedded into any  $\ell_p$ ,  $p < \infty$ . In contrast, we show that it is possible to ordinally embed any distance matrix into  $\ell_p$  for any fixed  $1 \leq p \leq \infty$ . We show that the shortest-path metric of an

---

<sup>1</sup>This observation settles an open problem posed in [BL04] asking for the worst-case ordinal dimension of any metric on  $n$  points, which they showed was between  $n/2$  and  $n - 1$ . Ultrametrics show that the answer is  $n - 1$ .

unweighted tree can be ordinally embedded into  $d$ -dimensional Euclidean space with relaxation  $\tilde{O}(n^{1/d})$ . We also show that relaxation  $\Omega(n^{1/(d+1)})$  is sometimes necessary. In contrast, the best bounds on the worst-case distortion required are  $O(n^{1/(d-1)})$  and  $\Omega(n^{1/d})$  [Gup00]. The proof techniques required for the ordinal case are also substantially different (in particular because the usual “packing” arguments fail) and lead to approximation algorithms described below. We show that ultrametrics can be ordinally embedded into  $O(\lg n)$ -dimensional  $\ell_p$  space with relaxation 1. In contrast, the best known metric embedding of ultrametrics into  $c \lg n$ -dimensional space has distortion  $1 + \Omega(1/\sqrt{c})$  [BM04], and ordinary (no-relaxation) ordinal embeddings require  $n - 1$  dimensions. For general metrics, we show a lower bound of  $\Omega(\lg n / (\lg d + \lg \lg n))$  on the relaxation of any ordinal embedding into  $d$ -dimensional  $\ell_p$  space for fixed integers  $p$  or  $p = \infty$ . In particular, for  $d = \Theta(\lg n)$ , this lower bound is  $\Omega(\lg n / \lg \lg n)$ , leaving a gap between the upper bound of  $O(\lg n)$  which follows from Bourgain embedding. In contrast, for metric embeddings, there is an  $\Omega(\lg n)$  lower bound on distortion for  $d = \Theta(\lg n)$  [LLR95, Mat97].

We also develop approximation algorithms for finding the minimum possible relaxation for an ordinal embedding of a specified metric. Specifically, we give a 3-approximation for ordinal embedding of the shortest-path metric of a specified unweighted tree into the line. In contrast, only  $O(n^{1/3})$ -approximation algorithms are known for the same problem with distortion [BDG<sup>+</sup>05]. In general, approximation algorithms for embedding are a central challenge in the field, and few are known [HIL03, Iva00, B03, ABF<sup>+</sup>99, FK99, BDHI04]. We also expect that our techniques will extend to obtain approximation algorithms for more general ordinal embedding problems.

## 2 Definitions

In this section, we define ordinal embeddings and relaxation, as well as the standard notions of metric embeddings and distortion.

Consider a finite metric  $D : P \times P \rightarrow [0, \infty)$  on a finite point set  $P$ —the *source metric*—and a class  $\mathcal{T}$  of metric spaces  $(T, d) \in \mathcal{T}$  where  $d$  is the distance function for space  $T$ —the *target metrics*. An *ordinal embedding (with no relaxation)* of  $D$  into  $\mathcal{T}$  is a choice  $(T, d) \in \mathcal{T}$  of a target metric and a mapping  $\phi : P \rightarrow T$  of the points into the target metric such that every comparison between pairs of distances has the same outcome: for all  $p, q, r, s \in P$ ,  $D(p, q) \leq D(r, s)$  if and only if  $d(\phi(p), \phi(q)) \leq d(\phi(r), \phi(s))$ . Equivalently,  $\phi$  induces a monotone function  $D(p, q) \mapsto d(\phi(p), \phi(q))$ , and for this reason ordinal embeddings are also called *monotone embeddings*. An *ordinal embedding with relaxation  $\alpha$*  of  $D$  into  $\mathcal{T}$  is a choice  $(T, d) \in \mathcal{T}$  and a mapping  $\phi : P \rightarrow T$  such that every comparison between pairs of distances not within a factor of  $\alpha$  has the same outcome: for all  $p, q, r, s \in P$  with  $D(p, q)/D(r, s) > \alpha$ ,  $d(\phi(p), \phi(q)) > d(\phi(r), \phi(s))$ . Equivalently, we can view a relaxation  $\alpha$  as defining a partial order on distances  $D(p, q)$ , where two distances  $D(p, q)$  and  $D(r, s)$  are comparable if and only if they are not within a factor of  $\alpha$  of each other, and the ordinal embedding must preserve this partial order on distances.

An ordinal embedding with relaxation 1 is a different notion from ordinal embedding with no relaxation, because the former allows violation of equalities between pairs of distances. Indeed, we will show in Section 6.1 that the two notions have major qualitative differences. We define ordinal embedding with relaxation in this way, instead of making the  $> \alpha$  inequality non-strict, because otherwise our notion of relaxation 1 would have to be phrased as “relaxation  $1 + \varepsilon$  for any  $\varepsilon > 0$ ”. Another consequence is that we can define the minimum possible relaxation  $\alpha^* = \alpha^*(D, \mathcal{T})$  of an ordinal embedding of  $D$  into  $\mathcal{T}$ , instead of having to take an infimum. (The infimum will be realized provided the space  $\mathcal{T}$  is closed.)

We pay particular attention to contrasts between ordinal embedding and “standard” embedding,

which we call “metric embedding” for distinction. A *contractive metric embedding with distortion  $c$*  of a source metric  $D$  into a class  $\mathcal{T}$  of target metrics is a choice  $(T, d) \in \mathcal{T}$  and a mapping  $\phi : P \rightarrow T$  such that no distance increases and every distance is preserved up to a factor of  $c$ : for all  $p, q \in P$ ,  $1 \leq D(p, q)/d(\phi(p), \phi(q)) \leq c$ . Similarly, we can define an *expansive metric embedding with distortion  $c$*  with the inequality  $1 \leq d(\phi(p), \phi(q))/D(p, q) \leq c$ . When  $c = 1$ , these two notions coincide to require exact preservation of all distances; such an embedding is called a *metric embedding with no distortion* or an *isometric embedding*. In general,  $c^* = c^*(D, \mathcal{T})$  denotes the minimum possible distortion of a metric embedding of  $D$  into  $\mathcal{T}$ . (This definition is equivalent for both contractive and expansive metric embeddings, by scaling.)

### 3 Comparison between Distortion and Relaxation

The following propositions relate  $\alpha^*$  and  $c^*$ .

**Proposition 1** *For any source and target metrics,  $\alpha^* \leq c^*$ .*

**Proof:** Consider a contractive metric embedding  $\phi$  into  $(T, d)$  with distortion  $c$ . We show that  $\phi$  is also an ordinal embedding into  $(T, d)$  with relaxation  $\alpha \leq c$ . Consider a pair of distances  $D(p, q)$  and  $D(p', q')$  with ratio  $D(p, q)/D(p', q')$  larger than  $c$ . (Thus, in particular, we label  $p, q, p', q'$  so that  $D(p, q) > D(p', q')$ .) Then  $d(\phi(p), \phi(q))/d(\phi(p'), \phi(q')) \geq D(p, q)/(cD(p', q'))$  by expansiveness of  $D(p, q)$  and distortion of  $D(p', q')$ . Thus  $d(\phi(p), \phi(q))/d(\phi(p'), \phi(q')) > 1$ , so  $d(\phi(p), \phi(q)) > d(\phi(p'), \phi(q'))$  as desired.  $\square$

Next we show that  $c^*$  and  $\alpha^*$  can have an arbitrarily large ratio, even when the target metric is the real line.

**Proposition 2** *Embedding a uniform metric (where  $D(p, q) = 1$  for all  $p \neq q$ ) into the real line has  $c^* = n - 1$  and  $\alpha^* = 1$ .*

**Proof:** The mapping  $\phi(p) = 0$ , for all  $p \in P$ , is an ordinal embedding with no relaxation, because every distance remains equal (albeit 0). Any expansive metric embedding into the real line must have distance at least 1 between consecutively embedded points, so the entire embedding must occupy an interval of length at least  $n - 1$ . The two points embedded the farthest away from each other therefore have distance at least  $n - 1$ , for a distortion of at least  $n - 1$ . On the other hand, any embedding in which consecutively embedded points have distance exactly 1 has distortion  $n - 1$ .  $\square$

Next we give a general bound on  $\alpha^*$  that is essentially always finite. Define the *diameter*  $\text{diam}(D)$  of a metric  $D$  to be the ratio of the maximum distance to the minimum distance. (If the minimum distance is zero and the maximum distance is positive, then  $\text{diam}(D) = \infty$ ; if both are zero, then  $\text{diam}(D) = 1$ .)

**Proposition 3** *For any source metric  $D$  and any target metrics,  $\alpha^* \leq \text{diam}(D)$ .*

**Proof:** The mapping  $\phi(p) = 0$ , for all  $p \in P$ , has ordinal relaxation  $\text{diam}(G)$ , because all non-equal comparisons between distances are violated, and the largest ratio between any two distances is precisely  $\text{diam}(D)$ .  $\square$

No such general finite upper bound exists for  $c^*$ , as evidenced by “cut metrics”. A *cut metric* is defined by a partition  $P = A \cup B$  of the point set  $P$  into two disjoint sets  $A$  and  $B$ . The metric assigns a distance of 0 between pairs of points in  $A$  and pairs of points in  $B$ , and assigns a distance

of 1 between other pairs of points. If the source metric  $D$  has no zero distances and the target metrics are the cut metrics, then  $c^* = \infty$ , because some distance must become 0 which requires infinite distortion.

In contrast,  $\alpha^*$  remains at most  $\text{diam}(D)$ , and in some sense measures the quality of a clustering of the points into two clusters. Furthermore, the optimal  $\alpha^*$  and clustering can be computed efficiently:

**Proposition 4** *The minimum-relaxation ordinal embedding of a specified metric into a cut metric can be computed in polynomial time.*

**Proof:** First we guess the optimal relaxation  $\alpha^*$  among  $O(n^4)$  possibilities (the ratio of the distance between any two pairs of points). Second we guess a pair  $(p, q)$  of points on different sides of the cut and with minimum distance  $D(p, q)$ . Thus all pairs  $(r, s)$  of points with smaller distance  $D(r, s) < D(p, q)$  must have  $r$  and  $s$  on the same side of the cut. Also, if there is any ordinal embedding of relaxation  $\alpha^*$ , there cannot be pairs  $(r, s)$  of points with distance larger by a factor of  $\alpha^*$ , i.e., with  $D(r, s) > \alpha^* D(p, q)$ , because such distances will be mapped to a distance smaller or equal to 1, the mapped distance of  $(p, q)$ . Similarly, there cannot be pairs  $(r, s)$  and  $(r', s')$  of points with distance less than  $D(p, q)$  and with  $D(r, s) > \alpha^* D(r', s')$ , because those pairs are forced to map to equal distances of 0. Finally, all pairs  $(r, s)$  of points with  $D(p, q) \leq D(r, s) \leq \alpha^* D(p, q)$  must have  $r$  and  $s$  on different sides of the cut if there is another distance  $D(r', s') < D(r, s)/\alpha^*$ , and otherwise are unconstrained.

All constraints of the form “ $r$  and  $s$  must be on the same side of the cut” and “ $r$  and  $s$  must be on different sides of the cut” can be phrased as a 2-SAT instance. Each point  $r$  has a variable  $x_r$  which is 0 if it placed in set  $A$  and 1 if it placed in set  $B$ . Each constraint thus has the form  $x_r = x_s$  or  $x_r \neq x_s$ , which can be phrased in 2-CNF. Thus we can find an ordinal embedding into a cut metric with relaxation at most the guessed value of  $\alpha^*$ , if one exists.  $\square$

Next we consider the related problem of ordinal embedding into the real line, which is a generalization of cut metrics. First we show that we can decide whether  $\alpha^* = 1$  in this case. The algorithm requires more sophistication (namely, guessing) than the trivial algorithm for metric embedding with distortion 1, where one can incrementally build an embedding in any Euclidean space in linear time.

**Proposition 5** *In polynomial time, we can decide whether a given metric can be ordinally embedded into the line with relaxation 1.*

**Proof:** The algorithm guesses the leftmost point  $p$  and greedily places every point  $q$  at position  $D(p, q)$  on the line. (In particular, the algorithm places  $p$  at position 0.) It is easy to show that this embedding has ordinal relaxation 1 whenever such an embedding exists.  $\square$

Next we consider the worst case for ordinal embedding into the line. We show in particular that the cycle requires large relaxation. The cycle also requires large distortion into the line, but the proof technique for ordinal relaxation is very different from the usual “packing argument” that suffices for metric distortion.

**Proposition 6** *Ordinal embedding of the shortest-path metric of an unweighted cycle of even length  $n$  into the line requires relaxation at least  $n/2$ .*

**Proof:** Suppose to the contrary that there is an ordinal embedding  $\phi$  of the cycle into the line with relaxation less than  $n/2$ . Label the vertices of the cycle 1 through  $n$  in cyclic order. Assume without

loss of generality that  $\phi(1) < \phi(n/2 + 1)$ . We must also have  $\phi(2) < \phi(n/2 + 1)$ , because otherwise  $|\phi(2) - \phi(1)| \geq |\phi(n/2 + 1) - \phi(1)|$ , contradicting that  $\alpha < n/2$ . Similarly,  $\phi(2) < \phi(n/2 + 2)$ , because otherwise  $|\phi(n/2 + 2) - \phi(n/2 + 1)| \geq |\phi(n/2 + 2) - \phi(2)|$ , again contradicting that  $\alpha < n/2$ . Repeating this argument shows that  $\phi(3) < \phi(n/2 + 3)$ , etc., and finally that  $\phi(n/2 + 1) < \phi(1)$ , a contradiction.  $\square$

Section 5 shows that some trees also require  $\Omega(n)$  ordinal relaxation into the line.

## 4 $\ell_p$ Metrics are Universal

In this section we show that every distance matrix can be ordinally embedded without relaxation into  $\ell_p$  space of a polynomial number of dimensions, for any fixed  $1 \leq p \leq \infty$ . This result is surprising in comparison to metric embeddings. Every metric can be embedded into  $\ell_p$  using  $O(\lg n)$  distortion [Bou85, LLR95], and in the worst case  $\Omega(\lg n)$  distortion is necessary for any  $p < \infty$ , as proved in [LLR95] for  $p = 2$  and in [Mat97] for all other values of  $p$ . In particular, the shortest-path metric of a constant-degree expander graph requires  $\Omega(\lg n)$  distortion.

**Theorem 1** *Every distance matrix can be ordinally embedded without relaxation into  $O(n^5)$ -dimensional  $\ell_p$  space, for any fixed  $1 \leq p \leq \infty$ .*

The same result was established independently in [BL04] using an algebraic proof. Specifically, they show that every distance matrix can be ordinally embedded into  $(n - 1)$ -dimensional Euclidean space, and then use the property that any Euclidean metric can be isometrically embedded into any  $\ell_p$  space with at most  $\binom{n}{2}$  dimensions. In contrast, our proof is purely combinatorial.

We can also reduce the number of dimensions for some values of  $p$ . For example, for  $p = 2$ , a simple rotation reduces the number of dimensions to  $n - 1$ .

Our proof proceeds in two steps. First we show that 0/1 Hamming metrics are universal in the same sense as Theorem 1. To conclude the proof, we note that there is an ordinal embedding without relaxation from 0/1 Hamming metrics into any  $\ell_p$  metric. In fact, the  $p$ th root of the distances in a 0/1 Hamming metric can be metrically embedded without distortion into  $\ell_p$  with the same number of dimensions. This second part is merely an observation, so the main work is in showing that 0/1 Hamming metrics are universal:<sup>2</sup>

**Lemma 1** *Every distance matrix can be ordinally embedded without relaxation into a 0/1 Hamming metric with  $O(n^5)$  dimensions. In other words, any desired ordering on the distances between pairs of  $n$  points can be realized by a 0/1 Hamming metric on those  $n$  points.*

**Proof:** Given a partial order  $\mathcal{P}$  on a set of distances, we construct a 0/1 Hamming metric  $H$  such that  $P_{i,j} < P_{k,l}$  implies  $H_{i,j} < H_{k,l}$ . If  $\mathcal{P}$  is non-total, then we can take any topological sort of  $\mathcal{P}$  and realize it as a Hamming metric. This ordinal embedding will satisfy the original partial order, so from now on, we assume that  $\mathcal{P}$  is a total order. Because  $\mathcal{P}$  is an order on distances, defined by pairs of points, we can define it as a sequence of pairs  $\mathcal{P} = [(a_0, b_0), (a_1, b_1), \dots, (a_{\binom{n}{2}}, b_{\binom{n}{2}})]$ , where in each pair, we arbitrarily select which node is  $a$  and which is  $b$ .

<sup>2</sup>Note that finite 0/1 Hamming metrics and finite Hamming metrics are essentially the same, because one can be converted into the other with a dimension blowup that is multiplicative in the number of points. Thus our result could have been established with general Hamming metrics. However, our construction directly yields a 0/1 Hamming metric, so we do not need this extra conversion detail.

We now must produce a 0/1 vector for each point of the space so that the Hamming metric induced preserves the order  $\mathcal{P}$ . We assume that  $n$  is a power of 2; otherwise we can simply round  $n$  up to the next power of two.

Our main tool will be Hadamard matrices, defined as follows. Let  $H_0 = [0]$ , and

$$H_{i-1} = \begin{pmatrix} H_i & H_i \\ H_i & \overline{H_i} \end{pmatrix}$$

where  $\overline{H_i}$  is the bitwise negation of  $H_i$ . Notice that the first row is the all-0 vector, denoted  $\vec{0}$ . Also, each row other than the first row consists of half 0s and half 1s. More strikingly, any two rows of  $H_i$  have Hamming distance  $2^{i-1}$ , that is, they differ in half their positions. Finally,  $\vec{1}$  has Hamming distance  $2^{i-1}$  from any row except the first row, with which it has Hamming distance  $2^i$ .

We generate a set of dimensions that code for each distance  $\mathcal{P}_i$  and concatenate all the dimensions at the end. To code for distance  $\mathcal{P}_i = (a_i, b_i)$ , we set  $a_i$ 's bits to be  $0^{in}$  and  $b_i$ 's bits to be  $1^{in}$ . Every other point in the space besides these two gets a distinct row from the Hadamard matrix, repeated  $i$  times. Now the induced distances are  $in/2$  for any pair of points except for  $a_i$  and  $b_i$ , which are at distance  $in$ .

Let the total number of dimensions be  $d = n \binom{n}{2} (\binom{n}{2} + 1)/2$ .

Consider now the distances between any pair  $a$  and  $b$  resulting from the concatenation of all  $d$  dimensions, and assume that  $a = a_i$  and  $b = b_i$ , that is, their pairwise distance is the  $i$ th in the list. Then their pairwise distance is  $(d + in)/2$ . Thus, this embedding assigns to the  $i$ th smallest distance in  $\mathcal{P}$  the  $i$ th smallest distance in the Hamming metric.  $\square$

## 5 Approximation Algorithms for Unweighted Trees into the Line

In this section, we give a 3-approximation algorithm for ordinal embedding the shortest-path metric induced by an unweighted tree into the line with approximately minimum relaxation. In contrast, the best approximation algorithm known for metrically embedding trees into the line with approximately minimum distortion is a recently discovered  $O(n^{1/3})$ -approximation [BDG<sup>+</sup>05].

First we find a structure for proving lower bounds on the optimal relaxation:

**Lemma 2** *Given  $n$  such that 3 divides  $n - 1$ , ordinal embedding of the shortest-path metric of an unweighted 3-spider with  $(n - 1)/3$  vertices on each leg of the spider (i.e., a 3-star with each edge subdivided into a path of  $(n - 1)/3$  edges) requires relaxation at least  $(n - 1)/3$ .*

**Proof:** Suppose to the contrary that there is an ordinal embedding  $\phi$  of the 3-spider into the line with relaxation  $\alpha < (n - 1)/3$ . Label the vertices as follows: 0 denotes the root, and  $a_1, \dots, a_{(n-1)/3}$ ,  $b_1, \dots, b_{(n-1)/3}$ , and  $c_1, \dots, c_{(n-1)/3}$  denote the nodes on the legs of the spider in order of their distance from the root 0. Because  $\alpha < (n - 1)/3$ ,  $|\phi(a_{(n-1)/3}) - \phi(0)| > 0$ , and the same holds for  $b_{(n-1)/3}$  and  $c_{(n-1)/3}$ . Because the spider has three legs, two of  $a_{(n-1)/3}, b_{(n-1)/3}, c_{(n-1)/3}$  are on the same side of the root 0 on the line. Without loss of generality, assume that the  $a$  and  $b$  legs are both to the right of 0, and that  $\phi(a_{(n-1)/3}) \geq \phi(b_{(n-1)/3}) > \phi(0)$ . Let  $k$  be such that  $\phi(a_k) < \phi(b_{(n-1)/3}) < \phi(a_{k+1})$  (where the label  $a_0$  refers to the root 0). Such a  $k$  exists because  $\alpha < (n - 1)/3$ , so  $\phi(a_k) \neq \phi(b_{(n-1)/3})$  for all  $k$ , and because  $\phi(0) < \phi(b_{(n-1)/3}) < \phi(a_{(n-1)/3})$ . It follows that  $|\phi(b_{(n-1)/3}) - \phi(a_{k+1})| < |\phi(a_{k+1}) - \phi(a_k)|$ . In contrast, in the 3-spider graph,  $b_{(n-1)/3}$  and  $a_{k+1}$  have distance at least  $(n - 1)/3$ , and  $a_{k+1}$  and  $a_k$  have distance 1. Therefore  $\alpha > (n - 1)/3$ .  $\square$

**Definition 1** Given a tree  $T$ , a tripod  $(a, b, c)$  is the union of shortest paths in  $T$  connecting every pair of vertices among  $\{a, b, c\}$ . The root  $r$  of the tripod is the common vertex among all three shortest paths. The length of the tripod is  $k = \min\{D(r, a), D(r, b), D(r, c)\}$ .

Any tripod of length  $k$  induces a 3-spider with  $k$  vertices on each leg, by truncating all longer arms of the tripod to length  $k$ . Thus by Lemma 2, any tree with a tripod of length  $k$  must have ordinal relaxation at least  $k$ . Using this lower bound, we obtain a constant-factor approximation algorithm.

**Theorem 2** Given a tree  $T$ , there is an ordinal embedding  $\phi : T \rightarrow \mathbb{R}$  of  $T$  into the line with relaxation  $2k + 1$ , where  $k$  is the length of the largest tripod of  $T$ . The embedding can be computed in polynomial time.

**Proof:** If there are at most two leaves in the tree  $T$ , then  $T$  can be trivially embedded into the line without distortion or relaxation. Otherwise,  $T$  has a tripod. Let  $(A, B, C)$  be a longest tripod, let  $r$  be its root, and let  $k$  be its length. We view  $T$  as rooted at  $r$ . Let  $(a, b, c)$  be a tripod rooted at  $r$  that maximizes  $D(r, a) + D(r, b) + D(r, c)$ . This tripod corresponds to taking the longest three paths starting from different neighbors of  $r$ . In particular all three paths have length at least  $k$ , so the tripod  $(a, b, c)$  has length  $k$ . Relabel  $\{a, b, c\}$  so that  $D(r, a) = k$ .

**Claim 1** For any  $d \in \{a, b, c\}$ , for any  $d' \neq r$  on the path from  $r$  to  $d$ , and for any descendant  $x$  of  $d'$ ,  $D(d', x) \leq D(d', d)$ .

**Proof:** Assume, to the contrary, that  $D(d', x) > D(d', d)$ . If  $d = a$ , then there would be a larger tripod  $(x, b, c)$  rooted at  $r$ . Otherwise, assume without loss of generality that  $d = b$ . Then there would be a tripod  $(a, x, c)$ , of the same length, and such that  $D(r, a) + D(r, x) + D(r, c) > D(r, a) + D(r, b) + D(r, c)$ , a contradiction.  $\square$

**Claim 2** For any  $d \in \{b, c\}$ , for any  $d' \neq r$  on the path from  $r$  to  $d$ , and for any descendant  $x$  of  $d'$ , such that the path from  $x$  to  $d'$  intersects the path from  $r$  to  $d$  only at vertex  $d'$ ,  $D(d', x) \leq k$ .

**Proof:** Suppose to the contrary that  $D(d', x) > k$ . By the definition of  $d'$ ,  $D(d', a) > D(r, a) = k$ . By Claim 1,  $D(d', d) \geq D(d', x)$ . If  $D(d', d) \leq k$ , then  $D(d', x) \leq D(d', d) \leq k$ , a contradiction. If  $D(d', d) > k$ , then the tripod  $(x, d, a)$  (rooted at  $d'$ ) has length at least  $k + 1$ , which is again a contradiction.  $\square$

Now we construct the embedding  $\phi$  as follows. For every vertex  $x$  on the shortest path between  $b$  and  $c$ , we contract every subtree that intersects the path only at  $x$  into the single vertex  $x$ . The resulting graph is the same path from  $b$  to  $c$ , but where each vertex represents several vertices of the original graph. We embed this path into the line, placing the  $i$ th vertex along the path at coordinate  $i$ . This embedding places several vertices of the original graph at the same point in the line.

We claim that the depth of each contracted tree is at most  $k$ . For each subtree rooted at  $r$  (e.g., the one containing  $a$ ), no vertex  $x$  in the subtree can have  $D(r, x) > k$  because then we could have chosen that vertex as  $a$  and increase the objective function  $D(r, a) + D(r, b) + D(r, c)$ , a contradiction. For each subtree rooted at another node  $b' \neq r$  on the path from  $b$  to  $c$ , we can apply Claim 2 and obtain that  $D(b', x) \leq k$  for any vertex  $x$  in the subtree rooted at  $b'$ . Therefore the depth of each contracted tree is at most  $k$ .

Finally we claim that the ordinal relaxation of this mapping is at most  $2k + 1$ . Consider two vertices  $x$  and  $y$  belonging to contracted subtrees rooted at  $s$  and  $t$ , respectively. Their original distance is at most  $2k + D(s, t)$ , and their new distance is  $D(s, t)$ . Therefore the distance changes order with respect to distances at least  $D(s, t)$ , for a worst-case ratio of  $(2k + D(s, t))/D(s, t)$ . This ratio is maximized when  $D(s, t) = 1$  in which case it is  $2k + 1$ .  $\square$

**Corollary 1** *There is a polynomial-time algorithm to find  $\phi$  of Theorem 2. The algorithm is a 3-approximation algorithm for ordinally embedding trees into a line.*

**Proof:** The proof of Theorem 2 is constructive, thus it gives an algorithm. Since the length of the largest tripod is a lower bound of embedding ordinally the tripod into a line, we obtain that the algorithm is a  $(2 + 1/k)$ -approximation algorithm.  $\square$

## 6 Ultrametrics

In this section we establish several results about ordinal embedding when the source metric or the target metrics are ultrametrics.

### 6.1 Ultrametrics into $\ell_p$ with Logarithmic Dimensions

First we demonstrate that ultrametrics can be ordinally embedded into  $O(\lg n)$ -dimensional  $\ell_p$  space, for any fixed  $1 \leq p \leq \infty$ , with relaxation 1. Here we exploit the minor difference between “relaxation 1” and “no relaxation”—that equality constraints can be violated—because, as described in the introduction, any ordinal embedding without relaxation of any ultrametric into Euclidean space requires  $n - 1$  dimensions. Thus the ordinal dimension of an ultrametric is “just barely”  $n - 1$ ; the slightest relaxation allows us to obtain a much better embedding. Our result also contrasts metric embeddings where ultrametrics can be embedded into Euclidean space with  $1 + \varepsilon$  distortion, but such an embedding requires  $\varepsilon^{-2} \lg n$  dimensions [BM04]. The number of dimensions in our ordinal embeddings is independent of any such  $\varepsilon$ .

Our construction is based on monotone stretching of the discrepancy between different distances:

**Lemma 3** *For any  $k > 1$ , and for any ultrametric  $M = (P, D)$ , there is an ultrametric  $M' = (P, D')$  such that, for any  $p, q, r, s \in P$ , if  $D(p, q) = D(r, s)$ , then  $D'(p, q) = D'(r, s)$ , and if  $D(p, q) > D(r, s)$ , then  $D'(p, q) \geq kD'(r, s)$ .*

**Proof:** Because  $M$  is an ultrametric, we can construct a weighted tree  $T$ , with  $P$  forming the set of leaves, such that the weights are nondecreasing along any path of  $T$  starting from the root. Moreover, for any  $u, v \in P$ , the ultrametric distance  $D(u, v)$  is equal to the maximum weight of an edge along the path from  $u$  to  $v$  in  $T$ .

For  $u, v \in P$ , define  $r(D(u, v)) = i$  where  $D(u, v)$  is equal to the  $i$ th smallest distance in  $M$ . Consider now the weighted tree  $T'$  obtained from  $T$  by replacing an edge of weight  $w$  by an edge of weight  $k^{r(w)}$ . Let  $M'$  be the resulting ultrametric induced by  $T'$ . If  $D(p, q) = D(r, s)$ , then  $r(D(p, q)) = r(D(r, s))$ , so  $D'(p, q) = D'(r, s)$ . Finally, if  $D(p, q) > D(r, s)$ , then  $r(D(p, q)) \geq r(D(r, s)) + 1$ , so  $D'(p, q) \geq kD'(r, s)$ .  $\square$

We combine this lemma with a result for the metric case:

**Lemma 4 (Bartal and Mendel [BM04])** *For any  $1 \leq p \leq \infty$ , any  $n$ -point ultrametric can be metrically embedded into  $O(\varepsilon^{-2} \lg n)$ -dimensional  $\ell_p$  space with distortion at most  $1 + \varepsilon$ .*

Now we are ready to prove the main result of this subsection:

**Theorem 3** *For any  $1 \leq p \leq \infty$ , any  $n$ -point ultrametric can be ordinally embedded into  $O(\lg n)$ -dimensional  $\ell_p$  space with relaxation 1.*

**Proof:** Given an ultrametric  $M = (P, D)$ , by Lemma 3, we can obtain an ultrametric  $M' = (P, D')$  such that, for any  $p, q, r, s \in P$ , if  $D(p, q) = D(r, s)$ , then  $D'(p, q) = D'(r, s)$ , and if  $D(p, q) > D(r, s)$ , then  $D'(p, q) \geq 2D'(r, s)$ . Applying Lemma 4 with  $\varepsilon = 1/2$ , we obtain a contractive metric embedding  $\phi$  of  $P$  into  $O(\lg n)$ -dimensional  $\ell_p$  space such that, for any  $p, q, r, s \in P$ , if  $D(p, q) > D(r, s)$ , then  $\|\phi(p) - \phi(q)\| \geq \frac{2}{3}D'(p, q) \geq \frac{4}{3}D'(r, s) \geq \frac{4}{3}\|\phi(r) - \phi(s)\|$ . Therefore  $\phi$  is an ordinal embedding with relaxation 1.  $\square$

## 6.2 Arbitrary Distance Matrices into Ultrametrics

In this subsection, we give a polynomial-time algorithm for computing an ordinal embedding of an arbitrary metric into an ultrametric with minimum possible relaxation.

We will show that the optimal ordinal embedding of a distance matrix  $M$  into an ultrametric is the *subdominant* of  $M$  [FKW95]. One recursive construction of the subdominant is as follows. First, we compute a partition  $P = P_1 \cup P_2 \cup \dots \cup P_k$ , for some  $k \geq 2$ , such that the minimum distance between any  $P_i$  and  $P_j$  is maximized. Such a partition can be found by computing a minimum spanning tree  $T$  of  $M$ , and partitioning the points by removing all the edges of  $T$  of maximum length. Let  $\Delta$  be the maximum distance between any two points in  $P$ . We create a hierarchical tree representation for an ultrametric by starting with a root  $v_P$  and  $k$  children  $v_{P_1}, \dots, v_{P_k}$ . The length of the edge  $\{v_P, v_{P_i}\}$  is equal to  $\Delta$  for each  $i \in \{1, 2, \dots, k\}$ . We recursively compute hierarchical tree representations for the metrics induced by the point sets  $P_1, P_2, \dots, P_k$ , and then we merge these trees by identifying, for each  $i \in \{1, 2, \dots, k\}$ , the root of the tree for  $P_i$  with the node  $v_{P_i}$ . In fact this entire construction can be carried out with a single computation of the minimum spanning tree, and thus takes linear time.

**Lemma 5** *Let  $\Delta = \max_{p, q \in P} D(p, q)$  and let  $\delta$  be the minimum distance between two points in different sets  $P_i$  and  $P_j$ . Then any ordinal embedding has relaxation at least  $\Delta/\delta$ .*

**Proof:** Suppose that the maximum distance  $\Delta$  is attained by points  $u, v$  with  $u \in P_i$  and  $v \in P_j$ , where  $i \neq j$ . Consider an optimal ordinal embedding  $\phi$  of  $M$  into a hierarchical tree representation  $T$  of an ultrametric. Thus the distance between two leaves  $p$  and  $q$  is equal to the maximum length of an edge along the unique path between  $p$  and  $q$ . No matter how  $\phi$  splits  $P$  into subsets at the root of  $T$ , there exist  $r, s \in P$  such that  $D(r, s) = \delta$  and the path from  $r$  to  $s$  in  $T$  visits the root of  $T$ . Thus the path from  $r$  to  $s$  passes through the maximum edge in  $T$ . Hence, the maximum distance along the path between  $u$  and  $v$  in  $T$  cannot be larger than the maximum distance along the path between  $r$  and  $s$  in  $T$ . Therefore  $d(\phi(u), \phi(v)) \leq d(\phi(r), \phi(s))$ , while  $D(u, v) = \Delta > \delta = D(r, s)$ , so the relaxation is at least  $\Delta/\delta$ .  $\square$

**Theorem 4** *Given any distance matrix  $M$ , we can compute in polynomial time an optimal ordinal embedding of  $M$  into an ultrametric.*

**Proof:** Let  $\phi$  be the ordinal embedding of  $M = (P, D)$  computed by the algorithm, with a hierarchical tree representation  $T$ . The maximum relaxation  $\alpha$  of  $\phi$  is attained for some  $p, q, r, s \in P$  such that  $D(p, q) \geq \alpha D(r, s)$  and  $d(\phi(p), \phi(q)) < d(\phi(r), \phi(s))$ . It follows that there exists an

internal node  $v$  of  $T$ , with children  $v_1$  and  $v_2$ , such that leaves  $p$  and  $q$  are descendants of  $v_1$ , while only one of the leaves  $r$  or  $s$  is a descendant of  $v_1$ . Assume without loss of generality that  $r$  is a descendant of  $v_1$  and  $s$  is a descendant of  $v_2$ .

Consider the recursive call of the algorithm on a subset of points  $P' \subseteq P$  in which the node  $v$  was created. Because  $r$  and  $s$  are in different subtrees of  $v$ , it follows that, in the partition of the set  $P'$  of points computed by the algorithm, the minimum distance between distinct sets is at most  $D(r, s)$ . On the other hand, the maximum distance between pairs of points in  $P'$  is at least  $D(p, q)$ . Thus, by Lemma 5, the optimal relaxation for ordinal embedding of  $M$  into an ultrametric is at least  $D(p, q)/D(r, s) \geq \alpha$ .  $\square$

By a similar argument it can be shown that the same algorithm also computes a metric embedding of  $M$  into an ultrametric with minimum possible distortion. Furthermore, the distortion is equal to the relaxation in this embedding. In the next section we show that ultrametries are essentially the only case where this can happen universally.

### 6.3 When Distortion Equals Relaxation

Finally we show that, in a certain sense, ultrametries are the only target metrics that have equal values of  $\alpha^*$  and  $c^*$ , or even a universally bounded ratio between  $\alpha^*$  and  $c^*$ .

**Theorem 5** *If a set  $\mathcal{T}$  of target metrics is closed under inclusion (i.e., closed under taking the submetric induced on a subset of points), and there is a constant  $C$  such that every distance matrix  $D$  has  $c^*/\alpha^* \leq C$  (when embedding  $D$  into  $\mathcal{T}$ ), then every metric in  $\mathcal{T}$  is an ultrametric.*

**Proof:** Consider any metric  $M$  in  $\mathcal{T}$ . We claim that  $M$  has more than one diameter pair. Suppose to the contrary that only  $p$  and  $q$  attain the maximum distance in  $M$ . Let  $M_{+d}$  be the distance matrix identical to  $M$  except for  $M_{+d}(p, q) = M(p, q) + d$ . Let  $d$  be any positive real greater than the sum of the second- and third-largest distances. Then  $M_{+d}$  is not in  $\mathcal{T}$  because it violates the triangle inequality and  $\mathcal{T}$  is a family of metrics. Because no other distance in  $M$  is equal to  $M(p, q)$ ,  $M_{+d}$  can be ordinally embedded with no relaxation into  $\mathcal{T}$  simply by taking  $M$ . However,  $M_{+d}$  cannot be metrically embedded into  $\mathcal{T}$  without distortion, because  $M_{+d}$  is not in  $\mathcal{T}$ . Furthermore  $M_{+cd}$  cannot be metrically embedded into  $\mathcal{T}$  with distortion less than  $c$ , because any contractive metric embedding must reduce the distance between  $p$  and  $q$  by a factor of  $c$ . Therefore the ratio between the minimum metric distortion  $c^*$  and the minimum ordinal relaxation  $\alpha^*$  cannot be bounded.

Now by inclusion, any submetric of  $M$  induced by three points is also in  $\mathcal{T}$ , and therefore has a non-unique maximum edge. Thus all triangles in  $M$  are tall isosceles, which is one characterization of  $M$  being an ultrametric.  $\square$

In fact, this theorem needs only that the set  $\mathcal{T}$  of target metrics is closed under taking the induced metric on any triple of points.

## 7 Worst Case of Unweighted Trees into Euclidean Space

In this section, we consider the worst-case relaxation required for ordinal embedding of the shortest-path metric of an unweighted tree  $T$  into  $d$ -dimensional  $\ell_2$  space. Our work is motivated by that of Gupta [Gup00] and Babilon, Matoušek, Maxová, and Valtr [BMMV03]. We show that, for any  $d \geq 2$ , and for any unweighted tree  $T$  on  $n$  nodes,  $\alpha^* = \tilde{O}(n^{1/d})$ . We complement this result by exhibiting a family of trees with optimal ordinal relaxation  $\Omega(n^{1/(d+1)})$ . In contrast, the best bounds on the worst-case distortion required are  $\tilde{O}(n^{1/(d-1)})$  and  $\Omega(n^{1/d})$  [Gup00]. These ranges

overlap at the endpoint of  $\tilde{\Theta}(n^{1/d})$ , but it seems that ordinal embedding and metric embedding behave fundamentally differently, in particular because different proof techniques are required for both the upper and lower bounds.

First we prove the upper bound. At a high level, the algorithm finds nodes that can be contracted to a single point, which can be an effective ordinal embedding, unlike metric embedding where it causes infinite distortion.

**Theorem 6** *Any weighted tree can be ordinally embedded into  $d$ -dimensional  $\ell_2$  space with relaxation  $\tilde{O}(n^{1/d})$ .*

**Proof:** Let  $T = (V(T), E(T))$  be an unweighted tree with  $|V(T)| = n$ . We show how to obtain an ordinal embedding of  $T$  into  $d$ -dimensional  $\ell_2$  space with relaxation  $\tilde{O}(n^{1/d})$ .

We construct a new tree  $T'$  as follows. Initially, we set  $T'_0 := T$ . For  $i = 1, \dots, n^{1/d}$ , we repeat the following process: Set  $T'_i := T'_{i-1}$ . For any leaf  $v$  of  $T'_{i-1}$ , we remove  $v$  from  $T'_{i-1}$ . Let  $T' := T'_{n^{1/d}}$ .

Define the function  $p : V(T) \rightarrow V(T')$ , such that for any  $v \in V(T) \setminus V(T')$ ,  $p(v)$  is the node in  $V(T')$ , which is closest to  $v$ , and for any  $v \in V(T')$ ,  $p(v) = v$ . It is easy to see that for every leaf  $v$  of  $T'$ , there are at least  $n^{1/d}$  nodes  $u \in V(T) \setminus V(T')$ , with  $p(u) = v$ . Thus, the number of leaves of  $T'$  is at most  $n^{\frac{d-1}{d}}$ .

It follows that using Gupta's algorithm [Gup00], we can compute an expansive metric embedding  $\phi'$  of  $T'$  into  $d$ -dimensional  $\ell_2$  space with distortion at most  $kn^{1/d}$ , for some  $k = \text{polylog}(n)$ . To obtain an embedding  $\phi$  of  $T$ , we simply set  $\phi(v) = \phi'(p(v))$  for each  $v \in V(T)$ .

It remains to show that  $\phi'$  has ordinal relaxation  $\tilde{O}(n^{1/d})$ . Let  $v_1, v_2, v_3, v_4 \in V(T)$ , with  $v_3 \neq v_4$  and

$$d_T(v_1, v_2) > (2 + k)n^{1/d}d_T(v_3, v_4).$$

We have

$$\begin{aligned} \|\phi(v_1) - \phi(v_2)\| &= \|\phi'(p(v_1)) - \phi'(p(v_2))\| \\ &\geq d_{T'}(p(v_1), p(v_2)) \\ &\geq d_T(v_1, v_2) - 2n^{1/d} \\ &> (2 + k)n^{1/d}d_T(v_3, v_4) - 2n^{1/d} \\ &\geq kn^{1/d}d_T(v_3, v_4) \\ &\geq kn^{1/d}d_{T'}(p(v_3), p(v_4)) \\ &\geq \|\phi'(p(v_3)) - \phi'(p(v_4))\| \\ &= \|\phi(v_3) - \phi(v_4)\|. \end{aligned}$$

Thus, we obtain that  $\phi$  has ordinal relaxation at most  $(2 + k)n^{1/d} = \tilde{O}(n^{1/d})$ .  $\square$

Next we prove the worst-case lower bound. The main novelty here is a new packing argument for bounding relaxation. Let  $F(m, L)$  denote the  $m$ -spider with arms of length  $L$ , that is, an  $m$ -star with each edge refined into a path of length  $L$ .

**Lemma 6** *Any ordinal embedding of  $F(m, L)$  into  $d$ -dimensional  $\ell_2$  space requires relaxation  $\Omega(\min\{L, m^{1/d}\})$ .*

**Proof:** Let  $T = F(m, L)$ , and let  $r \in V(T)$  be the only vertex of  $T$  with degree greater than 2. For any  $i$ , with  $0 \leq i \leq L$ , let  $U_i = \{v \in V(T) \mid d_T(r, v) = i\}$ .

Consider an optimal embedding  $\phi : V(T) \rightarrow \mathbb{R}^d$  with relaxation  $\alpha$ . We define

$$\begin{aligned}\mu_i &= \min_{u,v \in V(T)} \{\|\phi(u) - \phi(v)\| \mid d_T(u,v) = i\}, \\ \lambda_i &= \max_{u,v \in V(T)} \{\|\phi(u) - \phi(v)\| \mid d_T(u,v) = i\}.\end{aligned}$$

Observe that, if  $\mu_{2L} = 0$ , then there exist  $u, v \in U_L$  such that  $\phi(u) = \phi(v)$ . It follows that, if  $\alpha < 2L$ , then for any  $\{x, y\} \in E(T)$ ,  $\phi(x) = \phi(y)$ , which implies that all the vertices are mapped to the same point, and thus  $\alpha = \Omega(L)$ .

It remains to show that the assertion is true in the case  $\mu_{2L} > 0$ . Consider the nodes of  $U_L$ . For any  $u, v \in U_L$ , we have  $d_T(u, v) = 2L$ , and thus  $\|\phi(u) - \phi(v)\| \geq \mu_{2L}$ . For any  $v \in U_L$ , let  $B_v$  be the ball of radius  $\mu_{2L}/2$  centered at  $\phi(v)$ . It follows that, for any  $u, v \in U_L$ , the balls  $B_u, B_v$  can intersect only on their boundary. Thus,

$$\begin{aligned}\left| \bigcup_{v \in U_L} B_v \right| &= \sum_{v \in U_L} |B_v| \\ &= \Omega(m\mu_{2L}^d)\end{aligned}$$

By a packing argument, we obtain that there exist  $u, v \in U_L$  such that  $\|\phi(u) - \phi(v)\| = \Omega(m^{1/d}\mu_{2L})$ , which implies

$$\lambda_{2L} = \Omega(m^{1/d}\mu_{2L}). \quad (1)$$

Now consider two nodes  $u, v \in U_L$  such that  $\|\phi(u) - \phi(v)\| = \lambda_{2L}$ , and let  $p$  be the path from  $u$  to  $v$  in  $T$ . It follows that there exist nodes  $x, y \in p$  with  $d_T(x, y) = 2L/\alpha$  and  $\|\phi(x) - \phi(y)\| \geq \lambda_{2L}/\alpha$ . Thus

$$\lambda_{2L/\alpha} \geq \frac{\lambda_{2L}}{\alpha}. \quad (2)$$

Also, by the definition of the ordinal relaxation, we have

$$\mu_{2L} > \lambda_{2L/\alpha}. \quad (3)$$

Combining (1), (2), and (3), we obtain  $\alpha\lambda_{2L/\alpha} = \Omega(m^{1/d}\mu_{2L}) = \Omega(m^{1/d}\lambda_{2L/\alpha})$ . Thus we have shown that, if  $\mu_{2L} > 0$ , then  $\alpha = \Omega(m^{1/d})$ . The lemma follows.  $\square$

**Theorem 7** *For any  $n > 0$  and any  $d \geq 2$ , there is a tree  $T$  on  $n$  nodes for which every ordinal embedding has relaxation  $\Omega(n^{1/(d+1)})$ .*

**Proof:** The theorem follows from Lemma 6, for  $T = F(n^{d/(d+1)}, n^{1/(d+1)})$ .  $\square$

## 8 Arbitrary Metrics into Low Dimensions

By Lemma 1, a general  $O(\lg n)$  upper bound on relaxation carries over from metric embeddings of any  $n$ -point metric space into  $O(\lg n)$ -dimensional Euclidean space, using theorems of Bourgain and of Johnson and Lindenstrauss. For metric distortion, this bound is tight [LLR95], but one might suspect that the ordinal relaxation can be smaller. Here we show that it cannot be much smaller: some  $n$ -point metric spaces require relaxation  $\Omega(\log n / \log \log n)$ . This claim is a special case of the following result.

**Theorem 8** *There is an absolute constant  $c > 0$  such that, for every  $d$  and  $n$ , there is a metric space  $T$  on  $n$  points such that the relaxation of any ordinal embedding of  $T$  into  $d$ -dimensional Euclidean space is at least  $\frac{\log n}{\log d + \log \log n + c} - 1$ .*

The proof is based on two known results. The first is a bound of Warren on the number of sign patterns of a system of real polynomials. The second is the existence of dense graphs with no short cycles. We first state these two results.

Let  $P_j = P_j(x_1, \dots, x_\ell)$ ,  $j = 1, \dots, m$ , be  $m$  real polynomials. For a point  $u = (u_1, \dots, u_\ell) \in \mathbb{R}^\ell$ , the *sign pattern* of the  $P_j$ 's at  $u$  is the  $m$ -tuple  $(\varepsilon_1, \dots, \varepsilon_m) \in (-1, 0, 1)^m$ , where  $\varepsilon_j = \text{sign } P_j(u)$ . Let  $s(P_1, P_2, \dots, P_m)$  denote the total number of sign patterns of the polynomials  $P_1, P_2, \dots, P_m$ , as  $u$  ranges over all points of  $\mathbb{R}^\ell$ .

The following result is derived in [Alo95] as a slight modification of a theorem of Warren [War68].

**Theorem 9** *Let  $P_1 \dots P_m$  be  $m$  real polynomials in  $\ell$  real variables, and suppose the degree of each  $P_j$  does not exceed  $k$ . If  $2m \geq \ell$ , then  $s(P_1 \dots P_m) \leq (8ekm/\ell)^\ell$ .*

The following statement follows from a result of Erdős and Sachs [ES63], and can be also proved directly by a simple probabilistic argument.

**Lemma 7** *For every  $g \geq 3$  and every  $n \geq 3$ , there are (connected) graphs on  $n$  vertices with at least  $\frac{1}{4}n^{1+1/g}$  edges, and with no cycle of length at most  $g$ .*

We note that there are slightly better known results based on certain algebraic constructions, but for our purpose here the above estimate suffices.

We can now prove Theorem 8. Throughout the proof and the rest of the section, we assume that  $n$  is large, whenever this is needed, and omit all floor and ceiling signs whenever these are not crucial.

**Proof of Theorem 8:** Without trying to optimize the constants, define  $g = \frac{\log n}{\log d + \log \log n + 5}$ . We will show that some  $n$ -point metric spaces require relaxation at least  $g - 1$ . Without loss of generality, assume  $g - 1$  is bigger than 1, as otherwise there is nothing to prove. By Lemma 7, there is a graph  $G = (V, E)$  on a set  $V = \{1, 2, \dots, n\}$  of  $n$  labeled vertices, with  $m \geq \frac{1}{4}n^{1+1/g} > 7nd \log n$  edges, and with no cycles of length at most  $g$ . For every subset  $E' \subset E$  of precisely  $m/2$  edges, the subgraph  $(V, E')$  of  $G$  defines a metric space  $T(E')$  on the set  $V$  (if the subgraph is disconnected, some distances can be defined to be infinite; alternatively, we can fix a spanning tree in  $G$  and include it in all subgraphs to make sure they are all connected). This gives us a collection of  $2^{(1+o(1))m}$  metric spaces on  $V$ , with the following property.

(\*) For every two distinct spaces  $(T, d)$  and  $(T', d')$  in the collection, there are two pairs of points  $x, y$  and  $z, w$  so that  $d(x, y) = 1$  and  $d'(x, y) > g - 1$ , whereas  $d'(z, w) = 1$  and  $d(z, w) > g - 1$ .

Indeed, this follows from the fact that, for every two distinct subgraphs in our collection, there is an edge  $\{x, y\}$  belonging to the first one and not to the second, and vice versa. As the shortest cycle in  $G$  is of length exceeding  $g$ , the claim in (\*) follows.

Fix a space  $T$  in our collection, and let  $\phi_T$  be a minimum relaxation embedding of it into  $d$ -dimensional Euclidean space. Let  $\phi_T(i) = (x_{i,1}^T, \dots, x_{i,d}^T)$ . Then the square of the Euclidean distance between each two points in the embedding can be expressed as a polynomial of degree 2 in the  $dn$  variables  $x_{i,j}^T$ . The difference between two such squares of distances is thus also a polynomial of degree 2 in these variables. It follows that the order of all  $\binom{n}{2}$  distances is determined by the

signs of  $\binom{n}{2}^2 < n^4/4$  polynomials of degree 2 each, in  $dn$  variables. By Theorem 9, the total number of such orders is at most

$$\left(\frac{16en^4}{4dn}\right)^{dn} \leq n^{(3+o(1))dn} = 2^{(3+o(1))nd \log n}.$$

This is smaller than the number of spaces in our collection, and hence, by the pigeonhole principle, there are two distinct spaces  $T$  and  $T'$  in our collection, so that the orders of the distances in their embeddings are the same. This, together with (\*), implies that the relaxation in at least one of these embeddings is at least  $g - 1$ , completing the proof.  $\square$

The last proof easily extends to embeddings into  $d$ -dimensional  $\ell_p$  space for any even integer  $p$ . The only difference is that, in this case, the  $p$ th power of the distance between a pair of given points in the embedding is a polynomial of degree  $p$  in the  $dn$  variables describing the embedding. Working out the computation in the proof above, this yields the following result.

**Theorem 10** *There is an absolute constant  $c > 0$  such that, for every  $d$  and  $n$ , and for every even integer  $p$ , there is a metric space  $T$  on  $n$  points such that the relaxation in any ordinal embedding of  $T$  into  $d$ -dimensional  $\ell_p$  space is at least  $\frac{\log n}{\log d + \log(\log n + \log p) + c} - 1$ .*

The above argument, combined with an additional trick, can in fact be extended to handle ordinal embeddings into  $d$ -dimensional  $\ell_p$  space for odd integers  $p$ , as well as embeddings into  $d$ -dimensional  $\ell_\infty$  space.

**Theorem 11** (i) *For every  $n \geq d$ , there is a metric space  $T$  on  $n$  points such that the relaxation in any ordinal embedding of  $T$  in  $d$ -dimensional  $\ell_\infty$  space is at least  $\frac{\log n}{\log d + \log \log n + O(1)} - 1$ .*

(ii) *For every  $n \geq d$ , and for every odd positive integer  $p$ , there is a metric space  $T$  on  $n$  points such that the relaxation of any ordinal embedding of  $T$  into  $d$ -dimensional  $\ell_p$  space is at least  $\frac{\log n}{\log(2d^2 + 3d \log n + d \log p + O(d))} - 1$ .*

**Proof:** As before, the result is proved by a counting argument: we prove that the number of possible orders between all distances in a set of  $n$  points in the relevant spaces is not too large, and use the fact that there are many significantly different metric spaces on  $n$  points, concluding that for two such metric spaces the embedding orders the distances identically, and hence deriving the required lower bound on relaxation.

(i) We start by bounding the number of possible orders of all distances in a set  $X$  of  $n$  points in  $d$ -dimensional  $\ell_\infty$  space. Given such a set, define, for each ordered set  $(x, y, z, w)$  of (not necessarily distinct) four points of  $X$ , and for each two indices  $i, j$  in  $\{1, 2, \dots, d\}$ , the following linear polynomial in the  $dn$  variables representing the coordinates of the points:  $(x_i - y_i) - (w_j - z_j)$ . By Theorem 9 these  $d^2 n^4$  polynomials have at most  $(O(1)dn^3)^{dn} \leq 2^{(4+o(1))dn \log n}$  sign patterns. (In fact, because the polynomials here are linear, there is a slightly better, and simpler, estimate than the one provided by Warren's Theorem here—see [Har67]—but the asymptotic of the logarithm in this estimate is the same.) We claim that the signs of all these polynomials determine completely the order on all the  $\binom{n}{2}$  distances between pairs of the points. Indeed, the signs of the polynomials  $(x_i - y_i) - (x_j - y_j)$ ,  $(x_i - y_i) - (y_j - x_j)$  (and their inverses) determines a coordinate  $i$  such that  $\|x - y\|_\infty$  is  $x_i - y_i$  or  $y_i - x_i$  (as this is simply the maximum of all  $2d$  differences of the form  $(x_i - y_i), (y_i - x_i)$ ). Suppose, now, that  $\|x - y\|_\infty = x_i - y_i$  and  $\|w - z\|_\infty = w_j - z_j$ . Then the sign of  $(x_i - y_i) - (w_j - z_j)$  determines which of the two distances is bigger. It follows that the total number of orders of the distances of  $n$  points in  $d$ -dimensional  $\ell_\infty$  space is at most  $2^{(4+o(1))dn \log n}$ .

Define  $g = \frac{\log n}{\log d + \log \log n + 5}$ , take a graph  $G = (V, E)$  as in the proof of Theorem 8, and construct a collection of  $2^{(1+o(1))7nd \log n}$  metric spaces on a set of  $n$  points satisfying (\*). The desired result follows, just as in the proof of Theorem 8.

(ii) As in the proof of part (i), we first bound the number of possible orders of all distances in a set  $X$  of  $n$  points in  $d$ -dimensional  $\ell_p$  space. Given such a set, define, for each two (not necessarily distinct) pairs  $\{x, y\}$  and  $\{z, w\}$  of points, and each two sign vectors

$$(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_d), (\delta_1, \delta_2, \dots, \delta_d) \in \{-1, 1\}^d,$$

the following polynomial in the  $dn$  coordinates of the points:

$$\sum_{i=1}^d \varepsilon_i (x_i - y_i)^p - \sum_{j=1}^d \delta_j (z_j - w_j)^p.$$

This is a set of  $2^{2d}n^4$  polynomials, each of degree  $p$ , and thus, by Theorem 9, the number of their sign patterns is bounded by

$$2^{2d^2n + 3dn \log n + dn \log p + O(dn)}. \quad (4)$$

As before, it is not difficult to see that the signs of all these polynomials determine completely the order of all distances between pairs of points. Therefore, the number of such orders does not exceed (4). The desired result now follows as before, by considering metrics induced by subgraphs with half the edges of a graph on  $n$  vertices with at least  $\frac{1}{4}n^{1+1/g}$  edges, and no cycles of length at most  $g$ , where  $g = \frac{\log n}{\log(2d^2 + 3d \log n + d \log p + O(d))}$ .  $\square$

## 9 Conclusion and Open Problems

We have introduced minimum-relaxation ordinal embeddings and shown that they have distinct and sometimes surprising behavior. Yet many problems remain to be explored in this context; our hope is that this paper forms the foundation of a fruitful body of research. Here we highlight some of the more important directions for future exploration.

An important line of study is to continue comparing ordinal embeddings with metric embeddings. One interesting question is whether the dimensionality-reduction results of Bourgain [Bou85] and Johnson and Lindenstrauss [JL84] can be improved for ordinal relaxation. From Theorem 8 and Proposition 1, we know that the optimal worst-case relaxation for an ordinal embedding of a general metric into  $O(\lg n)$ -dimensional Euclidean space is between  $\Omega(\lg n / \lg \lg n)$  and  $O(\lg n)$ . Closing this  $\Theta(\lg \lg n)$  gap is an intriguing open problem; a better upper bound would improve on Bourgain-based metric embeddings into  $O(\lg n)$  dimensions. Another problem is how much relaxation is required for dimensionality *reduction* of a metric already embedded in arbitrary dimensional  $\ell_p$  space. For  $p = 2$ , we obtain an ideal relaxation of  $1 + \varepsilon$  using Johnson-Lindenstrauss combined with Proposition 1. For  $p = \infty$ , dimensionality reduction is impossible, by Theorem 11(i), because  $\ell_\infty$  is universal in the metric sense. For  $p \neq 2, \infty$ , the problem is open; in contrast, it is known for metric embeddings that dimensionality reduction is impossible for  $\ell_1$  [BC03, LN04].

Another important direction is to develop more approximation algorithms for minimum-relaxation ordinal embedding. Unlike general upper bounds on distortion, existing approximation algorithms for minimum-distortion metric embedding do not carry over to ordinal embedding because the optimum solution is generally smaller. Our  $O(1)$ -approximation result in Section 5, and the lack of a

matching result for metric embedding despite much effort, shows that in some contexts ordinal embedding problems may prove more easily approximable than metric embedding. We expect that our approximation result can be generalized using similar techniques to unweighted graphs, weighted trees, and/or higher dimensions, and that it can be strengthened to a PTAS. A related open problem is to consider trees as target metrics, and find the tree metric into which a given metric can be ordinally embedded with approximately minimum relaxation. Another family of approximation problems arise with the related notion of *additive relaxation*, in contrast to (multiplicative) relaxation, where pairs of distances within an additive  $\alpha$  must have their relative order preserved. In some cases, approximation results may be harder for ordinal embedding than metric embedding. For example, in the problem of approximating the minimum additive distortion/relaxation for an ordinal embedding of an arbitrary metric into the line, the simple greedy algorithm of Proposition 5 is a 3-approximation for metric embedding but can be arbitrarily bad for ordinal embedding.<sup>3</sup>

A final direction to consider is finding other applications of ordinal embedding. In particular, in the context of approximation algorithms for other problems, when are low-relaxation ordinal embeddings as useful as (and more powerful than) low-distortion metric embeddings? Nearest neighbor is a simple example where only the order of distances is relevant, but we expect there are several other such problems.

## Acknowledgments

We thank Amir Ben-Dor, Piotr Indyk, Howard Karloff, and Yuval Rabani for helpful discussions. In particular, Amir Ben-Dor contributed key ideas to the proof of universality (Theorem 1). Part of this work was done during a visit of the first, third, and fifth authors at Microsoft Research in Redmond, WA; we thank our hosts at Microsoft for their hospitality.

## References

- [ABF<sup>+</sup>99] Richa Agarwala, Vineet Bafna, Martin Farach, Mike Paterson, and Mikkel Thorup. On the approximability of numerical taxonomy (fitting distances by tree metrics). *SIAM J. Comput.*, 28(3):1073–1085, 1999.
- [AFR85] Noga Alon, Peter Frankl, and Vojtech Rödl. Geometrical realization of set systems and probabilistic communication complexity. In *Proceedings of the 26th Annual Symposium on Foundations of Computer Science*, pages 277–280, Portland, Oregon, 1985.
- [Alo95] Noga Alon. Tools from higher algebra. In R. L. Graham, M. Grötschel, and L. Lovász, editors, *Handbook of combinatorics*, volume 2, chapter 32, pages 1749–1783. MIT Press, 1995.
- [BC03] Bo Brinkman and Moses Charikar. On the impossibility of dimension reduction in  $\ell_1$ . In *Proceedings of the 44th Symposium on Foundations of Computer Science*, pages 514–523, 2003.

---

<sup>3</sup>The example is as follows. The graph has four points  $a, b, p, q$  and  $D(p, q) = 50$ ,  $D(p, a) = 100$ ,  $D(p, b) = 100 + \varepsilon$ ,  $D[q, a] = 70$ ,  $D(q, b) = 60$ , and  $D(a, b) = 10$ . The optimum ordinal embedding can place the points in the order  $p, q, b, a$ , and has additive relaxation  $\varepsilon$ , while greedy places the points in the order  $p, q, a, b$  and has additive relaxation 10, resulting in an arbitrarily large ratio.

- [BDG<sup>+</sup>05] Mihai Bădoiu, Kedar Dhamdhere, Anupam Gupta, Yuri Rabinovich, Harald Raecke, R. Ravi, and Anastasios Sidiropoulos. Approximation algorithms for low-distortion embeddings into low-dimensional spaces. In *Proceedings of the 16th Annual ACM-SIAM Symposium on Discrete Algorithms*, Vancouver, British Columbia, Canada, January 2005.
- [BDHI04] Mihai Bădoiu, Erik D. Demaine, MohammadTaghi Hajiaghayi, and Piotr Indyk. Low-dimensional embedding with extra information. In *Proceedings of the 20th Annual ACM Symposium on Computational Geometry*, Brooklyn, New York, June 2004.
- [BL04] Y. Bilu and N. Linial. Monotone maps, sphericity and bounded second eigenvalue. arXiv:math.CO/0401293, January 2004.
- [BM04] Y. Bartal and M. Mendel. Dimension reduction for ultrametrics. In *Proceedings of the 15th ACM-SIAM Symp. on Discrete Algorithms*, pages 664–665, 2004.
- [BMMV03] Robert Babilon, Jiří Matoušek, Jana Maxová, and Pavel Valtr. Low-distortion embeddings of trees. *Journal of Graph Algorithms and Applications*, 7(4):399–409, 2003.
- [Bou85] J. Bourgain. On Lipschitz embedding of finite metric spaces in Hilbert space. *Israel J. Math.*, 52(1-2):46–52, 1985.
- [Bǒ3] M. Bădoiu. Approximation algorithm for embedding metrics into a two-dimensional space. In *Proceedings of the 14th Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 434–443, 2003.
- [CS74] J. P. Cunningham and R. N. Shepard. Monotone mapping of similarities into a general metric space. *Journal of Mathematical Psychology*, 11:335–364, 1974.
- [CS98] B. Chor and M. Sudan. A geometric approach to betweenness. *SIAM Journal on Discrete Mathematics*, 11(4):511–523, 1998.
- [ES63] Paul Erdős and Horst Sachs. Reguläre Graphen gegebener Tailenweite mit minimaler Knotenzahl. *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe*, 12:251–257, 1963.
- [FK99] Martin Farach and Sampath Kannan. Efficient algorithms for inverting evolution. *J. ACM*, 46(4):437–449, 1999.
- [FKW95] M. Farach, S. Kannan, and T. Warnow. A robust model for finding optimal evolutionary trees. *Algorithmica*, 13(1-2):155–179, 1995.
- [Gup00] A. Gupta. Embedding tree metrics into low-dimensional Euclidean spaces. *Discrete Comput. Geom.*, 24(1):105–116, 2000.
- [Har67] E. F. Harding. The number of partitions of a set of  $N$  points in  $k$  dimensions induced by hyperplanes. *Proceedings of the Edinburgh Mathematical Society, Series II*, 15:285–289, 1966/1967.
- [HIL03] Johan Håstad, Lars Ivansson, and Jens Lagergren. Fitting points on the real line and its application to RH mapping. *Journal of Algorithms*, 49(62):42–62, 2003.

- [Hol72] W. Holman. The relation between hierarchical and euclidean models for psychological distances. *Psychometrika*, 37(4):417–423, 1972.
- [IM04] Piotr Indyk and Jiří Matoušek. Low-distortion embeddings of finite metric spaces. In J. E. Goodman and J. O’Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 8, pages 177–196. CRC Press, second edition, 2004.
- [Iva00] L. Ivansson. *Computational Aspects of Radiation Hybrid*. PhD thesis, Department of Numerical Analysis and Computer Science, Royal Institute of Technology, 2000.
- [JL84] W. B. Johnson and J. Lindenstrauss. Extensions of lipshitz mapping into hilbert space. *Contemporary Mathematics*, 26:189–206, 1984.
- [Kru64a] J. B. Kruskal. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*, 29:1–28, 1964.
- [Kru64b] J. B. Kruskal. Non-metric multidimensional scaling. *Psychometrika*, 29:115–129, 1964.
- [LLR95] Nathan Linial, Eran London, and Yuri Rabinovich. The geometry of graphs and some of its algorithmic applications. *Combinatorica*, 15(2):215–245, 1995.
- [LN04] James R. Lee and Assaf Naor. Embedding the diamond graph in  $L_p$  and dimension reduction in  $L_1$ . *Geometric and Functional Analysis*, 14(4):745–747, 2004.
- [Mat97] Jiří Matoušek. On embedding expanders into  $l_p$  spaces. *Israel Journal of Mathematics*, 102:189–197, 1997.
- [Opa79] J. Opatrny. Total ordering problem. *SIAM J. Computing*, 8:111–114, 1979.
- [SFC04] R. Shah and M. Farach-Colton. On the complexity of ordinal clustering. *Journal of Classification*, 2004. To appear.
- [She62] R. N. Shepard. Multidimensional scaling with unknown distance function I. *Psychometrika*, 27:125–140, 1962.
- [Tor52] W. S. Torgerson. Multidimensional scaling I: Theory and method. *Psychometrika*, 17(4):401–414, 1952.
- [War68] Hugh E. Warren. Lower bounds for approximation by nonlinear manifolds. *Transactions of the American Mathematical Society*, 133:167–178, 1968.