

Problem Set no. 1

Given: March 16, 2016

Due: March 30, 2016, Box 288

Exercise 1.1 Let $\mathbf{x} = (x_0, x_1, \dots, x_{n-1}) \in \mathbb{R}^n$ and let $\mathbf{y} = (y_0, y_1, \dots, y_{n-1}) \in \mathbb{C}^n$ be such that $\mathbf{y} = DFT(\mathbf{x})$. Prove that $y_{n-j} = y_j^*$, for $j = 1, \dots, n-1$. (Here z^* is the *conjugate* of $z \in \mathbb{C}$, i.e., if $z = a + ib$, where $a, b \in \mathbb{R}$, then $z^* = a - ib$.)

Exercise 1.2 Let $\mathbf{x} = (f(0), f(\frac{1}{32}), \dots, f(\frac{31}{32})) \in \mathbb{R}^{32}$, where $f(x) = \sin(9(2\pi x)) + 3\sin(2(2\pi x)) + 2\cos(2\pi x)$. Compute $DFT(\mathbf{x})$. (Hint: no complicated calculations are necessary. Use the relations $\cos x = \frac{1}{2}(e^{ix} + e^{-ix})$, $\sin x = \frac{1}{2i}(e^{ix} - e^{-ix})$, and the fact that Fourier basis is an orthonormal basis.)

Exercise 1.3 The *chirp transform* of a vector $(x_0, x_1, \dots, x_{n-1}) \in \mathbb{C}^n$, with respect to an arbitrary complex number $z \in \mathbb{C}$, is defined as follows: $y_k = \sum_{j=0}^{n-1} x_j z^{jk}$, for $k = 0, 1, \dots, n-1$.

(a) Show that the DFT is a special case of the chirp transform. (For which z ?)

(b) Use the relation $y_k = z^{k^2/2} \sum_{j=0}^{n-1} (x_j z^{j^2/2})(z^{-(k-j)^2/2})$ to express the chirp transform as a convolution. (Use caution. In the sum given, j ranges from 0 to $n-1$, for every value of k , while this is *not* the case for the non-cyclic convolution.)

(c) Show that the convolution of two vectors of length n can be computed in $O(n \log n)$ time for every value of n , not necessarily a power of 2.

(d) Use the previous results to show that the DFT of a vector of length n can be computed in $O(n \log n)$ time for any value of n .

Exercise 1.4 The *cross-correlation* of $\mathbf{x} = (x_0, x_1, \dots, x_{n-1})$ and $\mathbf{y} = (y_0, y_1, \dots, y_{m-1})$ is defined to be $\mathbf{z} = (z_{-(m-1)}, \dots, z_{n-1})$ such that $z_k = \sum_j x_{j+k} y_j$. (The sum here is over j such that $0 \leq j+k < n$ and $0 \leq j < m$.) Show that the *cross-correlation* of two vectors of lengths n and m respectively, where $m \leq n$, can be computed in $O(n \log m)$ time.

Exercise 1.5 Let T be a text of length n and let P be a pattern of length m over a finite and small alphabet Σ . Let D be a $|\Sigma| \times |\Sigma|$ matrix such that $D(a, b)$ specifies the *similarity* or *dissimilarity* of $a, b \in \Sigma$. For every $k = 0, 1, \dots, n-m-1$ define $d_k = \sum_{j=0}^{m-1} D(T[k+j], P[j])$ to be the total pattern-text dissimilarity when the 0-th pattern character is aligned with the k -th text character.

(a) Show that d_k , for $k = 0, 1, \dots, n-m-1$, can be computed in $O(|\Sigma| n \log m)$ time.

(b) Suppose now that $\Sigma \subset \mathbb{Z}$, i.e., that each character is actually an integer, and that $D(a, b) = ab$, for every $a, b \in \Sigma$. How fast can the d_k 's be computed?

(c) Suppose that we again have $\Sigma \subset \mathbb{Z}$ but this time $D(a, b) = (a-b)^2$, for every $a, b \in \Sigma$. How fast can the d_k 's be computed?